

Relationship Interleukin-8 levels, Malassezia, and sebum fatty acids in dandruff severity

Yosse Rizal¹, Nuzulia Irawati^{2*}, Irma D. Rosyanto³, Netti Suharti⁴

¹Program Doctor Biomedical Science, Faculty of Medicine, Universitas Andalas, Padang, Indonesia. ²Department of Parasitology, Faculty of Medicine, Universitas Andalas, Padang, Indonesia. ³Department of Dermatology and Venerology, Faculty of Medicine Universitas of Sumatera Utara, Medan, Indonesia. ⁴Department of Microbiology, Faculty of Medicine, Universitas Andalas, Padang, Indonesia.

Correspondence: Nuzulia Irawati, Department of Parasitology, Faculty of Medicine, Universitas Andalas, Padang, Indonesia. nuzulairawati@med.unand.ac.id

ABSTRACT

Dandruff is one of the skin diseases that cause problems for patients with complaints of itching. Apart from feeling itchy, it also causes discomfort for patients. This study aims to examine *Malassezia* species in patients with dandruff by PCR-RT. This is a cross-sectional observational study with a sample size of 60 dandruff patients. Patients were divided into mild/moderate and severe severity based on Severity Dandruff Score. *Malassezia* species examination was performed by real-time polymerase chain reaction sequencing. Lipid types were determined by gas chromatography-mass spectrometry. IL-8 levels were examined using the Elisa method. The results showed 26 patients with mild-moderate dandruff and 34 with severe dandruff. *M.restricta* topped the list with 73.3%, followed by *M.globosa* (23.3%). The most common sebaceous lipid type was hexanediol acid (41.6%), followed by octadecanoic acid (31.7%). IL-8 levels were associated with the severity of dandruff, where in mild-moderate degree, there was an association of $p=0.03$, while in severe degree, there was a highly relevant association ($p=0.00001$). This study shows that elevated IL-8 levels are associated with the severity of dandruff. *M.restricta* is the most common cause of dandruff, while hexanedioic acid is the most common fatty acid that causes dandruff.

Keywords: Dandruff, GC-MS, Interleukin-8, *Malassezia*, RT-PCR

Introduction

Dandruff (*Pityriasis capitis*) is a problem in dermatology and venerology. It causes itching, the most common complaint felt by dandruff patients. Apart from causing itching, dandruff also causes psychological and aesthetic disorders and affects the patient's social quality of life. It is part of seborrheic dermatitis, but the two diseases differ in location and severity [1]. More than 50% of adults worldwide have suffered from dandruff in their lifetime [2]. The severity and clinical degree of dandruff sustained by a person varies. Men suffer from dandruff more than women, with a ratio of 3:2 [3]. *Malassezia fungus* is one of the

microorganisms that cause dandruff, but the virulence factor of *Malassezia* is not known with certainty [4, 5]. *Malassezia* is a normal fungus on healthy human skin spread over several body parts [3]. From several studies conducted on cases of seborrheic dermatitis, *Malassezia restricta*, and *Malassezia globosa* are the two most common fungal species found and play the most role in the occurrence of the disease [6, 7]. Using the Polymerase Chain Reaction identification technique, *Malassezia sympodialis* and *Malassezia sloffiae* were most commonly found in these cases [8]. Apart from causing dandruff and seborrheic dermatitis, *Malassezia* species also cause skin diseases such as *Malassezia folliculitis* and *Pityriasis Versicolor*.

Differences in lipid levels on the body's surface also affect the type of *Malassezia* species. Most *Malassezia* have a dominant preference for depending on exogenous sources of fatty acids for synthesizing fatty acids and metabolizing carbohydrates [9]. *Malassezia restricta* is more commonly found on the forehead and scalp, so this fungus is most often found in cases of dandruff and seborrheic dermatitis [6], while *Malassezia globosa* is more

Access this article online

Website: www.japer.in

E-ISSN: 2249-3379

How to cite this article: Rizal Y, Irawati N, Rosyanto ID, Suharti N. Relationship Interleukin-8 levels, *Malassezia*, and sebum fatty acids in dandruff severity. J Adv Pharm Educ Res. 2025;15(1):54-9. <https://doi.org/10.51847/TdgjvOwMq>

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

dominant on the chest and back, so it is often found in cases of *Pityriasis versicolor* [10]. IgG and IgM antibodies against *Malassezia* are detectable in most individuals, but healthy individuals are usually not as sensitive as atopic dermatitis patients [8]. Dandruff is often associated with the onset of puberty and concurrent with increased activity of the sebaceous glands of the scalp because all *Malassezia* species except *M. pachydermatis* require exogenous lipids as nutrition; they are often associated with sebum-rich areas of the skin, such as the scalp [11]. The interaction between *Malassezia* fungi, sebaceous gland lipids, and a person's immune mechanism will determine the change in the form of *Malassezia*, which was initially commensal to becoming a pathogen.

Malassezia can induce the production of proinflammatory cytokines released by keratinocyte cells¹. If sebum production decreases, the lipid layer in the stratum corneum is affected, and the protective function of the lipid barrier may be disrupted [12]. Sebum nourishes *Malassezia* fungi, generally found in human sebum-rich areas, whose multiplication produces more fatty acids that disrupt the scalp's integrity and reduce the skin barrier [13]. Lipid metabolites (linoleic acid), lipase enzymes, and phospholipase enzymes induce skin inflammation and irritation. Lipase and phospholipase enzymes are considered the central virulence of *Malassezia* fungi [14]. The lipase enzymes produced by *Malassezia globosa* have significantly increased activity in the skin. The patient's head has been proven in patients suffering from seborrheic dermatitis [15]. Previous studies reported the expression of the lipase gene *Malassezia restricta* (MRE-0242). The gene is like LIP-1 in *Malassezia globosa* [16]. Increased lipase and phospholipase secretion of *Malassezia* species in the skin can produce lipid metabolites in the form of indole compounds, which can damage the epidermis and activate local immune responses in seborrheic dermatitis and pityriasis versicolor [17]. Still, there is no data on these lipids' effect on dandruff.

The sensitization of *Malassezia* to human skin in activating the immune response can occur directly and indirectly. *Malassezia* stimulates an immunological response in the skin by activating dendritic cells and macrophages. Keratinocyte cells exposed to *Malassezia* produce cytokines and chemokines that play a role in inflammation and the severity of dandruff. Interleukin 8 (IL-8) produced by human keratinocyte cells in cases of seborrheic dermatitis exposed to *Malassezia* increased significantly compared to healthy people. In addition, keratinocytes also produce several other proinflammatory cytokines, such as IL-1, IL-6, IL-12, and TNF α [18, 19].

Intracellular signal recognition of human keratinocyte cells in response to exposure to *Malassezia* can be through signaling Toll-like receptor 2 (TLR2), C type Lectin receptor (CLR), dectin and minge, which belong to the glycoprotein group, which functions as a transmembrane surface receptor/Pattern Recognise Receptor (PRR) [20]. Toll-like receptors and CLRs are involved in the natural immune response against microorganisms. Toll-like receptors and C receptor lectins initiate signaling pathways that activate cytokines, chemokines, and antimicrobial peptides. Keratinocytes can express TLR1, TLR2, TLR3 and TLR5. Toll-like receptor two can recognize simple molecules (ligands) in fungi, namely zymosan [21, 22].

At this time, the species of *Malassezia* fungus, the levels of Interleukin-8, and the type of lipid that causes the severity of dandruff have not been identified. Therefore, this study was conducted to analyze the relationship between knowing levels of interleukin-8 and the severity of dandruff. This research is vital because dandruff treatment can be carried out properly by obtaining a relationship between these variables.

Materials and Methods

Samples collection

Samples were taken from dandruff patients who went to the Dermatology and Venereology Polyclinic at the Ahmad Muchtar Hospital, Bukittinggi, by scraping scales from the patient's scalp using a scalpel and a spirit comb. The samples were then inserted and stored in a 1.5 ml microtube until they were used.

Sampling

Samples were collected by scraping scales or scales from the patient's scalp using a scalpel and a spirit comb, then inserted and stored in a 1.5 ml microtube until use. Each sample will be separated into three microtubes, which will be used to identify *Malassezia* species, measure Interleukin-8 levels, and determine sebaceous lipid types.

Identification spesies malassezia

Dandruff samples were tested using the Alkaline Lysis method. The dandruff samples were centrifuged in PBS for 2 minutes. After removing the supernatant, 50 mM NaOH 180 μ l was added to the dandruff pellets and vortexed. They were incubated at 95°C for 10 minutes. Add 20 μ l Tris-HCl Ph 8.0, then vortex, centrifuge at 12,000 rpm for 5 minutes. Next, take the 0.5-2 μ l supernatant for PCR and 100 μ l transfer it to a new microtube. DNA isolation was carried out according to the kit procedure, which generally consisted of sample preparation, cell lysis, DNA binding, washing, and DNA elution. DNA extraction with enzyme polymerase using KOD Toyobo Fx Neo Japan Bio-Technology reagent.

The amplification steps were carried out as follows: 95°C for 14 minutes (predenaturation) followed by 40 cycles at 94 °C for 45 seconds (denaturation), 55 °C for 45 seconds (annealing), 72 °C for 1-minute extension and 72 °C for 7 minutes final extension. The 580-bp chain associated with the 26S rDNA was electrophoresed with agarose gel, then restriction was carried out for 5 hours, and finally, the DNA fragments were visualized by electrophoresis in 1.5% agarose gel.

Measurement of interleukin-8 levels

The levels of interleukin-8 samples were tested using the Human IL-8 ELISA kit. Each sample was tested by calculating its absorption capacity at a wavelength of 450 nm using a multi-mode Microplate reader (Biorad). The results obtained are expressed in units of ng/l.

Lipid type testing

The type of sebaceous lipid was tested using the GCMS analysis. At the preparation stage, the sample was mixed with 2 ml of aqua and then bound using chloroform and methanol with a ratio of 1:2. Vortex for 2 minutes, add 2 ml of chloroform and 2.5 ml of aqua, vortex for 30 seconds, and centrifuge for 6 minutes. After that, the precise phase was discarded. It was put into the GCMS tool. The remaining extract was added to 2.5 ml of chloroform and incubated in the oven at 90 °C for 120 minutes. Added 1 ml of distilled water and 2 ml of extraction solution, then extracted with 2 ml of a mixture of hexa and chloroform with a ratio of 4:1. The top layer of this solution was taken with repetition up to 3 times. The results are read as a peaks diagram chromatogram GCMS solution [23-29].

Statistical analysis

Before carrying out the bivariate test, a normality test is carried out for numerical data. If the data is normally distributed, a parametric test can be carried out; if it is not normally distributed, then a transformation is carried out. After being transformed, the normality test was carried out again. A parametric test is carried out if it is usually distributed, but if it is still not normal, a non-parametric test is carried out. The Kruskal-Wallis and Mann-Whitney tests were carried out to see the relationship between variables.

Ethical clearance

This research has received ethical consideration and approval from the Research Ethics Committee Team of the Faculty of Medicine, Universitas Andalas, with registration number 109/KEP/FK/2020.

Results and Discussion

Identification of *Malassezia* species in dandruff patients

The type of *Malassezia* found in the examination above can be seen in **Table 1**.

Table 1. Results of the identification of *Malassezia* as a cause of dandruff in research subjects

No	Type	f	%
1	<i>Malassezia globosa</i>	14	23.3
2	<i>Malassezia restricta</i>	44	73.3
3	<i>Malassezia furfur</i>	1	1.67
4	<i>Malassezia verfertilionis</i>	1	1.67
Total		60	100

Table 1 identified *Malassezia restricta* as the most common cause of dandruff in research subjects (73.3%).

In this study, it was found that the most common fungus, *Malassezia restricta*, was found, namely 44 (73.33%), followed by

Malassezia globosa 14 (23.33%). *Malassezia vespertilionis* was also found, which is unusual on the scalp and has never been reported as a cause of dandruff. *Malassezia vespertilionis* is a species of yeast-like fungus that lives on the skin of bats. It was described as a new species in 2018. The holotype was obtained from skin swabs of the long-eared bat (*Myotis septentrionalis*) hibernating in Wisconsin, USA. The name *vespertilionis* uses the Latin *vespertilio* (bat) to refer to its host [30].

The results of this study are research conducted, who conducted a survey to identify *Malassezia* species in 21 dandruff patients using the PCR method. The results showed that there were ten species of *Malassezia*, with the most species being *M. sympodialis* (23.8%), *M. slooffiae* (9.5%), *M. furfur* (4.8%), *M. globosa* (4.8%), *M. restricta* and *M. sympodialis* (4.8%) [31]. In this research, no *Malassezia vespertilionis* species were found in all patients.

Sebaceous lipid types in dandruff patients

The results of research on sebaceous lipid types in research subjects can be seen in **Table 2**.

Table 2. Results of research on sebaceous lipid types in research subjects

No	Lipid Type	f	%
1	Octa decenoic acid	19	31.7
2	Benzenedicarboxylic acid	3	5
3	Hexa dioic acid	25	41.6
4	Hexa decanoic acid	7	11.7
5	Penta decanoid acid	1	1.7
6	Hepta decanoid acid	4	6.6
7	Tri decanoid acid	1	1.7
Total		60	100

Table 2 shows that Hexa nedioic acid is the most common type of lipid (41.6%), followed by octadecenoic acid (31.7%) in dandruff patient study subjects.

The results of this study are also from a survey conducted by Honnavar et al. [32] in India on 200 patients with seborrheic dermatitis on the head. The results showed that *M. globosa* was found to be dominant at 36.2%, followed by *M. restricta* at 31.3%, *M. furfur* at 15.7%, and a mixture of *M. globosa* and *M. restricta* at 12% or *M. arunalokei* at 4.8%. Researchers [33] also conducted a study to identify *Malassezia* species in 57 dandruff patients, and the results showed that *M. sympodialis* (62.96%) was the most common *Malassezia* species, followed by *M. furfur* (25.93%) and *M. globosa* (11.11%).

Study to identify *Malassezia* species that can cause dandruff. This research was conducted on 145 patients with dandruff; then, samples were taken by scraping the scalp and cultured. Of the 145 patients, 90.3% showed positive results for *Malassezia* species, including *M. furfur* (70.2%) and *M. globosa* (51.9%) [34]. Revealed that *M. restricta* was the most common dandruff-causing species in North India at 37.8% and *M. furfur* at 46.4% in South India [35]. Using the PCR technique, Zaree et al. conducted a study to identify *Malassezia* species in 65 dandruff patients. The

results showed that 58.4% of *M. restricta* was predominantly on the scalp and 41.6% on *M. globosa* [36].

Dandruff severity degree

According to Rogers *et al.*, 2003, the severity of dandruff is based on clinical symptoms. Based on this, the severity is divided into mild-moderate and severe.

1. Mild-moderate with a value of 0 to small with 4
2. Severe with a value of 4-5

The study results of dandruff's severity can be seen in **Table 3**.

No	Degree of Severity	f	%
1	Mild-moderate	26	43
2	Severe	34	57
Total		60	100

Table 3 shows the severity of dandruff, mild-moderate (43%) and severe (57%).

Distribution of lipid type examination

In this study, the most common type of lipid, hexanedioic acid (adipic acid), was found in 25 samples (41.6%). Hexanedioic acid is a dicarboxylic acid, one of the components that forms dandruff. Until now, there has been no relationship between the type of lipid and the degree of severity of dandruff a person suffers. Analyze the fat on the scalp of patients with dandruff (10 people) and patients without dandruff (10 people) or controls. This study used gas chromatography coupled with mass spectrometry. The results showed that Squalene, the main sebum component, was significantly more oxidized in dandruff-affected scalps and resulted in a much higher ratio of squalene monohydroperoxide (SQOOH). This was observed when comparing the dandruff-affected areas of dandruff patients with unaffected areas and control patients [11].

Previous research to detect lipid production by Malassezia using Nile red stain. The results showed that the average lipase synthesis rate increased by 30.7% and 29.8% in cells that were given *M. globosa* and *M. restricta* protein extracts, and *M. sympodialis* could increase lipase synthesis by 13.6% [7]. The results of another study by Lee *et al.* also showed that *M. restricta* could increase lipase synthesis. This study was conducted by taking swabs from the scalp and cheeks of 18 patients with seborrheic dermatitis and then analyzing them using the RT-PCR technique; the results showed that *M. restricta* increased lipase and phospholipase levels [14].

Correlation between IL-8 levels and the severity of dandruff in research subjects

The relationship between IL-8 levels and the severity of dandruff in research subjects can be seen in **Table 4**.

No	Degree of Severity	F	IL-8 Levels	p
1	Mild-moderate	26	249.53±123.34	0.36
2	Severe	34	482.80±71.16	0.0001

Table 4 shows that in patients with mild-moderate degrees, there is no significant relationship with IL-8 levels ($p=0.36$); in patients with severe dandruff, there is a significant relationship with IL-8 levels ($p=0.0001$).

The relationship between interleukin 8 levels and the severity of dandruff

From the results of the ELISA examination, it was found that 26 samples had IL-8 levels below 400 ng/l, and 34 samples had IL-8 levels above 400 ng/l. IL-8 levels above 400 ng/l are categorized as high and vice versa. The average IL-8 level in severe dandruff was 484.27 ng/l, with a standard deviation of 70.64, while the IL-8 level in mild dandruff was 249.88, with a standard deviation of 123.76. The statistical test results obtained $p = 0.05$, meaning there was a significant difference in the average IL-8 level between mild and severe degrees of dandruff. The results of this study are also from a survey conducted by Honnavar *et al.* in India on 200 patients with seborrheic dermatitis on the head. The results showed that *M. globosa* was found to be dominant at 36.2%, followed by *M. restricta* at 31.3%, *M. furfur* at 15.7%, and mixed *M. globosa* dan *M. restricta* 12% or *M. arunalokei* at 4.8% [31]. Researchers [32] also conducted a study to identify Malassezia species in 57 dandruff patients, and the results showed that *M. sympodialis* (62.96%) was the most common Malassezia species, followed by *M. furfur* (25.93%) and *M. globosa* (11.11%).

The results of this study are based on research conducted, which showed that the concentration of IL-8 increased in all cultures treated with Malassezia species extracts. Cells treated with *M. globosa* extract showed the most significant increase, followed by *M. restricta*. In *M. globosa*, the concentration of IL-8 increased almost threefold, whereas in *M. restricta*, it caused a 2.77-fold increase. IL-8 production rate induced by *M. sympodialis*, *M. dermatis*, and *M. slooffiae* also increased significantly, respectively 2.58 times, 2.31 times, and 2.38 times compared to control [7]. This research is also supported by the results of previous research, which conducted a study to identify cytokine production by keratinocytes that Malassezia modulates. This study was undertaken to assess pro-inflammatory (IL-6, IL-8, IL-1a, and Tumor Necrosis Factor- α /TNF- α) and anti-inflammatory (IL-10) cytokines using enzyme-linked immunosorbent assays (ELISA) techniques [21]. The results showed that *M. globosa* acapsular caused a 66-fold increase in IL-8 production ($P < 0.001$), and *M. furfura* capsular caused a 38-fold increase in IL-6 production ($P < 0.001$), as well as an increase in IL-10 production. as much as 12 times.

Conclusion

From the study results, it can be concluded that *Malassezia restricta* is the most common species that causes dandruff, followed by *Malassezia globosa*. Hexa deconoid acid is the most common type of fat that causes dandruff, followed by Okta decenoid acid. There is a relationship between Interleukin-8 levels of dandruff severity, where in mild-moderate degrees, this relationship is not significant, but in severe degrees, this relationship is substantial. From this research, it is necessary to provide anti-inflammatory and anti-fungal drugs.

Acknowledgments: None

Conflict of interest: None

Financial support: None

Ethics statement: This research has received ethical consideration and approval from the Research Ethics Committee Team of the Faculty of Medicine, Universitas Andalas, with registration number 109/KEP/FK/2020.

References

- Schwartz JR, Messenger AG, Tosti A, Todd G, Hordinsky M, Hay RJ, et al. A comprehensive pathophysiology of dandruff and seborrheic dermatitis - towards a more precise definition of scalp health. *Acta Derm Venereol.* 2013;93(2):131-7. doi:10.2340/00015555-1382
- Grimshaw SG, Smith AM, Arnold DS, Xu E, Hoptroff M, Murphy B. The diversity and abundance of fungi and bacteria on the healthy and dandruff-affected the human scalp. *PLoS One.* 2019;14(12):e0225796. doi:10.1371/journal.pone.0225796
- Manuel F, Ranganathan S. A new postulate on two stages of dandruff: a clinical perspective. *Int J Trichology.* 2011;3(1):3-6. doi:10.4103/0974-7753.82117
- Turner GA, Hoptroff M, Harding CR. Stratum corneum dysfunction in dandruff. *Int J Cosmet Sci.* 2012;34(4):298-306. doi:10.1111/j.1468-2494.2012.00723.x
- Zhang H, Ran Y, Xie Z, Zhang R. Identification of *Malassezia* species in patients with seborrheic dermatitis in China. *Mycopathologia.* 2013;175(1-2):83-9. doi:10.1007/s11046-012-9606-z
- Hiruma M, Cho O, Hiruma M, Kurakado S, Sugita T, Ikeda S. Genotype analyses of human commensal scalp fungi, *Malassezia globosa*, and *Malassezia restricta* on the scalps of patients with dandruff and healthy subjects. *Mycopathologia.* 2014;177(5-6):263-9. doi:10.1007/s11046-014-9748-2
- Kim SY, Kim SH, Kim SN, Kim AR, Kim YR, Kim MJ, et al. Isolation and identification of *Malassezia* species from Chinese and Korean patients with seborrheic dermatitis and in vitro studies on their bioactivity on sebaceous lipids and IL-8 production. *Mycoses.* 2016;59(5):274-80. doi:10.1111/myc.12456
- Saunte DML, Gaitanis G, Hay RJ. *Malassezia*-associated skin diseases, the use of diagnostics and treatment. *Front Cell Infect Microbiol.* 2020;10:112. doi:10.3389/fcimb.2020.00112
- Park M, Cho YJ, Lee YW, Jung WH. Whole genome sequencing analysis of the cutaneous pathogenic yeast *Malassezia restricta* and identification of the major lipase expressed on the scalp of patients with dandruff. *Mycoses.* 2017;60(3):188-97. doi:10.1111/myc.12586
- Zhang E, Tanaka T, Tajima M, Tsuboi R, Nishikawa A, Sugita T. Characterization of the skin fungal microbiota in patients with atopic dermatitis and in healthy subjects. *Microbiol Immunol.* 2011;55(9):625-32. doi:10.1111/j.1348-0421.2011.00364.x
- Jourdain R, Moga A, Vingler P, El Rawadi C, Pouradier F, Souverain L, et al. Exploration of scalp surface lipids reveals squalene peroxide as a potential actor in dandruff condition. *Arch Dermatol Res.* 2016;308(3):153-63. doi:10.1007/s00403-016-1623-1
- Yoon JS, Shim J, Lim JM, Park SG. Biophysical characteristics of dandruff-affected scalp categorized on the basis of sebum levels. *J Cosmet Dermatol.* 2021;20(3):1002-8. doi:10.1111/jocd.13626
- Mondon P, Ringenbach C, Doridot E, Genet V. Reinforcement of barrier function and scalp homeostasis by Senkyunolide A to fight against dandruff. *Int J Cosmet Sci.* 2017;39(6):617-21. doi:10.1111/ics.12417
- Lee YW, Lee SY, Lee Y, Jung WH. Evaluation of expression of lipases and phospholipases of *malassezia restricta* in patients with seborrheic dermatitis. *Ann Dermatol.* 2013;25(3):310-4. doi:10.5021/ad.2013.25.3.310
- Dawson TL Jr. *Malassezia globosa* and *restricta*: breakthrough understanding of the etiology and treatment of dandruff and seborrheic dermatitis through whole-genome analysis. *J Investig Dermatol Symp Proc.* 2007;12(2):15-9. doi:10.1038/sj.jidsymp.5650049
- Fuchsberger C, Flannick J, Teslovich TM, Mahajan A, Agarwala V, Gaulton KJ, et al. The genetic architecture of type 2 diabetes. *Nature.* 2016;536(7614):41-7. doi:10.1038/nature18642
- Gaitanis G, Magiatis P, Hantschke M, Bassukas ID, Velegraki A. The *Malassezia* genus in skin and systemic diseases. *Clin Microbiol Rev.* 2012;25(1):106-41. doi:10.1128/CMR.00021-11
- Watanabe S, Kano R, Sato H, Nakamura Y, Hasegawa A. The effects of *Malassezia* yeasts on cytokine production by human keratinocytes. *J Invest Dermatol.* 2001;116(5):769-73. doi:10.1046/j.1523-1747.2001.01321.x
- Kerr K, Darcy T, Henry J, Mizoguchi H, Schwartz JR, Morrall S, et al. Epidermal changes associated with symptomatic resolution of dandruff: biomarkers of scalp

- health. *Int J Dermatol.* 2011;50(1):102-13. doi:10.1111/j.1365-4632.2010.04629.x
20. Baroni A, Orlando M, Donnarumma G, Farro P, Iovene MR, Tufano MA, et al. Toll-like receptor 2 (TLR2) mediates intracellular signaling in human keratinocytes in response to *Malassezia furfur*. *Arch Dermatol Res.* 2006;297(7):280-8. doi:10.1007/s00403-005-0594-4
21. Thomas DS, Ingham E, Bojar RA, Holland KT. In vitro modulation of human keratinocyte pro- and anti-inflammatory cytokine production by the capsule of *Malassezia* species. *FEMS Immunol Med Microbiol.* 2008;54(2):203-14. doi:10.1111/j.1574-695X.2008.00468.x
22. Ermertcan AT, Öztürk F, Gündüz K. Toll-like receptors and skin. *J Eur Acad Dermatol Venereol.* 2011;25(9):997-1006. doi:10.1111/j.1468-3083.2011.04049.x
23. Noor H, Coşte A, Micu A, Pirvut V. Cognitive disorders around surgery and its prevention strategies. *Arch Pharm Pract.* 2024;15(1-2024):33-9.
24. Akbari M, Moghadam RS, Mohamadpour A, Leili EK, Azaripour E. A randomized clinical trial on topical ketorolac administration following phaco-surgery for pseudoexfoliation syndrome. *Arch Pharm Pract.* 2024;15(1-2024):40-6.
25. El Hadad S, Alzahrani S, Alhebshi A, Alrahimi J. Impact of carrageenan-soy protein combination on CXCR-4 expression, cell viability, and apoptosis in HCT-116 cells. *Arch Pharm Pract.* 2024;15(1-2024):53-62.
26. Zeb-un-Nisa Z, Ali SI, Shahnaz S, Mumtaz T, Swaleh MM. Estimation of quality characteristics for sustained releasing and acting formulation of domperidone. *Pharmacophore.* 2021;12(1-2021):57-64.
27. Yousaf M, Khan MM, Paracha AT. Leading professionally diverse workgroups of healthcare professionals for improving quality of care. *J Organ Behav Res.* 2021;6(1-2021):106-19.
28. Kianimoghadam AS, Arani AM, Mohraz M, Bakhtiari M, Manshadi SA, Alinaghi SA. Psychometric properties of persian version of patient health questionnaire (PHQ-9) in Iranian HIV-infected patients. *J Organ Behav Res.* 2021;6(2):46-57.
29. Adjdir S, Benariba N, Adida H, Kamila G, El Hacı IA, Terki M, et al. Phenolic compounds and antimicrobial activity of *Ziziphus jujuba* Mill. fruit from Tlemcen. *J Biochem Technol.* 2021;12(1-2021):40-4.
30. Lorch JM, Palmer JM, Vanderwolf KJ, Schmidt KZ, Verant ML, Weller TJ, et al. *Malassezia vespertilionis* sp. nov.: a new cold-tolerant species of yeast isolated from bats. *Persoonia.* 2018;41:56-70. doi:10.3767/persoonia.2018.41.04
31. Jusuf NK, Nasution TA, Ulllyana S. Diagnostic value of nested-PCR for identification of *Malassezia* species in dandruff. *IOP Conf Ser Earth Environ Sci.* 2018;125:012050.
32. Honnavar P, Chakrabarti A, Dhaliwal M, Dogra S, Handa S, Lakshmi PV, et al. Sociodemographic characteristics and spectrum of *Malassezia* species in individuals with and without seborrhoeic dermatitis/dandruff: a comparison of residents of the urban and rural populations. *Med Mycol.* 2021;59(3):259-65. doi:10.1093/mmy/myaa050
33. Archana BR, Beena PM, Kumar S. Study of the Distribution of *Malassezia* species in patients with pityriasis versicolor in Kolar Region, Karnataka. *Indian J Dermatol.* 2015;60(3):321. doi:10.4103/0019-5154.156436
34. Biobelemonye N, Ogba MO, Bassey NL. Dandruff aetiology and the effects of edible lipids on the growth of Isolates. *Eur J Pharma Med Res.* 2016;3(9):71-6.
35. Rudramurthy SM, Honnavar P, Dogra S, Yegneswaran PP, Handa S, Chakrabarti A. Association of *Malassezia* species with dandruff. *Indian J Med Res.* 2014;139(3):431-7.
36. Zareei M, Mohammadi SR, Shahbazi S. Molecular identification of malassezia species with direct DNA extraction from scalp of patients with dandruff and seborrheic dermatitis. *J Pure Appl Microbiol.* 2015;9(4):2781.