

Investigating a baby food prepared from *Opuntia ficus indica* in Wistar rats

Maha Hanafy Mahmoud¹, Hany Mohamed Ahmed Wahba^{1*}, Marwa Hanafy Mahmoud²

¹Nutrition and Food Science Dept., National Research Centre, 33 EL Bohouth St., Dokki, Giza, Egypt, P.O.12622. ²Food Tech. Dept., National Research Centre, 33 EL Bohouth St., Dokki, Giza, Egypt, P.O.12622.

Correspondence: Hany Mohamed Ahmed Wahba, Nutrition and Food Science Dept., National Research Centre, Dokki, Giza, Egypt.
Email: hanywahba248@gmail.com

ABSTRACT

Vitamin A and calcium deficiencies are major health problems worldwide and represent one of the main reasons for malnutrition in childhood. *Opuntia* fruit is a wild plant rich in minerals and vitamins and can replenish vitamin A and calcium deficiency. The present study was aimed to prepare baby food from *Opuntia ficus indica* fruit after selecting a good drying technique that maintains the vitamin content of the dried fruit and to evaluate its biological impact in Wistar rats with either vitamin A depletion or calcium deficiency. Three drying methods were tested for the *Opuntia* fruit; the oven, the microwave, and solar energy. Sensory evaluation and proximate analysis were done for the prepared baby food from dried *Opuntia* fruit. Then, thirty-six male Wistar rats with a body weight of 70 ± 10 g were divided into 6 groups, 6 rats in each group. The groups were: a control negative group, a control group + 150 g dried *Opuntia* fruit, a group fed on calcium-deficient diet, a calcium-deficient diet group + 150 g dried *Opuntia* fruit, a group fed on vitamin A depleted diet, and a group fed on vitamin A depleted diet + 150 g dried *Opuntia* fruit. The experiment lasted for four weeks, then, serum vitamin A, serum ionized calcium, total calcium, serum urea, creatinine, albumin, total protein, and hemoglobin were determined. The results revealed very good acceptability of the prepared baby food, and the solar energy method was the best method of drying that gave the highest β -carotene content. Feeding rats with *Opuntia* fruit restored the serum calcium in the group, fed on a calcium-deficient diet and restored the serum's vitamin A in the group, fed on vitamin A depleted diet. Thus, it is recommended to safely introduce the prepared baby food for babies as it doesn't have any drawbacks on body weight gain or different body functions and it can protect babies from calcium and vitamin A deficiencies.

Keywords: *Opuntia ficus indica*, vitamin A deficiency, calcium deficiency, baby food, drying methods.

Introduction

Vitamin A deficiency and calcium deficiency are among the most frequent reasons for childhood malnutrition. Approximately 228 million children are affected by vitamin A deficiency. It is the most common form of malnutrition leading to blindness and co-morbidity of childhood worldwide. It was reported that 1 to 3 million child deaths are due to eye disease; 5 to 10 million cases among whom half a million children get

blinded every year. Vitamin A deficiency has many clinical manifestations, ranging from xerophthalmia (a spectrum of ocular disease arising from VAD) to susceptibility to infections and anaemia^[1]. The World Health Organization (WHO) recommends breastfeeding since breast milk is a good natural source for vitamin A, also it recommends vitamin A supplementation along with increasing the intake of vitamin A-rich diets and vitamin A fortified foods^[2].

Calcium deficiency represents a public health problem in developing countries and in many developed countries^[3]. Calcium is important for many biological processes, such as provision of structural support for teeth and bones, muscle contraction, enzyme regulation, signal transduction, blood clotting in addition to other important roles^[4]. Its deficiency leads to rickets in children and osteoporosis in adults^[3].

Many plants either edible or non-edible have been known and used for their medicinal values since thousands of years ago in traditional medicine. Recently, in the era of medicine, lights are spot on the natural compounds to replace some of the natural

Access this article online

Website: www.japer.in

E-ISSN: 2249-3379

How to cite this article: MahaHanafy Mahmoud, Hany Mohamed Ahmed Wahba, Marwa Hanafy Mahmoud. Evaluation of a new baby food prepared from *Opuntia ficus indica* fruit in Wistar rats with either calcium deficient or vitamin A depletion. J Adv Pharm Edu Res 2020;10(1):92-99.
 Source of Support: Nil, Conflict of Interest: None declared.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

compounds of plant origin with synthetic drugs because of a lower cost and fewer side effects^[5]. *Opuntia ficus indica* is Among those medicinal plants, which is a plant of wide distribution native to Mexico^[5]. It grows wild in arid and semiarid climates and in the semi-dry countries such as Italy, South-western USA, Tunisia, Morocco, Argentina, Brazil, Iran, Peru, Spain, and South Africa^[6]. It belongs to Cactaceae family^[7] and it is evergreen modified stem plant. *Opuntia ficus indica* is also known as prickly pear or cactus pear and it is consumed as food and in folk medicine due to its nutritional and beneficial properties^[8, 9]. The fruits of different varieties are of many colors including red, yellow, orange, and green differing in their physical and chemical composition and properties^[10]. From the health point of view, opuntia contains many compounds of health values including polyphenols, fibers, minerals, and vitamins. Its fiber content is of considerable amount and of distinct health benefits such as lowering blood cholesterol, weight control, diabetes control and reducing the risk of some types of cancer including the colon cancer^[11, 12]. Opuntia fruit is a good source of vitamins and minerals^[5, 13]. The red and yellow varieties comprise considerable amounts of vitamin A and carotenes, which are known for their antioxidant properties and not only having an essential role in the processes of vision, reproduction, and cellular growth but also needed for the maintenance of healthy skin and mucosa and protects against lung cancer^[14]. Vitamin A deficiency leads to blindness, xerophthalmia, and premature death^[15]. Moreover, Opuntia is a good source of minerals such as calcium, manganese, magnesium, and iron. Calcium and magnesium are required for bone-strengthening. It was reported previously that consuming cactus improves the density of bone mineral in women with low bone mass^[16]. Cactus pear is a crop of economic importance that has numerous uses^[15]. Recently, this plant has been used to prepare some beverages and many other food products^[6]. Opuntia has been used in the elaboration of jellies, candies and beverages^[17]. Also, it has been incorporated in cake and bread^[18]. Bad feeding practices and the lack of suitable, convenient, and affordable baby foods are responsible mainly for malnutrition in children especially in developing countries^[19]. Nowadays, there is an increasing demand for complementary baby foods all over the world. There are a wide variety of complementary baby foods in the market including the cereal-based baby foods, vegetables, and fruits baby foods either ready-to-use or prepared by rehydration in milk or water^[20]. Recently, powdered food products are increasingly gaining more preference among consumers' choices as they meet their everyday life demands for convenient, smart, and easy-to-use applications^[21]. The aim of the present study was to prepare a low cost powdered ready-to-use baby food from *Opuntia ficus indica* fruit which is a wild plant rich in minerals and vitamins and has good acceptability after selecting the best drying method for the fruit to ensure safety, longer shelf life, as well as better and easier handling of the prepared product. Also, to evaluate its biological effect in Wistar rats, which were calcium deficient or

vitamin A depleted on different biochemical parameters as well as body weight gain and feed intake.

Materials and Methods

Materials

Opuntia ficus indica fruits were obtained from the Egyptian local market during their ripening season in 2018.

Materials used for chemical analysis of the *Opuntia ficus* fruit

All solvents that were used for determination of calcium and β -carotene in opuntia fruit and those which were used for determination of vitamin A in serum by HPLC technique were of the analytical grade for HPLC and were obtained from Sigma-Aldrich Chemical Co. (St. Louis, Mo, 63103, USA).

Materials used for the animal experiment

Most of the components of the diet, introduced to the rats (table 1) were obtained from the local market, while, casein was obtained from Scerma Co., (France). The constituents of the salt and the vitamin mixtures used for the formulation of the diet were of analytical grade and obtained from Fluka (Germany) and BDH (England) Chemical Companies.

Animals used in this experiment were Wistar male albino rats obtained from the Central Animal House, National Research Centre (NRC), Egypt. The animal experiment was carried out according to the guidelines for Animal Care and Ethics Committee of the NRC (Egypt) and the study protocol was approved by the committee.

Materials used for biochemical analysis

Kit used for determination of blood hemoglobin was obtained from Biodiagnostic Co., (Egypt). Serum urea and creatinine determination kits were purchased from Chronolab Systems S.L., (Barcelona, Spain). Kits used for determination of total calcium, total protein, and albumin were obtained from Olympus Company for automated Olympus Autoanalyzer, Ireland.

Methods

Technological treatments

1. Preparation of the dried opuntia fruit

All tested samples of opuntia fruit were peeled, one sample was kept fresh and the other samples were dried on perforated trays by using the different drying methods according to Mahmoud et al.,^[22] as follows:

- a) Fresh opuntia fruit.
- b) Air oven drying: Opuntia fruit was placed on stainless steel trays in one layer and then placed on the oven and dried at a temperature of 40 °C for 8 hrs, the trays were removed out of the oven when the weight of samples was being constant.
- c) Solar drying: One layer of opuntia fruit was placed on stainless steel trays and dried for 48 hrs at 40 °C with ambient air temperature of 19.4 °C, relative humidity of 57%, relative humidity inside the dryer of 10% and global solar radiation of 845 W/m², the trays were removed out of the solar dryer when the weight of the sample was being constant.

- d) Microwave drying: One layer of opuntia fruit was placed in a microwave oven which was Samsung (Model MF245) with oven cavity dimensions of 419 × 245 × 428 mm and operation frequency of 2.450 MHz, with a power source of 230 V-50 Hz. The nominal microwave power outputting was 900 Ws, the air temperature was 40 °C /6 min.

2. Preparation of the baby food from opuntia fruit

Baby food was prepared according to Szulc and Lenart^[23] with some modifications. Solar dried opuntia was collected after fine milling blended to powder form (0.5 mm) in a high-speed blender (Braun KMM 30 mill, type 3045, CombiMax, Germany) and kept in a dry place to prevent aggregation of granules. A sample of baby food powder weighing 25 g was tested for rehydration as following: 25 g of powder was rehydrated in 100 ml water and examined for its taste.

Chemical analysis

1. Proximate composition

Proximate composition of the opuntia fruit samples was analyzed for moisture, protein, ash, and fat content according to the methods described in the AOAC^[24]. Their analyses were performed in triplicate.

2. Determination of β -carotenoids in *Opuntia ficus indica* fruit

According to Neves et al.,^[25] with some modifications, one g of the sample was placed in a test tube with 50 mL of an acetone extraction solution, then the tube was covered with aluminum foil. The extract was stirred for 1 min, incubated for 9 min, filtered through cotton, and immediately read in triplicate with a spectrophotometer (Shimadzu UV-2401 PC, Australia) at a wavelength of 450 nm. The calibration curve was obtained using β -carotene as a standard (Sigma-Aldrich Chemical Co., St. Louis, Mo, 63103, USA). The results were expressed as mg β -carotene/100 g sample.

3. Determination of calcium in *Opuntia ficus indica* fruit

The samples were subjected to analyze for the determination of calcium according to the method of Sahari et al.,^[26] using Optima 2000 DV inductively coupled plasma atomic emission spectrometer with full PC control (Perkin Elmer-Germany). The concentrations were obtained based on the calibration curve. High purity Merck standards were used.

Animal experiment

Thirty-six white male Wistar albino rats with a body weight ranging from 60 to 80 g were used. The standard control diet was prepared according to Reeves et al.^[27]. Animals were divided into 6 groups, each comprising 6 rats as follows;

Group 1: was given the control diet.

Group 2: was given the control diet + 15% dried opuntia fruit.

Group 3: was given the control diet which was calcium deficient as 1 g Ca/Kg diet according to Hunt et al.,^[28].

Group 4: was given the calcium deficient diet + 15% dried opuntia fruit.

Group 5: was given the control vitamin A depleted diet.

Group 6: was given the vitamin A depleted diet + 15% dried opuntia fruit.

Rats were housed individually in separate cages. The experiment lasted for 4 weeks during which, food and water were allowed ad-libitum to each rat. Body weight was measured twice a week. Food intake of each rat was daily recorded. At the end of the experiment, body weight gain, feed intake, and feed efficiency ratio for each rat were calculated (feed efficiency ratio = body weight gain/feed intake).

At the end of the experimental period, rats were fasted overnight and in the early morning, each rat in different groups was subjected to slight anesthesia by diethyl ether. Blood was withdrawn by open heart puncture and each sample was delivered into one heparinized tube and one dry tube.

Biochemical analysis

Blood hemoglobin concentration was immediately measured for each sample according to Betke and Savelsberg^[29]. Serum was separated from the second tube for each sample after centrifugation at 3000 rpm for 15 min and delivered into clean tubes, then, stored at -20 °C until analysis.

Serum urea and creatinine were assessed as described previously by Fawcett and Scott^[30] and Murray^[31], respectively. Serum total calcium, total protein, and albumin were measured according to the manufacturer's instructions using auto-analyzer Olympus Au 400. Serum ionized calcium was calculated based on data obtained for serum total calcium, total protein, and albumin from the following equation as described previously by Sava et al.^[32]:

$$\text{Ionized Ca (mg/dl)} = \frac{((6 * \text{Ca}) - ((0.19 * \text{T.P}) + \text{Alb}/3))}{(6 + ((0.19 * \text{T.P}) + \text{Alb}))/4}$$

where Ca is total calcium, T.P is total protein, and Alb is albumin.

Determination of serum vitamin A by HPLC analysis

Vitamins A in serum was determined by the HPLC technique as described previously by Hussein et al.^[33].

Sample extraction

First, 400 μ l of serum was added to 400 μ l of ethanol and 400 μ l of hexane, mixed well-using vortex and centrifuged for 5 min at 5000 rpm using cooling centrifuge (Laborzentrifugen, 2K15, Sigma, Germany). Then, the upper layer was removed and the precipitate was washed by 400 μ l of hexane. This step was repeated to collect the upper layer (hexane layer), which was then dried by nitrogen stream. Finally, the extract was redissolved in 500 μ l of mobile phase (ethanol and acetonitrile) and filtered by hydrophilic 0.45 μ m PVDF filter for removing any insoluble materials.

HPLC condition

Determination of vitamin A was carried out using a high-performance liquid chromatography (HPLC) system, Agilent technologies 1100 series (Agilent Technologies, Hewlett-Packard-Strasse 8, 76337 Waldbronn, Germany) equipped with a quaternary pump (Quat pump, G1310A model), G1328A manual injector (MI), UV detector, and a reversed phase

column (C18-25cm, 5 μ m particle size). Twenty μ l of the filtrate was injected onto the reversed phase column and isocratically eluted with acetonitrile/ethanol 50:50 (v/v) with 0.1% triethylamine as a mobile phase and was delivered at a flow rate of 1ml/min. The UV detector was set at 325 nm. Serial dilutions of vitamin A standard were injected into HPLC; the peak areas were determined. A linear standard curve was constructed by plotting peak areas vs the corresponding standard concentrations. The concentrations of the samples were obtained from the standard curve.

Statistical analysis

Data were statistically analyzed using the computerized program SPSS software, version "20" for Windows. The one way ANOVA test was done followed by the Duncan test. Data were represented as mean \pm SE. The level of significance $P < 0.05$ was considered as significant, otherwise non-significant.

Results and Discussion

Effect of drying on the chemical composition of opuntia

The primary goal of the drying was to reach the minimum moisture content in the samples to increase the shelf-life of the products but taking into account and being cautious not to burn the samples and also to avoid loss of nutrients during processing to maintain the nutritional quality of the prepared baby food product and therefore to support a satisfactory nutritional status when being introduced to babies [34]. The analysis of the proximate composition of the opuntia fruit either fresh or dried (Table 1) showed that there was a significant difference between the fresh and dried (powder) forms in the content of moisture, protein, fat, ash, and carbohydrates ($P < 0.05$). The obtained data for the proximate composition of the dried fruit was more or less similar to that obtained in other studies [35]. The ash content of the dried fruit was considered relatively high due to the high mineral content of opuntia and in particular, the calcium content, which was detected as 380.2 mg/Kg dry weight (Table 2) that was considered relatively high when compared to other plants. It was mentioned that the calcium content of opuntia fruit was relatively high which was in agreement with our results [36].

Table 1: Proximate composition of opuntia fruit.

	Moisture %	Protein %	Fat %	Ash %	Carbohydrates* %
Fresh	83.93 \pm 0.27a	0.96 \pm 0.06b	2.96 \pm 0.02b	0.38 \pm 0.01b	11.78
Powder	4.65 \pm 0.15b	4.54 \pm 0.23a	14.78 \pm 0.15a	1.89 \pm 0.04a	74.14

All values are represented as means of triplicate determinations \pm standard deviation (SD). Values that share the same letters at the same column are significantly different ($P < 0.05$). *Total carbohydrates: calculated by difference (100—Sum of all components).

Table 2: Concentration of calcium in opuntia fruit.

Group	Calcium (mg/ Kg dry wt.)
Opuntia fruit	380.2

Evaluation of the prepared baby food

1. Proximate composition and calcium content of opuntia fruit

Opuntia ficus indica fruit was chosen in the present study to prepare baby food as it is a rich source for vitamins and minerals particularly vitamin A and calcium [5]. Moreover, the opuntia fruit is mostly a wild plant, which means that no fertilizers and no pesticides are added to it in addition to its low cost that renders opuntia to be the plant of choice and the most suitable source for the baby food. In addition to its benefits, the sugar content of the opuntia fruit is relatively high [15], which makes it of a very good taste and acceptable when introduced to babies. Also, the form of the prepared baby food as a dried powder, which is ready to use by rehydration, easy to market, and has relatively longer shelf-life than other types of baby foods give advantages to the prepared baby food in the present study over other commercial types of baby foods, found in the local market. It can be used as it is or it can be added to cereal-based baby food.

2. Sensory evaluation of the prepared baby food

Table (3) summarizes the results of the sensory evaluation of Opuntia baby food sample formulated compared to the commercial one. Data indicated that there were significant differences ($P < 0.05$) between samples for the sensory attributes of color, taste, odor, flavor, and overall-acceptability. The results showed that Opuntia baby food had an acceptable and very good panel scores that may be due to its high sugar content besides its accepted flavor.

Table 3: Sensory evaluation of baby food formulated from opuntia powder.

Treatment	Color	Taste	Odor	Flavor	Overall acceptability
Opuntia powder	8.0 \pm 0.8 ^b	8.2 \pm 0.8	7.7 \pm 0.1 ^b	8.4 \pm 0.3 ^b	8.6 \pm 0.82 ^b
Com. baby food	8.2 \pm 0.5 ^a	8.7 \pm 0.5 ^a	8.8 \pm 0.8 ^a	8.6 \pm 0.6 ^a	8.8 \pm 0.83 ^a

3. Effect of different drying methods on β -carotene content of the prepared baby food

β -carotene content was detected in the present study to have varying concentrations according to the method of drying. As shown in fig. (1) drying opuntia fruit with any of the methods selected for drying, which were solar energy, oven, and microwave kept β -carotene content in the samples but with different concentrations. It is obvious that solar energy was the best method of drying, which gave the highest β -carotene concentration followed by microwave method, and then the oven method gave the least β -carotene concentration. So, solar energy was chosen to be the method for drying to prepare baby food. These obtained results are in accordance with the results obtained in a previous study for our team for the effect of drying methods on the antioxidant activity of orange peel, which revealed that solar energy is the best drying method [22].




Figure 1: Effect of drying methods on beta-carotene content

It is the worth mentioning that the fresh sample recorded a value for beta-carotene content of 3.19 mg/100 g sample, which is less than that for the sample that was dried by solar energy and also less than any of the values detected for dried samples by different drying techniques (Table 4).

Table 4: Concentration of beta-carotene in fresh opuntia fruit and dried opuntia fruit as baby food.

sample	beta-carotene (mg/100 g sample)
Fresh fruit	3.19
Dried fruit (solar energy)	8.24

Biological evaluation of the baby food

1. Effect of feeding opuntia fruit on body weight gain, feed intake, and feed efficiency ratio

Data obtained for the feed intake, body weight gain and feed efficiency ratio (FER) of the different groups (Table 5) showed a significant reduction in the body weight gain and feed intake for the group with calcium deficiency compared to the control negative group and the control group that received the opuntia fruit. These results seem to be in accordance with those obtained by Alomaim et al., [37] who reported that low dietary calcium reduced body weight and fat mass. Meanwhile, the group that with calcium deficiency and received opuntia fruit showed a rise in the body weight gain compared to the group with calcium deficiency, which did not receive opuntia fruit, but it was still significantly lower than the control negative group. It was previously reported that opuntia fruit is a rich source of calcium [18, 36]. Moreover, Aguilera-Barreiro et al., [16] mentioned that the consumption of opuntia in women with low bone mass improved the density of bones mineral.

Table 5: Feed intake, body weight gain, and food efficiency ratio (FER) of different groups compared to the control group.

Group	B. wt. gain(g)	Feed intake(g)	FER
Control (-ve)	124.50±4.42 ^a	397.33±12.99 ^a	0.489±0.02 ^{ab}
Control+ OF	125.83±5.61 ^a	395.83±11.64 ^a	0.516±0.01 ^{ab}
Vit. A def.	118.33±5.91 ^{ac}	365.00±1.59 ^a	0.528±0.01 ^b

Vit. A def. + OF	124.33±7.96 ^a	357.50±5.55 ^a	0.537±0.02 ^b
Ca def.	92.67±26.74 ^b	276.67±17.35 ^b	0.402±0.03 ^c
Ca Def.+ OF	103.33±1.36 ^{bc}	290.00±31.31 ^b	0.473±0.02 ^a

Values are expressed as mean ± SE and the mean difference is significant at P<0.05. Values that share the same letters at the same column are not significant while values that share different letters at the same column are significant.

2. Effect of feeding opuntia fruit on serum total and ionized calcium

Determination of total calcium and ionized calcium in serum revealed that there was a significant drop in both total and ionized calcium for the group of rats with calcium deficiency, however, this drop was more or less normalized in the group of rats with calcium deficiency, which received the opuntia fruit (fig. 2 & 3). It is well known that calcium deficiency leads to so many problems in the body among which is low bone mineralization that may further lead to osteoporosis. It was previously reported that the consumption of opuntia boosts the density of bone mineral in women with low bone mass [16]. It seems that these results are in accordance with the results obtained in this study.




Figure 2: A- Total calcium and B- Ionized calcium of group with calcium deficiency and group with calcium deficiency + opuntia fruit and the control groups. Bars represent mean ± SE and the mean difference is significant at P<0.05. Bars that share the same letter are not significant, while, those that share different letters are significant.

3. Effect of feeding opuntia fruit on serum vitamin A levels

Vitamin A has an important role in visual health, development, skin and mucosa maintenance, fetal growth and immune function. Vitamin A deficiency represents a public health problem worldwide, especially in South-East Asia and Africa. It can lead to visual impairment as night blindness and in children, may rise the risk of illness and death due to childhood infections, including those causing diarrhea and measles [38]. Data obtained in the present study revealed that there was a very highly significant drop of serum vitamin A in the group of rats that were fed on vitamin A depleted diet. On the other hand, the group of rats that were fed on vitamin A depleted diet and received opuntia fruit showed a rise in serum vitamin A, which is significantly higher compared to the group fed on

vitamin A depleted diet, but still significantly lower than the control negative group. *Opuntia* was reported to contain β -carotene which is vitamin A precursor with a comparatively higher amount than other plants [39]. Also, from the data obtained in this study, it can be obvious that vitamin A precursor of the *opuntia* fruit has good bioavailability.

4. Effect of feeding *opuntia* fruit on kidney function, blood proteins, and hemoglobin

No significant change was noticed for serum urea, creatinine, or total protein for all groups compared to the control negative group indicating normal kidney function for all groups (Table 6). However, there is a slight non-significant decrease in serum albumin for vitamin A depletion, vitamin A depletion + *opuntia*, and calcium deficient groups compared to the control negative group, but this decrease was significant when compared to the control group that received the *opuntia* fruit. No significant change was noticed for hemoglobin concentration for all groups compared to the control negative group. However, the non-significant decrease noticed in the group with vitamin A depletion and the group with calcium deficiency was restored in the group with vitamin A depletion that received the *opuntia* fruit and also the group with calcium deficiency that received the *opuntia* fruit. It was previously reported that *opuntia* fruit contains vitamins and minerals such as iron, folate, and vitamin B12 that are necessary for hemoglobin synthesis [40].




Figure 3: Serum vitamin A by HPLC of the group with vitamin A deficiency, the group with vitamin A deficiency + *opuntia* fruit and the control groups. Bars represent mean \pm SE and the mean difference is significant at $P < 0.05$ level. Bars that share the same letter are not significant, while those that share different letters are significant.

Table 6: Serum urea, creatinine, total protein, albumin, and hemoglobin (Hb %) of all groups.

Group	s.Urea (mg/dl)	s.Creat. (mg/dl)	s.T.prot. (g/dl)	s.Alb (g/dl)	Hb (%)
Control (-ve)	41.57 \pm 4.01 ^a	0.48 \pm 0.16 ^a	9.38 \pm 1.38 ^a	4.37 \pm 0.60 ^{ab}	13.80 \pm 0.44 ^{abc}
Control + OF	42.86 \pm 5.04 ^a	0.45 \pm 0.08 ^a	9.57 \pm 0.55 ^a	5.54 \pm 0.40 ^a	14.99 \pm 0.53 ^a

Vit.A depleted	50.60 \pm 3.65 ^a	0.36 \pm 0.08 ^a	9.16 \pm 0.40 ^a	3.90 \pm 0.44 ^b	12.80 \pm 0.23 ^c
Vit.A dep.+ OF	40.50 \pm 4.54 ^a	0.42 \pm 0.03 ^a	9.44 \pm 0.14 ^a	4.00 \pm 0.49 ^b	13.61 \pm 0.12 ^{abc}
Ca def.	45.05 \pm 5.06 ^a	0.40 \pm 0.07 ^a	9.49 \pm 0.37 ^a	4.23 \pm 0.11 ^b	13.14 \pm 0.31 ^{bc}
Ca Def. + OF	42.69 \pm 2.63 ^a	0.36 \pm 0.07 ^a	9.24 \pm 0.31 ^a	4.36 \pm 0.17 ^{ab}	13.96 \pm 0.49 ^{abc}

Values are expressed as mean \pm SE and the mean difference is significant at $P < 0.05$. Values that share the same letters at the same column are not significant, while values that share different letters at the same column are significant.

Conclusion

It can be concluded from the obtained results in the present study that the prepared form of the baby food was of a very good and accepted sensory evaluation. Solar energy was found to be the best drying method that gave the highest value of β -carotene content among other used methods for drying, which were oven and microwave. The biological evaluation of the prepared baby food revealed that feeding rats with calcium deficient foods + dried *opuntia* fruit caused a rise in their serum total and ionized calcium since *opuntia* is a rich source of calcium. Also, feeding rats on a diet with vitamin A depletion + dried *opuntia* fruit had a good impact on serum vitamin A of these rats. Thus, it is recommended that the prepared baby food can be safely introduced for babies as it doesn't have any drawbacks on body weight gain or different body functions and it can protect babies from calcium and vitamin A deficiency. In addition, it can be used for babies in those populations that have vitamin A or calcium deficiency.

References

1. Chiu M, Dillon A, Watson S. Vitamin A deficiency and xerophthalmia in children of a developed country. *Journal of paediatrics and child health*. 2016 Jul;52(7):699-703.
2. UNICEF. Ready-to-use therapeutic food: Current outlook. UNICEF Supply Division. February 2017. Available at: https://www.unicef.org/supply/files/Ready-to-use_Therapeutic_Food_Current_Outlook.pdf.
3. Pettifor JM. What's new in hypophosphataemic rickets?. *European journal of Pediatrics*. 2008 May 1;167(5):493-9. doi: 10.1007/s00431-007-0662-1.
4. Bromage S, Ahmed T, Fawzi WW. Calcium deficiency in Bangladesh: burden and proposed solutions for the first 1000 days. *Food and nutrition bulletin*. 2016 Dec;37(4):475-93.
5. Aragona M, Lauriano ER, Pergolizzi S, Faggio C. *Opuntia ficus-indica* (L.) Miller as a source of bioactivity compounds for health and nutrition. *Natural*

- product research. 2018 Sep 2;32(17):2037-49. DOI: 10.1080/14786419.2017.1365073
6. Salehi E, Emam-Djomeh Z, Askari G, Fathi M. *Opuntia ficus indica* fruit gum: Extraction, characterization, antioxidant activity and functional properties. *Carbohydrate polymers*. 2019 Feb 15;206:565-72.
 7. Nharingo T, Moyo M. Application of *Opuntia ficus-indica* in bioremediation of wastewaters. A critical review. *Journal of environmental management*. 2016 Jan 15;166:55-72.
 8. Stintzing FC, Carle R. Cactus stems (*Opuntia* spp.): A review on their chemistry, technology, and uses. *Molecular nutrition & food research*. 2005 Feb;49(2):175-94.
 9. Pérez-Torrero E, Garcia-Tovar SE, Luna-Rodriguez LE, RodríguezGarcía ME. Chemical Composition of Prickly Pads from (*Opuntia ficus-indica* (L.) Miller Related to Maturity Stage and Environment. *International Journal of Plant Biology & Research*. 2017.
 10. Méndez LP, Flores FT, Martín JD, Rodríguez EM, Romero CD. Physicochemical characterization of cactus pads from *Opuntia dillenii* and *Opuntia ficus indica*. *Food chemistry*. 2015 Dec 1;188:393-8.
 11. Lansky EP, Paavilainen HM, Pawlus AD, Newman RA. *Ficus* spp.(fig): Ethnobotany and potential as anticancer and anti-inflammatory agents. *Journal of Ethnopharmacology*. 2008 Sep 26;119(2):195-213.
 12. Uebelhack R, Busch R, Alt F, Beah ZM, Chong PW. Effects of cactus fiber on the excretion of dietary fat in healthy subjects: a double blind, randomized, placebo-controlled, crossover clinical investigation. *Current Therapeutic Research*. 2014 Dec 1;76:39-44.
 13. Stintzing FC, Schieber A, Carle R. Phytochemical and nutritional significance of cactus pear. *European Food Research and Technology*. 2001 Mar 1;212(4):396-407.
 14. Valla A, Valla B, Le Guillou R, Cartier D, Dufossé L, Labia R. New Syntheses of Retinal and Its Acyclic Analog γ -Retinal by an Extended Aldol Reaction with a C6 Building Block That Incorporates a C5 Unit after Decarboxylation. A Formal Route to Lycopene and β -Carotene. *Helvetica chimica acta*. 2007 Mar;90(3):512-20.
 15. El Kharrassi Y, Mazri MA, Benyahia H, Benaouda H, Nasser B. Fruit and juice characteristics of 30 accessions of two cactus pear species (*Opuntia ficus indica* and *Opuntia megacantha*) from different regions of Morocco. *LWT-Food Science and Technology*. 2016 Jan 1;65:610-7.
 16. Aguilera-Barreiro MD, Rivera-Márquez JA, Trujillo-Arriaga HM, Tamayo y Orozco JA, Barreira-Mercado E, Rodríguez-García ME. Intake of dehydrated nopal (*Opuntia ficus indica*) improves bone mineral density and calciuria in adult Mexican women. *Food & nutrition research*. 2013 Jan 1;57(1):19106.
 17. Morales P, Ramírez-Moreno E, de Cortes Sanchez-Mata M, Carvalho AM, Ferreira IC. Nutritional and antioxidant properties of pulp and seeds of two *Opuntia joconostle* cultivars (*Opuntia joconostle* FAC Weber ex Diguët and *Opuntia matudae* Scheinvar) of high consumption in Mexico. *Food Research International*. 2012 Apr 1;46(1):279-85.
 18. Msaddak L, Abdelhedi O, Kridene A, Rateb M, Belbahri L, Ammar E, Nasri M, Zouari N. *Opuntia ficus-indica* cladodes as a functional ingredient: bioactive compounds profile and their effect on antioxidant quality of bread. *Lipids in health and disease*. 2017 Dec;16(1):32.
 19. Muhimbula HS, Issa-Zacharia A, Kinabo J. Formulation and sensory evaluation of complementary foods from local, cheap and readily available cereals and legumes in Iringa, Tanzania. *African Journal of Food Science*. 2011 Jan 31;5(1):26-31.
 20. da Silva ED, Leme AB, Cidade M, Cadore S. Evaluation of the bioaccessible fractions of Fe, Zn, Cu and Mn in baby foods. *Talanta*. 2013 Dec 15;117:184-8.
 21. Plati F, Matsakidou A, Kiosseoglou V, Paraskevopoulou A. Development of a dehydrated dressing-type emulsion with instant powder characteristics. *Food Structure*. 2019 Apr 1;20:100110. <https://doi.org/10.1016/j.foostr.2019.100110>.
 22. Mahmoud MH, Wahba HM, Mahmoud MH, Abu-Salem FM. Antagonizing the hazardous impact of increased oxidative stress in Wistar rats by biscuits with dried orange peel. *J Biol Sci*. 2018;18(1):21-31.
 23. Szulc K, Lenart A. Water vapour adsorption properties of agglomerated baby food powders. *Journal of food engineering*. 2012 Mar 1;109(1):135-41.
 24. Horwitz W, Chichilo P, Reynolds H. Official methods of analysis of the Association of Official Analytical Chemists. Official methods of analysis of the Association of Official Analytical Chemists.. 1970.
 25. Neves LC, da Silva VX, Pontis JA, Flach A, Roberto SR. Bioactive compounds and antioxidant activity in pre-harvest camu-camu [*Myrciaria dubia* (HBK) Mc Vaugh] fruits. *Scientia Horticulturae*. 2015 Apr 21;186:223-9.
 26. Sahari MA, Barzegar M, Radfar R. Effect of varieties on the composition of dates (*Phoenix dactylifera* L.)—note. *Food science and technology international*. 2007 Aug;13(4):269-75. DOI: 10.1177/1082013207082244
 27. Reeves PG, Nielsen FH, Fahey Jr GC. AIN-93 purified diets for laboratory rodents: final report of the American Institute of Nutrition ad hoc writing committee on the reformulation of the AIN-76A rodent diet.
 28. Hunt JR, Hunt CD, Zito CA, Idso JP, Johnson LK. Calcium requirements of growing rats based on bone mass, structure, or biomechanical strength are similar. *The Journal of nutrition*. 2008 Aug 1;138(8):1462-8. <https://doi.org/10.1093/jn/138.8.1462>.

29. Betke K, Savelsberg W. Stufenphotometrische Hemoglobinbestimmungsmittel aus Cyanohämoglobin. *Z. Biochem.* 1950; 320, 431.
30. Fawcett JK, Scott J. A rapid and precise method for the determination of urea. *Journal of clinical pathology.* 1960 Mar 1;13(2):156-9. DOI <https://doi.org/10.1136/jcp.13.2.156>
31. Murray, R. L. Creatinine. *Clinical Chemistry, Theory, Analysis and Correlation.* L. A. Kaplan, A. J. Pesce (Eds.). CV Mosby Co., St. Louis, Toronto, Princeton, 1984; 1247-1253.
32. Sava L, Pillai S, More U, Sontakke A. Serum calcium measurement: total versus free (ionized) calcium. *Indian Journal of Clinical Biochemistry.* 2005 Jul 1;20(2):158-61.
33. Hussein J, Abo-elmatty D, El-Khayat Z, Abdel Latif Y, Saleh S, Farrag AR, Abd-El-Ghany W. Kiwifruit extract attenuates DNA damage and vitamins reduction in indomethacin-induced experimental gastric ulcer. *Jokull J.* 2015;65:2-16.
34. Mesías M, Wagner M, George S, Morales FJ. Impact of conventional sterilization and ohmic heating on the amino acid profile in vegetable baby foods. *Innovative Food Science & Emerging Technologies.* 2016 Apr 1;34:24-8.
35. Astello-García MG, Cervantes I, Nair V, del Socorro Santos-Díaz M, Reyes-Agüero A, Guéraud F, Negre-Salvayre A, Rossignol M, Cisneros-Zevallos L, de la Rosa AP. Chemical composition and phenolic compounds profile of cladodes from *Opuntia* spp. cultivars with different domestication gradient. *Journal of Food Composition and Analysis.* 2015 Nov 1;43:119-30.
36. Hernández-Urbiola MI, Pérez-Torrero E, Rodríguez-García ME. Chemical analysis of nutritional content of prickly pears (*Opuntia ficus indica*) at varied ages in an organic harvest. *International journal of environmental research and public health.* 2011 May;8(5):1287-95. doi:10.3390/ijerph8051287.
37. Alomaim H, Griffin P, Swist E, Plouffe LJ, Vandelloo M, Demonty I, Kumar A, Bertinato J. Dietary calcium affects body composition and lipid metabolism in rats. *PloS one.* 2019 Jan 10;14(1):e0210760. <https://doi.org/10.1371/journal.pone.0210760>
38. Vitamin A supplementation in postpartum women, WHO, 2017.
39. Chahdoura H, Morales P, Barreira JC, Barros L, Fernández-Ruiz V, Ferreira IC, Achour L. Dietary fiber, mineral elements profile and macronutrients composition in different edible parts of *Opuntia microdasys* (Lehm.) Pfeiff and *Opuntia macrorhiza* (Engelm.). *LWT-Food Science and Technology.* 2015 Nov 1;64(1):446-51.
40. del Socorro Santos Díaz M, Barba de la Rosa AP, Héliès-Toussaint C, Guéraud F, Nègre-Salvayre A. *Opuntia* spp.: characterization and benefits in chronic diseases. *Oxidative medicine and cellular longevity.* 2017;2017. <https://doi.org/10.1155/2017/8634249>.