

Anti-inflammatory activity of Caralluma Fimbrata- raw 264 macrophage cell lines

P. Keshaav Krishnaa¹, Lakshmi Thangavelu^{2*}, Anitha Roy²

¹Under Graduate Student, Saveetha Dental College, SIMATS, Saveetha University, Chennai, India, ²Associate Professor, Department of Pharmacology, Saveetha Dental College, SIMATS, Saveetha University, Chennai, India.

Correspondence: Lakshmi Thangavelu, Associate Professor, Department of Pharmacology, Saveetha Dental College, SIMATS, Saveetha University, Chennai, India. E_mail: lakshmi085@gmail.com

ABSTRACT

Introduction: The aim of the study was to estimate the anti-inflammatory activity of Caralluma Fimbrata on Raw 264 Macrophage cell lines enhanced with LPS which stands for lipo polysaccharide. **Materials and Methods:** The ethanolic extract of Caralluma Fimbrata was obtained from Green Chem Herbal Extract, Bangalore as a gift sample. Raw 264 macrophage cell lines were obtained from NCSS, Pune for the study. The sample was then studied along with the obtained cell lines to estimate the expression of nitrous oxide. Nitrous oxide being a known substance involved in inflammation. The gene expression of iNOS was also studied. **Results:** The effects of different concentrations of Caralluma Fimbrata was assessed and the nitrous oxide levels was estimated. It was observed that with increased concentration of the ethanolic extract of Caralluma Fimbrata the nitrous oxide Level was lowered and thus it is proven that Caralluma Fimbrata possesses anti-inflammatory properties. **Conclusion:** Caralluma Fimbrata extract possesses anti-inflammatory activity showed by significantly decrease in production of pro-inflammatory mediator NO and also by the suppression by the iNOS gene expression even when enhanced with LPS.

Keywords: Anti-inflammatory, Anti inflammation, Caralluma fimbrata, Anti arthritic, Nitrous oxide.

Introduction

Nature has provided a complete store-house of remedies to cure all ailments of mankind [1]. This is where, nature provides us drugs in the form of herbs, plants and algae to cure the incurable diseases without any toxic effect [2]. Research on medicinal plants is an important fact of biochemical research in India because of several reasons. Inflammation is a disorder involving localized increases in the number of leukocytes and a variety of complex mediator molecules [3-7]. Prostaglandins are ubiquitous substances that indicate and modulate cell and tissue responses involved in inflammation [4, 8-12]. Their biosynthesis has also been implicated in the pathophysiology of cardiovascular diseases, cancer, colonic adenomas and Alzheimer's disease [5, 13-17]. Medicinal plants are believed to be an important source of new chemical substances with potential therapeutic effects [6-8, 18, 19].

Macrophages after LPS stimulation produce NO by up-regulating iNOS expression through mitogen-activated protein kinases (MAPK) and NF- κ B signaling pathways. In response to macrophage activation, LPS stimulates a Toll-like receptor 4 (TLR4)-mediated myeloid differentiation factor (MyD88)-dependent pathway, which in turn activates the transforming growth factor- β -activated protein kinase 1 (TAK1), which subsequently results in activation of nuclear factor- κ B (NF- κ B) and activating protein-1 (AP-1), and produces inflammatory cytokines including TNF- α , IL-6, and IL-1 β [9, 10, 20-22]. Therefore, inhibition of these inflammatory mediators has been considered as an effective strategy for the development of anti-inflammatory drugs.

The aim of the study is to evaluate the anti-inflammatory activity of Caralluma Fimbrata on Raw 264 Macrophage cell lines.

Materials and Methods

Plant Material

Caralluma fimbrata bark ethanolic extract used for the study were obtained from Green Chem Herbal extract and Formulations, Bengaluru as a gift sample.

Chemicals

Access this article online

Website: www.japer.in

E-ISSN: 2249-3379

How to cite this article: P. Keshaav Krishnaa, Lakshmi Thangavelu, Anitha Roy. Anti-inflammatory activity of Caralluma Fimbrata- raw 264 macrophage cell lines. J Adv Pharm Edu Res 2018;8(2):21-24.

Source of Support: Nil, Conflict of Interest: None declared.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Lipopolysaccharide (LPS), Phenol free Dulbecco's modified Eagle medium (DMEM), MTT, Dimethyl sulphoxide (DMSO), phosphate buffer saline (PBS), and antibiotic-anti mycotic solution (100U penicillin, 100µg streptomycin, and 0.25µg amphotericin B per ml) were purchased from Sigma-Aldrich. Fetal bovineserum was purchased from GIBCO/BRL Invitrogen.

Cell culture

Macrophage RAW 264.7 cells were obtained from the NCCS, Pune with Passage no 16. Cells were cultured in phenol red-free Dulbecco's modified Eagle medium (DMEM) supplemented with 100units/ml penicillin, 100µg/ml streptomycin, and 10%heat-inactivated fetal bovine serum at 37°C with5% CO2. Cells were washed with DMEM medium and detached with 0.25% trypsin-EDTA. The cells were re-suspended in DMEM medium at a density of 2 x 106cells/ml.

Estimation Nitric oxide (NO)

The presence of nitrite, a stable oxidized product of nitric oxide (NO), was determined in cell culture media using Griess reagent. Briefly, 50 µl of supernatant from the test culture was mixed with 50 µl of 1% (w/v) sulphanilic acid in 5% (v/v) phosphoric acid in a 96-well plate, followed by incubation for 10 min at room temperature. After that 50 µl 0.1% (w/v) N-1-naphthylethylenediamineHCl in distilled water was added and incubated for 10 min at room temperature. The optical density at 540 nm was measured with a micro plate reader. The NO concentration was calculated by comparison with a NaNO2 (0–100 µM) standard curve. The final concentration of DMSO was adjusted to less than 0.1% for all treatments. The results were expressed as inhibition of NO production compared to the control (LPS) using: $([\text{nitrite}]_c - [\text{nitrite}]_t)/[\text{nitrite}]_c$, where $[\text{nitrite}]_c$ and $[\text{nitrite}]_t$ are the nitrite concentration in the control and test sample, respectively.

RNA Isolation and q - PCR Analysis

RAW macrophages were treated with 30µg/ml, 60µg/ml and 90µg/ml of Caralluma Fimbrata extract with 1µg/ml of LPS and incubated for 24h. Total RNA was isolated using TRIzol reagent (Invitrogen) according to the manufacturer's protocol, and 2µg of RNA was used for complementary DNA synthesis using M-MLV reverse transcriptase (Promega, Madison, WI, USA). Quantitative real-time polymerase chain reaction (q-PCR) was performed in an ABI 7500 Real-Time System with SYBR Green PCR Master Mix (Takara). Reactions were initiated with an initial incubation at 50°C for two minutes and 94°C for 10 min, followed by40 cycles of 94°C for 5s, 60°C for 15s, and 72°C for 10 s. The relative gene expression levels werecalculated using the 2–ΔΔCt method. The specific primer sequences used were given below:

INOS: Forward: 5'-ATGTCCGAAGCAAACATCAC-3'

Reverse: 5'-TAATGTCCAGGAAGTAGGTG-3'

β-actin was used as an internal reference gene between different samples.

Statistical analysis

Data obtained from the experiments were expressed as Mean ± SEM. The Statistical analysis of the difference between the groups was evaluated by Dunnett's following one-way ANOVA Post hoc comparisons in Graph pad Prism 5.0 software version. p<0.001, p<0.01 and p<0.05 were considered to be statistically significant.

Results

Effect of Caralluma fimbriata on NO production

NO is a pluripotent signaling molecule produced by different isoforms of enzyme, Nitric enzyme synthase (NOS). Despite possessing various beneficial effects, overproduction of nitric oxide lead to various disorders including inflammatory diseases. Hence effect of different concentrations Carulluma extract was evaluated on LPS induced NO production. LPS significantly increased NO production in RAW macrophages. The level of NO increased by LPS induction was significantly decreased in a dose – dependent manner when treated with different concentrations of Carulluma extract.

Table 1: The effect of Carulluma fimbriata on NO production in LPS stimulated RAW 264.7 macrophages

S.No	Concentration (µg/ml)	% of NO production in LPS Stimulated RAW 264.7 macrophages
1	3.175	80.84±0.60
2	6.25	75.01±0.56
3	12.5	68.57±0.52
4	25	59.73±0.34
5	50	47.85±1.23
6	75	29.71±0.52
7	150	8.61±0.76

Values are expressed Mean±SEM (n = 3)





Figure 1: Graphical representation of the effect of Carulluma fimbriata on production of NO production in LPS stimulated RAW 264.7 macrophages

Gene Expression of iNOS

LPS stimulation of RAW macrophages strongly upregulated the iNOS gene expression levels. However, when Carulluma was added at three different doses of 25µg/ml, 50µg/ml and 75µg/ml, the iNOS levels was significantly suppressed compared to that of LPS treatment only.



Values are expressed Mean±SEM (n = 3)

Figure 2: The effect of Carulluma fimbriata on LPS – stimulated iNOS expression in RAW 264.7 macrophages.

Discussion

Several studies have demonstrated the properties of various compounds from plants tend to possess rich pharmacological properties that play beneficial roles in many different conditions, including inflammation-related diseases [23-25]. Inflammation is a dynamic process involving proinflammatory cytokines such as nitrous oxide and it acts as important biological response toward injury [26, 27]. In the present study, we have examined the anti-inflammatory activity of Caralluma fimbriata n Raw 264 macrophage cell lines enhanced with LPS.

NO is a free radical produced from L-arginine by nitric oxide synthases (NOSs), and an important cellular second messenger [28-31]. The modulation of iNOS-mediated NO release is one of the major contributing factors during the inflammatory process [32-37]. Thus suppression of the response of NO would in turn indicate the anti-inflammatory activity of the particular extract. NO has the property to modify or generate intercellular signals and thus it has an effect on Immune cells, tumour cells and the cells of different tissues or organs.

From the above results of the present study it is evident that Caralluma fimbriata as an anti-inflammatory activity from table 1. On increasing the concentration of the extract the nitrous oxide level continues to decrease and thereby inhibiting the effect of iNOS mediated inflammation. Thus the anti-inflammatory effect of Caralluma fimbriata has been confirmed. The limitations of the study would include that only the NO pro-inflammatory messenger system has been analyzed in the current study. Other secondary messengers [38-40] in case of inflammatory reactions can be analyzed with the same extract and a conclusive final analysis of the particular source can be obtained.

Conclusion

Caralluma fimbriata Extract possess anti-inflammatory activity showed by significantly decrease in production of pro-inflammatory mediator NO and also by the suppression by the iNOS gene expression even when enhanced with LPS.

References

- Kokate CK, Purohit AP, Gokhale SB. Textbook of Pharmacognosy, Nirali Prakasan: Pune; 2002, 18; p1-4.
- Trease GE, Evans MC. Text book of Pharmacognosy. 12th edition. Balliere, Tindall: London; 1983, p 343-383
- Chopra RN, Nayar SL, Chopra IC. Glossary of Indian Medicinal Plants. CSIR: New Delhi; 1956, p -229.
- Anti-nociceptive activity of leaf extract of Caralluma fimbriata, using albino mice; Sayanathan Ray; Journal of Pharmaceutical and Biomedical Sciences Volume 18 Issue 15.
- Turner R A; Screening Methods in Pharmacology; Academic Press; New York 1965,100.
- Report of KS Laddha, Medicinal Natural Products Research Laboratory, University of Mumbai, Matunga, Mumbai, India.
- Gupta M, Mazumder UK, Gomathi P, Thamilselvan V. Anti-inflammatory evaluation of leaves of Plumeria acuminata. BMC Complementary and alternative medicine,2006; 6(36):1472-6882.
- Wealth of India. A Dictionary of Indian Raw Materials and Industrial Products. 1992; 3:266-67.
- Hayashi K, Iida I, Nakao Yu, Nakao Yo, Kaneko K: Four pregnane glycosides, boucerosides, AI, AII, BI, and BII Phytochemistry. 1988; 27:3919- 3924.
- Abdel-Sattar E, Meselhy MR, Al-Yahya MAA: New oxypregnane glycosides from Caralluma penicillata. Planta Med. 2002; 68:430-434.
- Bader A, Braca A, De Tommasi N, Morelli I: Further constituents from Caralluma negevensis. Phytochemistry. 2003; 62:1277-81.
- Kamil M, Fjajaraj A, Ahmad F, Gunasekhar C, Samuel S, Chan K, Habibullah M: Identification and quantification of flavonoids from Caralluma arabica and its quality control studies. J Pharm Pharmacol,1999; 51:225.
- Farrokh-Eslamlou H, Oshnoui S, Alinejad V. Novel restricted access to vasectomy in Iran: addressing changing trends in vasectomy clients' characteristics over 16 years in northwestern Iran. Contraception. 2015 Nov;92(5):488-93. doi: 10.1016/j.contraception. 2015.07.010. Epub 2015 Jul 28.
- Saberi A, Hatamian HR, Nemati S, Banan R. Hearing statement in multiple sclerosis: a case control study using auditory brainstem responses and otoacoustic emissions. Acta Med Iran. 2012;50(10):679-83.
- Keshavarz P, Saberi A, Sharafshah A, Asgari K, Rezaei S. Association of BDNF G196A Gene Polymorphism with Ischemic Stroke Occurrence and its 6-Month Outcome in an Iranian Population. Top Stroke Rehabil. 2016 Aug;23(4):254-60. doi: 10.1080/10749357.2016.1141491. Epub 2016 Feb 17.
- Rezaei S, Asgari Mobarake K, Saberi A, Keshavarz P, Leili EK. Brain-derived neurotrophic factor (BDNF) Val66Met polymorphism and post-stroke dementia: a hospital-based study from northern Iran. Neurol Sci. 2016 Jun;37(6):935-42. doi: 10.1007/s10072-016-2520-2. Epub 2016 Apr 12.
- Malekzadeh S, Azami M, Mirzaei M, Motamedi F. Comparative Investigation of Health Literacy Level of Cardiovascular Patients Hospitalized in Private and Educational Hospitals of Kerman City, Iran. Acta Inform Med. 2016 Feb;24(1):56-60. doi: 10.5455/aim.2016.24.56-60. Epub 2016 Feb 2.
- Azami M, Fatahi SR. Database graphic user interface correspondence with Ellis information seeking behavior model. Information Sciences and Technology.2010, 25(2): 247-264
- Aazami M, Yousefzadeh E, Namdar F. Faculty members of Kerman University of medical sciences' participation in science production and investigation of effective factors. Wuhan Daxue Xuebao (Xinxi Kexue Ban)/Geomatics and Information Science of Wuhan University 2010; 35(2).
- Alamolhoda SH, Kariman N, Hoseinpanah F, Alavi Majd H. Relationship between maternal hemoglobin level in first

- trimester with gestational diabetes mellitus. 2010, Iranian Journal of Endocrinology and Metabolism. 11(6), pp. 661-666+734.
19. Kariman N, Alamolhoda SH, Hosein Panah A, Alavi Majd H. Relationship between cigarette smoking and gestational diabetes mellitus. 2010, Journal of Zanjan University of Medical Sciences and Health Services. 18(71), pp. 34-41.
 20. Kariman N, Hedayati M, Taheri Z, Fallahian M, Salehpoor S, Alavi Majd SHA. Comparison of ELISA and three rapid HCG dipsticks in diagnosis of premature rupture of membranes. Iran Red Crescent Med J. 2011 Jun;13(6):415-9. Epub 2011 Jun 1.
 21. Haidari M, Nazer MR, Ahmadinejad M, Almasi V, Khorramabadi MS, Pournia Y. Honey in the treatment of Fournier's gangrene as an adjuvant: a cross sectional study. J Pak Med Assoc. 2014 May;64(5):571-3.
 22. Hemmati L, Rojhani-Shirazi Z, Ebrahimi S. Effects of Plantar Flexor Muscle Static Stretching Alone and Combined with Massage on Postural Balance. Ann Rehabil Med. 2016 Oct;40(5):845-850. Epub 2016 Oct 31.
 23. Musa AM, Cooperwood SJ, Khan MOF. A review of coumarin derivatives in pharmacotherapy of breast cancer. Curr Med Chem. 2008;15(26):2664-2679.
 24. A. Trishala, T. Lakshmi; Anti-inflammatory activity of Morinda Citrifolia extract-an invivo study, International Journal of Pharmacy and Technology, Vol 9(1), 2015; 29282-29286.
 25. Ghayathri, T. Lakshmi; Anti-inflammatory activity of acacia catechu bark extract in vitro study; Journal of Chemical and Pharmaceutical Research, Vol 7(7), 2015,1184-1187.
 26. Dinarello CA. Anti-inflammatory agents present and future. National Institute of Health. 2010;140(6):935-950.
 27. Alinejad V, Shadmehr A. Investigation of the factors affecting the positivity of the transplant result. Social Sciences (Pakistan), Volume 11, Issue 6, 2016, Pages 906-909.
 28. Palmer RM, Ashton DS, Moncada S. Vascular endothelial cells synthesize nitric oxide from L-arginine. Nature. 1988; 333:664-6.
 29. Gandjalikhan Nassab SA, Bahrami A, Keshtkar MM, Saffaripour MH. Comparison between two methods for enhancing Focal neuropathies following percutaneous nephrolithotomy (PCNL)--preliminary study. Ger Med Sci. 2013 Jun 13;11: Doc07. doi: 10.3205/000175. Print 2013.
 30. g heat transfer in separated convection flows-second law analysis. 2014. Iranian Journal of Science and Technology - Transactions of Mechanical Engineering. Vol. 38, No. M1, pp 9-23.
 31. Nasseh H, Pourreza F, Saberi A, Kazemnejad E, Kalantari BB, Falahatkar S. Focal neuropathies following percutaneous nephrolithotomy (PCNL)--preliminary study. Ger Med Sci. 2013 Jun 13;11: Doc07. doi: 10.3205/000175. Print 2013.
 32. Kim ND, Kim EM, Kang KW, Cho MK, Choi SY, Kim SG. Ginsenoside Rg3 inhibits phenylephrine-induced vascular contraction through induction of nitric oxide synthase. Br J Pharmacol. 2003; 140:661-70.
 33. Araban M, Kariman N, Tavafian SS, Motesaddi S, Alavimajd H, Shokravi FA. Air pollution and low birth weight: a historical cohort study from Tehran. East Mediterr Health J. 2012 Jun;18(6):556-60.
 34. Kariman N, Afrakhte M, Hedayati M, Fallahian M, Alavi Majd H. Diagnosis of premature rupture of membranes by assessment of urea and creatinine in vaginal washing fluid. Iran J Reprod Med. 2013 Feb;11(2):93-100.
 35. Keshtkar MM, Talebizadeh P. Multi-objective optimization of cooling water package based on 3E analysis: A case study. Volume 134, 1 September 2017, Pages 840-849.
 36. Keshtkar MM. Performance analysis of a counter flow wet cooling tower and selection of optimum operative condition by MCDM-TOPSIS method. Applied Thermal Engineering. Volume 114, 5 March 2017, Pages 776-784.
 37. Shahrzad E, Zahiri S, Ghasemi F, Kargar Jahromi H. A study of effects of L-carnitine on morphology and apoptosis in cryopreserved sperm. 2013. Advances in Environmental Biology, 7(9): 2126-2134.
 38. Krenn L, Paper DH. Inhibition of angiogenesis and inflammation by an extract of red clover (*Trifolium pretense* L.) Phytomed. 2009; 16:1083-1088.
 39. Sahu D. 8-methoxy chromen-2-one alleviates collagen induced arthritis by down regulating nitric oxide, NFkB and proinflammatory cytokines. Science Direct. 2015;29(2):891-900.
 40. Heidari T, Kariman N, Afrakhteh M, Alavi Majd H. The study of relationship between menorrhagia and gestational diabetes. 2008, Koimesh. 9(2), pp. 147-153+22.