

Anti-diabetic activity of *Sesbania grandiflora* - alpha amylase inhibitory effect

Shreya Kothari¹, Lakshmi Thangavelu^{2*}, Anitha Roy²

¹Under Graduate Student, Saveetha Dental College, SIMATS, Saveetha University, Chennai, India, ²Associate Professor, Department of Pharmacology, Saveetha Dental College, SIMATS, Saveetha University, Chennai, India.

Correspondence: Lakshmi Thangavelu, Associate Professor, Department of Pharmacology, Saveetha Dental College, SIMATS, Saveetha University, Chennai, India. E_mail: lakshmi085@gmail.com

ABSTRACT

Introduction: Diabetes mellitus is characterised by high blood sugar levels caused due to ineffectiveness of the insulin produced at the peripheral tissues. Due to the side effects of insulin and oral hypoglycaemic agent, demand to use natural products with anti-diabetic activity has been increased recently. **Materials and Methodology:** *Sesbania grandiflora* extract used in the study is obtained from Green Chem Herbal Extracts and Formulations, Bengaluru, India. Acarbose, Dinitrosalicylic acid, DMSO is obtained from Sigma Aldrich Co. India. α -Amylase enzyme were procured from HiMedia, Mumbai. In vitro amylase inhibition was studied by the method of Bernfeld and the percentage inhibition of α -Amylase enzyme was calculated. **Results:** A maximum inhibition of 81% was observed at 1000 μ g/ml which was compared with that of standard Acarbose that showed 93% inhibition. The IC₅₀ of the extract was found to be 50.95 μ g/ml and for Acarbose 34.83 μ M. **Conclusion:** The study showed that this plant species exhibited anti-diabetic property and can be used to explore possibilities of developing a natural drug for the management of diabetes mellitus rather than using synthetic drugs which produce side effects.

Keywords: *Sesbania grandiflora*, anti-diabetic activity, glycation, diabetes mellitus, alpha amylase.

Introduction

Sesbania grandiflora commonly known as vegetable hummingbird is a small tree in the genus *Sesbania*. The tree thrives under full exposure to sunshine and is extremely frost sensitive [1]. This fast growing tree is cultivated in many parts of India and Sri Lanka and has many traditional uses [2]. The leaf extract inhibits the formation of advanced glycation end-products. It contains linolenic acid and aspartic acid, which were found to be the major compounds responsible for the anti-glycation potential of the leaf extract [3,4-8].

The flowers have an astringent action on the body used in liver and spleen disorders, rhinitis, night blindness, abdominal pain etc. and helps in the detoxification of the body. The leaves

possess tonic properties which is helpful in worm infestations and bleeding disorders. The leaf paste is also used to treat oral and throat problems. The root bark is externally applied to reduce pain and inflammation in arthritis and gout. This plant also possesses anti-anxiolytic and anti-proliferative properties against cancer cells in humans [5,9,10].

Diabetes mellitus is a complex chronic condition that is a major source of ill health worldwide. It is characterised by hyperglycaemia and imbalance of carbohydrate, protein, and fat metabolisms [6,11-15]. This may be attributed to insulin inactivity or resistance, as a direct result of destruction or dysfunction of the beta-cells of the pancreas [7,16,17].

India is emerging as the diabetic capital of the world. Facing the burden of the consequences that the disease brings; it is currently estimated that every fifth diabetic in the world is an Indian [8,18,19]. Moreover, statistical projection about India suggests that the number of diabetics will rise from 15 million in 1995 to 57 million in the year 2025, the highest number of diabetics in the world. Reasons for this rise include increase in sedentary lifestyle, consumption of energy-rich diet, obesity, higher life span, etc. [9,20]. Inhibition of α -amylase enzymes involved in the digestion of carbohydrates, can significantly decrease the postprandial increase of blood glucose after a mixed carbohydrate diet and therefore can be an important strategy in

Access this article online

Website: www.japer.in

E-ISSN: 2249-3379

How to cite this article: Shreya Kothari, Lakshmi Thangavelu, Anitha Roy. Anti-diabetic activity of *Sesbania grandiflora* - alpha amylase inhibitory effect. J Adv Pharm Edu Res 2017;7(4):499-502.

Source of Support: Nil, Conflict of Interest: None declared.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

the management of postprandial blood glucose level in type 2 diabetic patients and borderline patients [10,21,22].

Control of postprandial hyperglycaemia is the key to the management of type 2 diabetes mellitus and its associated complications. Conventional approach to this growing epidemic is associated with many side effects, and moreover, it is expensive and inaccessible to certain communities [11, 12,23]. Due to this, evaluation of plant products is of growing interest since they contain many bioactive substances. Currently there is renewed interest in plant-based medicines modulating physiological effects in the prevention and cure of diabetes and obesity [13,24,25].

Hence the aim of this research is to scientifically investigate the anti-diabetic activity of *Sesbania grandiflora* in order to develop a natural drug instead of a synthetic one.

Materials and Methodology

Plant materials:

Sesbania grandiflora extract used in the study obtained from Green Chem Herbal Extracts and Formulations, Bengaluru, India.

Chemicals used:

Acarbose, Dinitrosalicylic acid, DMSO is obtained from Sigma Aldrich Co. India. α -Amylase enzyme was procured from HiMedia, Mumbai.

In-vitro α -Amylase inhibitory assay (Bernfeld 1955):

In vitro amylase inhibition was studied by the method of Bernfeld. In brief, 100 μ l of the different concentrations of *Sesbania* (1,3,10,30,100,300 and 1000 μ g/ml) was allowed to react with 200 μ l of α -Amylase (Hi media RM 638) and 100 μ l of 2mM of phosphate buffer (pH- 6.9). After 20-minute incubation, 100 μ l of 1% starch solution was added. For the controls, the same process was done that 200 μ l of the enzyme was replaced by buffer. Acarbose had the role of being used as the reference standard. 500 μ l of Dinitrosalicylic acid reagent was added to both control and test after incubation for 5 minutes. 5 minutes was the period for keeping them in boiling water. Using spectrophotometer, the absorbance was recorded at 540 nm and using the following formula, the percentage inhibition of α -Amylase enzyme was calculated:

$$\% \text{inhibition} = [(Control - Test) / Control] * 100$$

Suitable reagent blank and inhibitor controls were carried out simultaneously.

Results

α - amylase inhibitory activity:

The results of the present study exhibited strong α - amylase inhibitory activity of the *Sesbania* extract which is compared with standard Acarbose. As shown in Figure 1, maximum inhibition of 81% was observed at a concentration of 1000 μ g/ml by *Sesbania* extract, which was compared with that of standard Acarbose that showed inhibition of about 93%. The IC_{50} of the

extract was found to be 50.95 μ g/ml and for Acarbose 34.83 μ M.




Figure 1. alpha amylase inhibitory assay

Discussion

Many herbal extracts are being used directly or indirectly for the preparation of many modern medicines. In this study, an in vitro alpha-amylase inhibitory effect of *Sesbania grandiflora* is performed to evaluate the anti-diabetic activity. It was seen in figure 1 that this plant had lowered the blood glucose level by showing a maximum inhibition of 81%. This is due to the presence of lignin, terpenes, tripenes and other constituents [26]. However, acarbose being the standard showed 93% inhibition. When the IC_{50} values of *Sesbania* and acarbose is compared, it was found that it's more than acarbose. Alpha-amylase catalyses the hydrolysis of alpha-1,4-glycosidic linkages of starch, glycogen and various oligosaccharides. The disaccharides will be broken down to simple sugars by alpha- glucose daze making it available for intestinal absorption. The inhibition of their activity in the digestive tract of humans is considered to be effective tool to control diabetes [27].

Other studies regarding the α - amylase inhibitory activity were also performed. It showed that *D. Villosa* showed an increase ranging from 10.71 to 71.88% with a concentration range of 1.5 μ g/ml - 1000 μ g/ml and the IC_{50} value was calculated as 72.44 μ g / ml when compared with acarbose, IC_{50} value 83.23 μ g/mL [28]. IC_{50} values of α -amylase inhibitory effects of ethanol and hexane extracts of *P. amarus* was also found. The hexane extract (at a concentration 100 μ g/mL) showed 75.32% of α -amylase inhibitory activity with IC_{50} value 48.92 μ g/mL. Both ethanol and hexane extracts showed appreciable α -amylase inhibitory effects when compared with acarbose [29]. In another study, a weak α -amylase inhibitory activity of 20% (v/v) extract of *A. paniculata* was seen. It showed similar α -glucosidase and amylase inhibitory effect. The IC_{50} values show that 20% (v/v) eth- anolic extract has a lower potency and has a preference for α -glucosidase over α -amylase [30].

Therefore, *Sesbania grandiflora* showed considerable α -amylase inhibitory effects when compared with acarbose. Several studies of anti-anxiolytic and anti-inflammatory properties have been performed too. From the above mentioned results, this extract can be used as an anti-diabetic agent and incorporated into the pharmacological preparations.

Conclusion

A good understanding of the anti-diabetic activity of the *Sesbania grandiflora* has been created and now is being utilised effectively for other research studies in the pharmacological arena. These newer discoveries will be very helpful in the treatment of one of the most common metabolic disorder- Diabetes. Thus, establishing a promising future in this field.

References

1. "Sesbania grandiflora", Natural Resources Conservation Service PLANTS Database. USDA. Retrieved 10 November 2015.
2. Kirtikar K. R. & B. D. Basu, Indian Medicinal Plants Vol-I, International Book Distributor & Publisher, Dehradun, Edition 2005, bks pp. 735–736.
3. Prasanna, Govindarajan. "Linolenic acid prevents early and advanced glycation end-products (AGEs) modification of albumin". International Journal of Biological Macromolecules. 95: 121–125, 2016.
4. Prasanna, Govindarajan. "Aspartic acid functions as carbonyl trapper to inhibit the formation of advanced glycation end products by chemical chaperone activity". Journal of Biomolecular Structure and Dynamics. 34 (5): 943–951, 2015.
5. Joshi SG (2000) Leguminosae: text book of medicinal plants. New Delhi: Oxford Publishing. 130 p.
6. Unwin N, Gan D, Whiting D, The IDF awareness and promoting action. Diabetes Res Clin Pract 2010; 87(1):2-3.
7. Nayak BS, Roberts L. Relationship between inflammatory markers, metabolic and anthropometric variables in the Caribbean type 2 diabetic patients with and without microvascular complications. J Inflamm (Lond) 2006; 3:17.
8. Kumar A, Goel MK, Jain RB, Khanna P, Chaudhary V. India towards diabetes control: Key issues. Australas Med J. 2013; 6:524–31.
9. Satyanarayana T., Hypoglycemic and antihyperglycemic effect of alcoholic extract of *Euphorbia leucophylla* and its fractions in normal and in alloxan induced diabetic rats. Pharmacog Mag, 2:244-53, (2006).
10. Ali H, Houghton PJ, Soumyanath A. α -Amylase inhibitory activity of some Malaysian plants used to treat diabetes; with particular reference to *Phyllanthus amarus*. J Ethnopharmacol 107: 449–455.
11. Chai TT, Khoo CS, Tee CS, Wong FC. Alpha glucosidase inhibitory and antioxidant potential of antidiabetic herb *Alternanthera sessilis*: Comparative analysis of leaf and callus solvent fractions. Pharmacogn Mag. 2016; 12:253–8.
12. Kibiti CM, Afolayan AJ. Herbal therapy: A review of emerging pharmacological tools in management of diabetes mellitus in Africa. PharmacognMag. 2015;11(Suppl S2):258–74.
13. Sharma UD., Cure of heart diseases with ayurvedic drugs. Sacitra Ayurveda, 47:95-6, (1994).
14. Shirinbaik SH, Samarai V, Alinejad V, Baghaei R. Studying on the Effects of Family-centered Care Program on the Incidence of Early Complications of Cataract Surgery Introduction and Purpose. Journal of Global Pharma Technology. 2016; 02(8):68-74.
15. Saberi A, Hatamian HR, Nemati S, Banan R. Hearing statement in multiple sclerosis: a case control study using auditory brainstem responses and otoacoustic emissions. Acta Med Iran. 2012;50(10):679-83.
16. Keshavarz P, Saberi A, Sharafshah A, Asgari K, Rezaei S. Association of BDNF G196A Gene Polymorphism with Ischemic Stroke Occurrence and its 6-Month Outcome in an Iranian Population. Top Stroke Rehabil. 2016 Aug;23(4):254-60. doi: 10.1080/10749357.2016.1141491. Epub 2016 Feb 17.
17. Rezaei S, Asgari Mobarake K, Saberi A, Keshavarz P, Leili EK. Brain-derived neurotrophic factor (BDNF) Val66Met polymorphism and post-stroke dementia: a hospital-based study from northern Iran. Neurol Sci. 2016 Jun;37(6):935-42. doi: 10.1007/s10072-016-2520-2. Epub 2016 Apr 12.
18. Malekzadeh S, Azami M, Mirzaei M, Motamedi F. Comparative Investigation of Health Literacy Level of Cardiovascular Patients Hospitalized in Private and Educational Hospitals of Kerman City, Iran. Acta Inform Med. 2016 Feb;24(1):56-60. doi: 10.5455/aim.2016.24.56-60. Epub 2016 Feb 2.
19. Azami M, Fatahi SR. Database graphic user interface correspondence with Ellis information seeking behavior model. Information Sciences and Technology. 2010, 25(2): 247-264
20. Aazami M, Yousefzadeh E, Namdar F. Faculty members of Kerman University of medical sciences' participation in science production and investigation of effective factors. Wuhan Daxue Xuebao (Xinxi Kexue Ban)/Geomatics and Information Science of Wuhan University 2010; 35(2).
21. Alamolhoda SH, Kariman N, Hoseinpanah F, Alavi Majd H. Relationship between maternal hemoglobin level in first trimester with gestational diabetes mellitus. 2010, Iranian Journal of Endocrinology and Metabolism. 11(6), pp. 661-666+734.
22. Kariman N, Alamolhoda SH, Hosein Panah A, Alavi Majd H. Relationship between cigarette smoking and gestational diabetes mellitus. 2010, Journal of Zanjan University of Medical Sciences and Health Services. 18(71), pp. 34-41.
23. Kariman N, Hedayati M, Taheri Z, Fallahian M, Salehpour S, Alavi Majd SHA. Comparison of ELISA and three rapid HCG dipsticks in diagnosis of premature rupture of membranes. Iran Red Crescent Med J. 2011 Jun;13(6):415-9. Epub 2011 Jun 1.

24. Haidari M, Nazer MR, Ahmadinejad M, Almasi V, Khorramabadi MS, Pournia Y. Honey in the treatment of Fournier's gangrene as an adjuvant: a cross sectional study. *J Pak Med Assoc.* 2014 May;64(5):571-3.
25. Hemmati L, Rohani-Shirazi Z, Ebrahimi S. Effects of Plantar Flexor Muscle Static Stretching Alone and Combined With Massage on Postural Balance. *Ann Rehabil Med.* 2016 Oct;40(5):845-850. Epub 2016 Oct 31.
26. McCue P, Vatter D, Shetty K. Inhibitory effect of clonal oregano extracts against porcine pancreatic amylase in vitro. *Asia Pac J Clin Nutr.* 2004; 13:401-8.
27. Van de Laar FA, Lucassen PL, Akkermans RP, Van de Lisdonk EH, Rutten GE, Van Well C. Alpha-glucosidase inhibitors for patients with type 2 diabetes: Results from a Cochrane systematic review and meta-analysis. *Diabetes Care* 2005; 28:166-75.
28. Roy A, Geetha RV. In vitro α -amylase and α -glucosidase inhibitory activities of the ethanolic extract of *Dioscorea villosa* tubers. *Int J Pharm Bio Sci.* 2013; 4:49-54.
29. Iniyar G. Tamil, B. Dineshkumar, M. Nandhakumar, M. Senthilkumar, and A. Mitra. In vitro study on alpha amylase inhibitory activity of an Indian medicinal plant, *Phyllanthus Amarus*. *Indian J Pharmacol.* 2010 Oct; 42(5): 280-282.
30. Rammohan Subramanian, M. Zaini Asmawi, Amirin Sadikun. In vitro α -glucosidase and α -amylase enzyme inhibitory effects of *Andrographis paniculata* extract and andrographolide. *Acta Biochimica Polonica* Vol.55 No.2/2008 391-398.