

# Management of perforation - A review

Mahalakshmi Nandakumar, Iffat Nasim\*

Department of Conservative Dentistry and Endodontics, Saveetha Dental College, Saveetha University, Chennai, Tamil Nadu, India.

**Correspondence:** Iffat Nasim, Department of Conservative Dentistry and Endodontics, Saveetha Dental College, Saveetha University, No.162, Poonamallee High Road, Chennai - 600 077, Tamil Nadu, India. Phone: +91-9940063567. E-mail: nasimiffat@gmail.com.

## ABSTRACT

The aim of this review is to discuss the etiology, classification, diagnosis, prognosis, and management of perforation. The objective is to review the literature and various related articles regarding the etiology, classification, diagnosis, prognosis, and management of perforation. The logical corollary is that perforation can be considered as an artificial communication between the root canal system to the supporting tissues of teeth or to the oral cavity. Successful treatment can be achieved provided that proper diagnosis and immediate sealing of the perforation are provided aimed at eliminating the risk of infection.

**Keywords:** Iatrogenic perforation, perforation repair material, MTA, bio dentine, intentional replantation

## Introduction

Root perforation is one of the most common mishaps of endodontic treatment that a clinician or even an endodontic will encounter on a regular basis. By definition, root perforations are artificial communication between the root canal systems and the supporting tissues of teeth or to the oral cavity.<sup>[1-7]</sup> It occurs in approximately 2-12% of endodontically treated teeth.<sup>[2, 8-12]</sup> This acts as a pathway for the ingress of microorganisms either from the oral cavity or periodontal tissues, thus contaminating the perforation site and preventing its healing process. If left untreated or unnoticed, it leads to secondary periodontal involvement, suppuration, fistula formation, and prognosis of such tooth becomes questionable and extraction becomes the most likely treatment option. This review discusses the etiology, classification, diagnosis, factors governing prognosis, and various treatment options.

## Etiology

Often, the causes of perforation can be iatrogenic or non-iatrogenic.<sup>[3, 13-19]</sup> Iatrogenic perforations occur mainly due to the lack of attention to the internal anatomy and failure to consider and anticipate the possible variations that can occur in the root canal system. These iatrogenic perforations can occur at any level of the endodontic treatment. During access cavity preparation, perforation tends to occur mainly at the coronal level; most common causes are misalignment of burs due to failure to appreciate the alignment and angulation of the tooth and its position, plunging the bur through relatively thin pulpal floor, or while negotiating canals in cases of completely calcified canals. During cleaning and shaping procedure, perforation tends to occur mainly at the

coronal-, middle-, or apical third of the root. Strip perforations occur at the coronal- third of small curved canals. The furcal area of the coronal- third in small curved roots is known as the “danger zone” as the dentin is relatively thin at these areas, in such cases, using a straight file or aggressive filing can cause strip perforation.<sup>[4, 20-24]</sup> Midroot perforation mainly occurs due to misuse of rotary instruments while preparing post-space or while trying to bypass a ledge or a separated instrument. Apical perforation occurs because of instrumentation beyond the apical foramen or due to the transportation of canals.

Non-iatrogenic causes include resorption,<sup>[5, 25-30]</sup> external, or internal, mostly secondary to trauma and caries that is extensively involving the furcal area.

## Classification

According to fess and trope, perforation can be classified based on time, size, and location.<sup>[1, 31-35]</sup>

### 1. Based on time:

- a. Fresh perforation - perforations that occurs at the same appointment characterized by fresh blood at the perforation site. These, if treated immediately or shortly after occurrence under aseptic conditions, have a good prognosis.
- b. Old perforation - previously not treated or unnoticed with likely bacterial infection, these have a questionable prognosis.

### 2. Based on the size:

- a. Small perforation - these are smaller than size 20 endodontic instruments because the mechanical damage to tissue is minimum, and the due to the ease of sealing, they have a good prognosis.
- b. Large perforation – it occurs mainly during postpreparation with significant tissue damage and obvious difficulty in providing adequate seal, there is obvious salivary contamination or coronal leakage, these have questionable prognosis.

### 3. Based on the location:

- a. Coronal perforation - perforations that occur coronal to the level of crestal bone and epithelial attachment, and they cause minimal damage to the supporting tissues and are easy to access and seal a good prognosis.

## Access this article online

Website: [www.japer.in](http://www.japer.in)

E-ISSN: 2249-3379

**How to cite this article:** Nandakumar M, Nasim I. Management of perforation - A review. J Adv Pharm Edu Res 2017;7(3):208-211.

**Source of Support:** Nil, Conflict of Interest: None declared.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

- b. Crestal perforation - perforations at the level of the epithelial attachment into the crestal bone, questionable prognosis.
- c. Apical perforation - perforations apical to the crestal bone and the epithelial attachment, minimal risk of salivary contamination, and hence, a good prognosis.

## Diagnosis

Accurate diagnosis of root perforation can be tricky. Sudden bleeding and pain during instrumentation are warning signs of a potential root perforation. Continuous and profuse bleeding will occur in case of perforation, and the patient will complain of severe pain. The appearance of blood on paper points after instrumentation may be indicative of either strip perforation or apical perforation. However, it is unreliable as the bleeding may originate from the apical foramen also. Radiographs taken at different angles with radiopaque materials such as calcium hydroxide or barium sulfate or radiopaque instruments in the root canal are better option and may confirm the presence of perforation.

Electronic apex locators can accurately determine the location of perforations, making them more reliable than radiographs. A dental operating microscope is another effective tool in detecting perforation during orthograde root canal therapy and in surgical endodontic treatment.

## Measures to Prevent Perforation

The following few precautions can be followed to prevent the complication of perforation:

- Before root canal access, a pre-operative x-ray must be well studied regarding the number of canals, angulation of the root and curvature of the canals, the distance between the floor and roof of the pulp chamber, the and presence of calcifications. [6, 36-40]
- During access preparation, understanding the tooth anatomy and applying the basic principles such as the location of pulp chamber centrally at the level of CEJ and the canals are located at the junction of floor and walls of the pulp chamber and the location of canals directly under their respective cusps. [7, 41-44]
- During cleaning and shaping, pre-curving the files for curved canals prevents strip perforation and transportation. Accurate determination of working length using both radiographs and apex locators to prevent apical perforation.

## Management of Perforation

The successful management of root perforation depends on early diagnosis, choice of treatment, materials used, and the experience of the practitioner. The main aim is to achieve a tight and permanent seal that will prevent bacteria and its by-products in the root canal from entering the surrounding tissues. Perforations can be managed either by non-surgical or orthograde approach and surgical approach or intentional replantation.

### Non-surgical approach

Non-surgical or orthograde approach is done in cases of small, uncomplicated perforations that are easily accessible. Perforations present at the crown portion of the tooth can be repaired by conventional filling material such as GIC or composite. Those that are present subgingivally can be treated by endodontic-orthodontic approach by orthodontically extruding the tooth so that the perforation site becomes visible after which repair is attempted. However, this technique is not followed because it is time-consuming. Perforations present at the coronal-third of the root can be filled nonsurgically, provided it is accessed easily.

Ideal requirements of the root repair materials are as follows: [9]

1. It should provide adequate seal.
2. It should be biocompatible.
3. It should have the ability to produce osteogenesis and cementogenesis.
4. It should be bacteriostatic and radiopaque, non-toxic, and noncarcinogenic.

5. It should provide an absorbable matrix over which the sealing material may be condensed.

Conventionally, amalgam, EBA, gutta-percha, and cavite were used as root repair material. However, with the immense research on material science, newer materials are being constantly introduced. The newer materials include GIC, composite, dentin chips, decalcified freeze-dried bone, calcium phosphate, calcium hydroxide, Portland cement, MTA, Biodentine, endosequence, Bioaggregate, and new endodontic cement (NEC).

### Amalgam

In studies conducted by Mahmoud et al. [10] and Benenati et al., [11] amalgam had superior sealing properties of furcal perforation than cavite, calcium hydroxide, and gutta-percha.

### Super EBA

Super EBA is an alumina-reinforced zinc oxide-eugenol cement. It is easy to use, biocompatible, and it is highly adhesive and adapts well to the dentinal walls. Kenneth Weldon et al. showed that Super-EBA allowed significantly less microleakage than MTA at 24 h. [12, 45-49]

### Glass Ionomer Cement

It exhibits a greater sealing ability compared to the other materials used previously because of its chemically adhesive nature. Alhadainy and Himel found that light-cured glass ionomer cement has superior sealing ability compared to chemically cured glass ionomer cement. [13, 50, 51]

Fuss et al. had used metal modified glass ionomer cement in treating furcation perforations and concluded that leakage with metal modified GIC was significantly less than amalgam. Resin-modified glass ionomer cement can also be used as a repair material, and it is said to be superior to the conventional, chemically set glass ionomer cement and composite resin when used for perforation repair. [14, 52-54]

### MTA

It was introduced by Mahmoud Torabinejad in 1992. MTA consists of fine hydrophilic particles of tricalcium silicate, tricalcium aluminate, tricalcium oxide, silicate oxide, calcium sulfate dihydrate, tetracalcium aluminoferrite, and small amounts of mineral oxides. [15] It has a setting time of  $165 \pm 5$  min, evokes cementogenesis, and is biocompatible with the periradicular tissues. When amalgam, IRM, and mineral trioxide aggregate were tested for repair of experimentally created root perforations by Torabinejad et al., the results showed that the MTA had significantly less leakage than IRM or amalgam. [16]

### Biodentine

Biodentine is a calcium silicate-based bioactive material composed of tricalcium silicate, dicalcium silicate, calcium carbonate and oxide, iron oxide, zirconium oxide as powder, and liquid consisting of calcium chloride and hydrosoluble polymer.

It is easy to handle, has short setting time approximately 12 min and high alkaline pH, and is a biocompatible material. [17] Guner et al. compared biodentine and other perforation repair materials and that biodentine had considerable performance as a perforation repair material even after being exposed to various endodontic irrigants as compared to MTA. [18]

### Endosequence

Endosequence is a bioceramic material. It has an initial working time of 30 min and can set in the presence of moisture with a final working time of 4 h. It is referred to as bioceramic because it has a property of simulating tissue fluid and can precipitate hydroxyapatite crystals that increase in size with time. In a study by Jeevani et al., endosequence showed better sealing ability when compared to MTA and biodentine. [19]

### Bioaggregate

Bioaggregate is also a bioceramic material; it induces mineralized tissue formation and precipitation of apatite crystals that become larger with increasing immersion time. [20] A study by Hashem and Amin concluded that MTA is more influenced by acidic pH than bioaggregate when used as perforation repair material. [21]

### NEC

Bioactive material consisting of different calcium compounds are referred to as calcium-enriched mixture (CEM). It is composed of different calcium compounds in higher concentration, therefore causes

rapid precipitation of hydroxyapatite and this would make CEM cement preferable as a furcal perforation repair material in close proximity to the exposed periodontium. Asgary et al. observed cementogenesis and periodontal regeneration when CEM was used as perforation repair material.<sup>[22]</sup>

For larger perforations, internal matrix technique has been suggested.<sup>[23]</sup> This can be done only if there is adequate access and visibility, the site should be sterile. In this technique, a matrix material such as plaster of Paris, hydroxyapatite, calcium sulfate, and resinmodified glass ionomer is placed at the perforation site initially which serves as an artificial floor barrier over which the repair material is condensed to adequate thickness, and this is done to minimize the risk of pushing the material into the surrounding tissues.

### Surgical Approach

Surgical intervention is required in cases of large perforations that are not easily accessible, such as due to resorption, failure to heal after non-surgical repair. The main purpose of surgical treatment is to provide a tight seal against the entry of bacteria and its by-products into the periodontium. The success of the surgical repair depends on the following factors:<sup>[24]</sup>

- Amount of remaining bone
- Extent of osseous destruction
- Periodontal disease status
- Duration of perforation
- Soft tissue attachment level
- Patients oral hygiene status
- Operator's expertise.

Rud et al. suggested that the slightest bridge of crestal bone present should be preserved. Buccal perforations are easy to repair compared to lateral and lingual perforations due to its ease of access.

In case of surgical intervention, it is advisable to properly treat and fill the root canals before repair. The flap is elevated at the perforation site, and the repair material is packed directly on to the defect carefully without extruding it to the adjacent tissues, after which the flap is approximated. In cases of apical perforations, resection of the apical root to sound root structure with adequate filling is recommended. Crestal perforations are tricky to repair, as it will certainly result in loss of epithelial attachment and apical migration of tissues leading to pocket formation and secondary periodontal involvement. In such scenario's, guided tissue regeneration has been attempted as the GTR membrane acts as a barrier for apical migration of the epithelium.

### Intentional Replantation

Intentional replantation may be considered in surgically inaccessible perforations or if the defect is extensively large.<sup>[25]</sup> The tooth is nontraumatically extracted without extensive damage to cementum or periodontal tissue, repair done externally and immediately replanted with minimal extraoral time to the risk of external resorption. Teeth that are periodontally compromised and those with long, curved canals, and divergent roots are contraindicated for intentional replantation. Inflammatory resorption and ankylosis are the complications that may occur.

### Conclusion

The prognosis of perforation repair varies greatly depending on the above-discussed factors. The practitioner should have a well-updated knowledge and expertise to treat perforation. The successful management of root perforation depends on early diagnosis, choice of treatment, materials used, and the experience of the practitioner.

### References

1. Fuss Z, Trope M. Root perforations: Classification and treatment choices based on prognostic factors. *Endod Dent Traumatol* 1996; 12:255-64.
2. Seltzer S, Bender IB, Smith J, Freedman I, Nazimov H. Endodontic failures-an analysis based on clinical, roentgenographic, and histologic findings. II. *Oral Surg Oral Med Oral Pathol* 1967;23:517-30.
3. Tsesis I, Fuss Z. Diagnosis and treatment of accidental root perforations. *Endod Top* 2006; 13:95- 107.
4. Alhadainy HA. Root perforations. A review of literature. *Oral Surg Oral Med Oral Pathol* 1994; 78:368-74.
5. Fuss Z, Tsesis I, Lin S. Root resorption-diagnosis, classification and treatment choices based on stimulation factors. *Dent Traumatol* 2003; 19:175- 82.
6. Moreinis SA. Avoiding perforation during endodontic access. *J Am Dent Assoc* 1979; 98:707- 12.
7. Krasner P, Rankow HJ. Anatomy of the pulp- chamber floor. *J Endod* 2004; 30:5-16.
8. Peeso FA. The "ABC" of crown and bridge work. *Dent Cosmos* 1903; 45:274-9.
9. Oswald RJ. Procedural accidents and their repair. *Dent Clin North Am* 1979; 23:593-616.
10. El-Deeb ME, El-Deeb M, Tabibi A, Jensen JR. An evaluation of the use of amalgam, cavite, and calcium hydroxide in the repair of furcation perforations. *J Endod* 1982; 8:459-66.
11. Benenati FW, Roane JB, Biggs JT, Simon JH. Recall evaluation of iatrogenic root perforations repaired with amalgam and gutta-percha. *J Endod* 1986; 12:161-6.
12. Weldon JK, Pashley DH, Loushine RJ, Weller RN, Kimbrough WF. Sealing ability of mineral trioxide aggregate and super-EBA when used as furcation repair materials: A longitudinal study. *J Endod* 2002; 28:467-70.
13. Alhadainy HA, Himel VT. Comparative study of the sealing ability of lightcured versus chemically cured materials placed into furcation perforations. *Oral Surg Oral Med Oral Pathol* 1993; 76:338-42.
14. Himel VT, Alhadainy HA. Effect of dentin preparation and acid etching on the sealing ability of glass ionomer and composite resin when used to repair furcation perforations over plaster of Paris barriers. *J Endod* 1995; 21:142-5.
15. Torabinejad M, Hong CU, Pitt Ford TR, Kettering JD. Antibacterial effects of some root end filling materials. *J Endod* 1995; 21:403-6.
16. Torabinejad M, Hong CU, McDonald F, Pitt Ford TR. Physical and chemical properties of a new root- end filling material. *J Endod* 1995; 21:349-53.
17. Jitaru S, Hodisan I, Timis L, Lucian A, Bud M. The use of bioceramics in endodontics-literature review. *Clujul Med* 2016; 89:470-3.
18. Guneser MB, Akbulut MB, Eldeniz AU. Effect of various endodontic irrigants on the push-out bond strength of biodentine and conventional root perforation repair materials. *J Endod* 2013; 39:380-4.
19. Jeevani E, Jayaprakash T, Bolla N, Vemuri S, Sunil CR, Kalluru RS. Evaluation of sealing ability of MM- MTA, endosequence, and biodentine as furcation repair materials: UV spectrophotometric analysis. *J Conserv Dent* 2014; 17:340-3.
20. Shokouhinejad N, Nekoofar MH, Razmi H, Sajadi S, Davies TE, Saghi MA, et al. Bioactivity of endo sequence root repair material and bioaggregate. *Int Endod J* 2012; 45:1127-34.
21. Hashem AA, Amin SA. The effect of acidity on dislodgment resistance of mineral trioxide aggregate and bio-aggregate in furcation perforations: An in vitro comparative study. *J Endod* 2012; 38:245-9.
22. Asgary S, Moosavi SH, Yadegari Z, Shahriari S. Cytotoxic effect of MTA and CEM cement in human gingival fibroblast cells. Scanning electronic microscope evaluation. *N Y State Dent J* 2012; 78:51-4.
23. Lemon RR. Nonsurgical repair of perforation defects. Internal matrix concept. *Dent Clin North Am* 1992; 36:439-57.
24. Guttman JL, Harrison JW. *Surgical Endodontics*. Boston MA: Blackwell Scientific Publication; 1991. p. 409.

25. Grossman LI. Intentional replantation of teeth. *J Am Dent Assoc* 1966; 72:1111-8.
26. Ghavimi MA, Talesh KT, Ghoreishizadeh A, Amani M, Divband A. Autogenous bone with collagen material and surgical for maxillary sinus membrane elevation (sinus lift) in human subjects. *Biomedical & Pharmacology Journal*. Vol. 6(2), 249-252 (2013).
27. Yazdani J, Hajizadeh S, Ghavimi MA, Pourghasem Gargari B, Nourizadeh A, Kananizadeh Y. Evaluation of changes in anthropometric indexes due to intermaxillary fixation following facial fractures. *J Dent Res Dent Clin Dent Prospects*. 2016 Fall;10(4):247-250. doi: 10.15171/joddd.2016.039.
28. Yazdani J, Ghavimi MA, Jabbari Hagh E, Ahmadpour F. The Role of E-Cadherin as a Prognostic Biomarker in Head and Neck Squamous Carcinoma: A Systematic Review and Meta-Analysis. *Mol Diagn Ther*. 2018 Jul 13. doi: 10.1007/s40291-018-0351-y.
29. Seyyed Vahid Dehnad, Mohammad Ali Ghavimi, Arezu Ghoreishizadeh, Parisa Falsefi, Ramin Negahdari, Amir Reza Babalou. The Comparison of Job Stress among Dental Professionals of Dentistry Faculty Members of Tabriz University. Volume 8, Supplementary Issue 1, 2016. <https://doi.org/10.18311/aprh/2016/7714>.
30. Samiei M, Shahi S, Abdollahi AA, Eskandarinezhad M, Negahdari R, Pakseresht Z. The Antibacterial Efficacy of Photo-Activated Disinfection, Chlorhexidine and Sodium Hypochlorite in Infected Root Canals: An in Vitro Study. *Iran Endod J*. 2016 Summer;11(3):179-83. doi: 10.7508/iej.2016.03.006.
31. Fakhrzadeh V, Gaffari T, Hamedirad F, Negahdari R, Nasrollah A. P, Eslami H. Influence of Preparation Design and Restorative Material to Marginal Discrepancy of Ceramic Laminate Veneers. *Biomed Pharmacol J* 2015;8(2).
32. Fakhrzadeh V, Eslami H, Nasrollah A P, Negahdari R, Falsafi P, Sabeti M. The Comparison of Dimensional Accuracy of Modified Two-Stage Putty-Wash Condensation S Polyvinyl Molding Method and Distance Two-Stage Putty-Wash Molding Method. *International Journal of Clinical Dentistry*. 2016, Vol. 9 Issue 4, p265-271. 7p.
33. Pournasrollah A, Fakhrzade V, Eslami H, Negahdari R. Evaluation of shear bond strength between direct composite and zirconia. 2015. *Der Pharmacia Lettre*. <http://localhost:8080/xmlui/handle/123456789/13525>.
34. Negahdari R, Rahbar M, Fakhrzadeh V, Eslami H, Akbari T, Bohluli S. Comparison of Proinflammatory Cytokine Levels in Gingival Crevicular Fluid around Dental Implants with Ceramic and Titanium Abutments. *J Contemp Dent Pract*. 2017 Sep 1;18(9):831-836.
35. Khosravi A, Mousavi-Moghadam R, Zarei S. Evaluation of medical ethics among patients with nose and abdominal surgery. *Life Science Journal*. Volume 9, Issue 4, 25 December 2012, Pages 5778-5781.
36. Mousavimoghadam SR, Hamedheydari S, Abbasalipour S. Examine the relationship between spiritual intelligence of employed mothers, teachers elementary with mental health of children aged 6-12 years in ILAM. *Der Pharmacia Lettre*. Volume 7, Issue 10, 2015, Pages 39-43.
37. Musavimoghadam SR, Haji S, Chegeni ME, Veiskarami M. Examine the relationship between religious orientation, intelligence, ethical, and mental health among students Azad University of ILAM. *Der Pharmacia Lettre*. Volume 7, Issue 7, 2015, Pages 182-186.
38. Musavimoghadam SR, Sabzeghabaei EA, Ebrahimpour M. Relationship between spiritual understanding and meaningful life with psychological distress in mothers of children exceptional. *International Journal of Tropical Medicine*. Volume 11, Issue 1, 2016, Pages 5-8.
39. Musavimoghadam SR, Mansouri Akbari. Examining the relationship of belief in God, self-esteem and quality of mate selection in Ilam young girls. *Research Journal of Applied Sciences*. Volume 11, Issue 3, 2016, Pages 72-76.
40. Mahmoudi T, Karimi K, Arkani M, Farahani H, Nobakht H, Dabiri R, Asadi A, et al. Lack of associations between vitamin D metabolism-related gene variants and risk of colorectal cancer. (2014) *Asian Pacific Journal of Cancer Prevention*, 15 (2), pp. 957-961.
41. Darvishi M. Antibiotic resistance pattern of uropathogenic methicillin-resistant staphylococcus aureus isolated from immunosuppressive patients with pyelonephritis. (2016) *Journal of Pure and Applied Microbiology*, 10 (4), pp. 2663-2667.
42. Darvishi M. Virulence factors profile and antimicrobial resistance of acinetobacter baumannii strains isolated from various infections recovered from immunosuppressive patients. (2016) *Biomedical and Pharmacology Journal*, 9 (3), pp. 1057-1062.
43. Sanayei A, Rajabion L. Critical successful factors contributing to e-commerce adoption among Iranian SMEs. (2009) *International Journal of Information Science and Management*, 7 (2), pp. 57-65.
44. Siyahi V, Habibi-Yangjeh A, Latifi-Navid S, Asadi A. Microwave-Assisted One-Pot preparation of AgBr/ZnO nanocomposites as highly efficient Visible-Light photocatalyst for inactivation of Escherichia coli. (2015) *Materials Express*, 5 (3), pp. 201-210.
45. Ravaji B, MashadizadeS, Hashemi A. Introducing optimized validated meshing system for wellbore stability analysis using 3D finite element method. *Journal of Natural Gas Science and Engineering*. Volume 53, May 2018, Pages 74-82.
46. Sotoudeh A, Jahanshahi A, Zareiy S, Darvishi M, Roodbari N, Bazzazan A. The influence of low-level laser irradiation on spinal cord injuries following ischemia-reperfusion in rats. *Acta Cir Bras*. 2015 Sep;30(9):611-6.
47. Rajabion L, Kheirabadi A. Information technology: It's role in facilitating the sharing and transfer of knowledge in small medium sized enterprises. (2011) *KMIS 2011 - Proceedings of the International Conference on Knowledge Management and Information Sharing*, pp. 275-282.
48. Tavan H, Sayehmiri K, Taghinejad H, Mousavi Moghadam SR. Factor Analysis of Spiritual Health on the Islam Viewpoint. *Iran J Public Health*. 2015 Nov;44(11):1572-3.
49. Mousavimughaddam SR. Healthy man and its role in the spiritual health and mental health from the viewpoints of rene descartes and allamah tabatabaei based on the interpretative attitude of almizan. 2014, *Journal of Zanjan University of Medical Sciences and Health Services*. 22(90), pp. 33-44
50. Negahdary M, Habibi-Tamijani A, Asadi A, Ayati S. Synthesis of zirconia nanoparticles and their ameliorative roles as additives concrete structures. (2013) *Journal of Chemistry*, art. no. 314862.
51. Mahmoudi T, Majidzadeh-A K, Farahani H, Mirakhorli M, Dabiri R, Nobakht H, Asadi A. Association of Vitamin D receptor gene variants with polycystic ovary syndrome: A case control study. (2015) *Iranian Journal of Reproductive Medicine*, 13 (12), pp. 793-800.
52. azar A, Shamsazar F, Asadi A, Rezaei-zarchi S. A glucose biosensor based on glucose oxidase enzyme and ZnO nanoparticles modified carbon paste electrode. (2016) *International Journal of Electrochemical Science*, 11 (12), pp. 9891-9901.
53. Bahador M, Keshtkar MM, Zariie A. Numerical and experimental investigation on the breast cancer tumour parameters by inverse heat transfer method using genetic algorithm and image processing. 2018, *Sadhana - Academy Proceedings in Engineering Sciences*. 43(9),142
54. Keshtkar MM, Khani AG. Exergoeconomic analysis and optimization of a hybrid system based on multi-objective generation system in Iran: a case study. 2018, *Renewable Energy Focus*. 27, pp. 1-13.