

Low-Level laser Diode on post modified Radical Mastectomy Lymphedema: a randomized controlled trial

Hussein Gamal Hussein Mogahed^{1*}, Manar Mohamed Badawy², Nesma Morgan Abdel Aziz¹

¹ Lecturer in the Department of Physical Therapy for Surgery, Faculty of Physical Therapy, Cairo University, Giza, Egypt. ² Lecturer of Physical Therapy for Cardiovascular, Pulmonary Disorders, and Geriatrics, Faculty of Physical Therapy, Misr University for Science and Technology, Giza, Egypt.

Correspondence: Hussein Gamal Hussein Mogahed; Lecturer in the Department of Physical Therapy for Surgery, Faculty of Physical Therapy, Cairo University, Giza, Egypt. Email: hmogahed85@gmail.com.

ABSTRACT

Objective: This work aimed to evaluate the effect of low-level laser diode therapy on post-mastectomy lymphedema and pain. **Methods:** Thirty female patients suffering from unilateral post-mastectomy shoulder lymphedema and pain were randomly divided into two equal groups. Group (A) received low-level laser diode therapy plus a traditional physical therapy program. Group (B) received placebo low-level laser diode therapy plus a traditional physical therapy program. Treatment sessions were three times per week for 3 months for each group. Volumetric measurement and visual analog scale were used to assess post-mastectomy lymphedema and pain respectively. Outcome measures were evaluated before and after the treatment period. **Results:** Lymphedema volume was decreased significantly in the laser group more than the placebo group (-108.4 ml) mean difference by volumetric measurement. The pain was decreased significantly in the laser group more than the placebo group (-0.1333 degree) mean difference by visual analog scale. **Conclusion:** Low-level laser diode therapy had a valuable effect on the treatment of post-mastectomy lymphedema and pain.

Keywords: Low-level laser diode therapy, Mastectomy, Lymphedema, volumetric measurement, and visual analogue scale

Introduction

Breast cancer is very rare before the age of 25. It reaches a high incidence in the decade between 40 to 50 and a continuous increase into old age. The most common cancers are lung cancer in men and breast cancer in women [1].

Cancer rehabilitation is a dynamic process directed towards enabling a patient to function at the maximum level of disease in terms of their physical, mental and emotional potential. Four major problems presented during treatment include post-

mastectomy lymphedema, pain, limited motion of the shoulder joint and psychological problems [2].

Tumor excision and radiotherapy may develop an obstruction in the lymphatic system of axilla that causes post-mastectomy lymphedema (PML). PML is an abnormal accumulation of proteins and interstitial rich fluid and an increase in capillary permeability. It can affect one or both arms, the head, and the neck. Swelling can worsen and become severe. Skin sores or other problems can develop [3-5].

Ordinary management of PML includes massage, complete decongestive lymphatic therapy, compression therapy, elevation, and intermittent pneumatic compression. They are expensive and time-consuming [6].

Recently, low-level laser therapy (LLLT) appeared to be a good choice for the treatment of some cases such as a hypertrophic scar, immunity diseases and PML [7-9].

Low-level laser diode helps lymphangiogenesis, enhancement of lymphatic drainage, stimulation of macrophage cells and facilitation of the immune system so, it has been used for the management of PML [10].

Access this article online

Website: www.japer.in

E-ISSN: 2249-3379

How to cite this article: Hussein Gamal Hussein Mogahed, Manar Mohamed Badawy, Nesma Morgan Abdel Aziz. Low-Level Laser Diode on Post Modified Radical Mastectomy Lymphedema: A Randomized Controlled Trial. J Adv Pharm Educ Res. 2020;10(4):105-9.
Source of Support: Nil, Conflict of Interest: No.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Low-level laser diode enhances the phagocytic and chemotactic activity of human leukocytes in vitro and it decreases infection risk which is considered one of its advantages ^[11, 12].

No study has illustrated that if any resolving of the PML symptoms would improve upper limb function. In our study low-level diode laser was administered only to the axilla because of breast cancer surgeries, radiotherapy and/or chemotherapy makes the blockage of lymphatic drainage causing PML symptoms ^[13].

There was a novelty in our study in solving post-mastectomy serious problems such as limitation of upper limb range of motion, pain, heaviness, and psychological morbidity because of the fewness of evidence-based studies that improve those critical symptoms.

Materials and Methods

Thirty female patients who had secondary (stage II, III) lymphedema in the unilateral upper limb after modified radical mastectomy with axillary lymph node dissection with or without radiotherapy intervention were included in the study. Lymphedema was diagnosed by isotopic lymphangiography. They were recruited from the National Cancer Institute. Their ages ranged from 40 to 55 years. Before the study procedures, they were examined carefully by the physician. This research was approved by the research ethics committee of the Faculty of Physical Therapy (No:P.T.REC/012/001156). The followed procedures complied with the Helsinki declaration of 1975.

Patients were excluded if they had distant metastasis, active infection, bilateral upper limb lymphedema, kidney problems, heart diseases, and diabetes. This study was a prospective, single-blinded, controlled clinical trial. The patients were randomly assigned to two equal groups (A and B). Group (A) (study group): included 15 patients who received low-level laser diode therapy plus traditional physical therapy program (manual lymphatic drainage, circulatory exercises with elevation, shoulder ROM exercises and pneumatic compression therapy) 3times/week/12 weeks. Group (B) (Control group): included 15 patients who received placebo laser therapy and traditional physical therapy program (manual lymphatic drainage, circulatory exercises with elevation, shoulder ROM exercises and pneumatic compression therapy) 3 times/week/12 weeks.

Ethical Consideration

All assessment and treatment procedures were explained in detail to patients and informed consent was signed from each patient before starting.

Measurements

Patients' demographic data such as age, weight, dominant arm, type of breast surgery done and the onset of PML arm volume change were written on the evaluation sheet of every patient.

The measurement procedures were started before starting treatment (first record) and at the end of the treatment after 3 months (second record).

Lymphedema Volume Measurement Procedure

Volumetric lymphedema measurement (VLM) was done before starting treatment (first record) and at the end of treatment (second record) by the physical therapist. The graded container was initially full but, not overflowing to a level of drain spot with water. Each patient was put in a standing position for volume measurement and asked to remove jewelry (if present). The affected upper limb was lowered into a container with an extended elbow. The affected upper limb was removed and dried, the graduated baker was used to detect displaced water volume by immersing upper limb and data were recorded. We repeated the previous steps with the sound upper limb, then comparisons between affected and sound limbs results ^[14].

Shoulder Pain Measurement Procedure

Pain assessment was done by visual analog scale (VAS) that graded from 0 to 10 score, before starting treatment (first record) and at the end of treatment (second record) ^[15].

Laser Treatment Procedure

A laser unit (Bravo Terza Serie, Model D; ASA S.r.l., Vicenza, Italy) with a scanning head was used. The unit comprises infrared laser head sources emitting at a wavelength of 905 nm. The average output was 24 mW, each with pulsed emission at a frequency varying from 1 to 10,000 Hz. A dose of 2 J/cm² was chosen because it lies within the therapeutic window of laser dose stated by the Arndt-Schultz Law. The scanner laser unit was used in our study because it can penetrate subcutaneously and into deeper tissues without elevation of tissue temperature^[7, 8].

All patients were placed in supine positions and LLLT was applied perpendicularly to the axillary region of the affected side. The plane of the emission of a laser scanner was parallel to the treatment area. The distance between the plane of a laser scanner and the treatment area was 50 cm. The aiming beam helped to locate the treatment region and to ensure the whole axillary region was covered.

Data Analysis

The Statistical Package for the Social Sciences (SPSS) version 23 (IBM Corp, New York, United States) was used for data analysis, and it was performed by an independent investigator. The paired *t*-test was calculated on the pre-test to post-test change within each group. Finally, to assess whether any difference that exists in the post-test scores, an independent *t*-test was calculated on the post-test change in both groups. The level of significance $p < 0.05$ was used.

Results

The descriptive characteristics of both groups are shown in (Table 1). There were no statistical differences between both groups regarding age.

Table 1. Descriptive statistics for age in both groups.

	Group (A)	Group (B)
Mean of age (years)	48.4	48.33
SD	4.0848	4.1346
Standard error	1.0547	1.0676
Mean difference	0.066667	
t. value	-0.07	
p. value	0.948	
Level of significance	Non-significant	

Comparison Between Pre-treatment and Post-treatment within The same Group

• Results of Lymphedema Volume

The mean value of the lymphedema volume of the first record in group A was 357.3 ± 74.0 ml. and the value of the second record was 248.7 ± 49.4 ml. The paired *t*-test demonstrated a statistically significant difference between pre-treatment and post-treatment for lymphedema volume in this group ($t = -15.35$, $p = 0.0001$).

The mean value of the lymphedema volume of the first record in group B was 357.2 ± 74.3 ml. and the value of the second record was 357.1 ± 79.8 ml. The paired *t*-test demonstrated a statistically no significant difference between pre-test and post-test for lymphedema volume in this group ($t = -0.01$, $p = 0.993$), Table (2), Fig. (1).

Table 2. Comparison of lymphedema volume (ml) pre-treatment and post-treatment in both groups.

	Group (A)		Group (B)	
	Pre	Post	Pre	Post
Mean (ml)	357.333	284.667	357.2	357.133
± SD	74.038	49.442	74.252	79.819
Standard error	19.117	12.766	19.172	20.609
Mean difference	-108.667		-0.066667	
t. value	-15.35		-0.01	
p. value	0.0001		0.993	
Level of significance	Significant		Non-significant	




Figure 1. Comparison of lymphedema volume (ml) pre-treatment and post-treatment in both groups.

• Results of Shoulder Pain

As observed in the table (3) and figure (2): There were non-significant differences in the first records of VAS, between study and control groups ($P > 0.05$).

As observed in the table (4) and figure (2): There was a significant decrease in the second records of VAS, between study and control groups ($P < 0.0001$).

Table 3. The first (pre-treatment) records of VAS in both groups.

	Group (A)	Group (B)
Mean in degrees	9	9.133
± Standard deviation	0.926	0.3519
Standard error	0.239	0.0909
Mean difference	- 0.1333	
t. value	- 0.52	
p. value	0.609	
Level of significance	Non-significant	

Table 4. The second (post-treatment) records of VAS in both groups.

	Group (A)	Group (B)
Mean in degrees	2.333	8.933
± Standard deviation	0.976	0.458
Standard error	0.252	0.118
Mean difference	- 6.6	
t. value	-23.71	
p. value	0.0001	
Level of significance	Significant decrease	




Figure 2. Mean values of VAS (in degrees) of pre-treatment and post-treatment records of both groups.

Discussion

This study illustrated that low-level laser diode therapy was significantly beneficial in reducing upper limb volume and decreasing shoulder pain as a post-mastectomy complication.

The results were agreed with Pillar and the lander, who found that with using (LLLT), there was a reduction in the amount of lymphedema and volume of the extracellular fluid through measuring by bioimpedance. Patients had an improvement in pain, tightness, and heaviness in addition to upper limb motion. 19.7% losing of arm lymphedema in the first 16 sessions, and about 7% loss over 6 months^[8].

Also, Thelander^[16], who investigated the effects of (LLLT) with a wavelength of 632.8 nm, an output intensity of 9 mW, and an energy density of 2.4 J/cm² on lymphedema with beneficial effects on lymphedema volume and shoulder mobility.

Carati *et al.*^[13], found that after three months of treatment, there was a 90 ml reduction in upper limb volume with better life quality in the group that administrated to active LLLT.

All studies used the infrared range of light through fluencies ranging from 1.5 to 2 J/cm². Wavelengths located in the near-infrared spectral range are the best recommended in treating more deeply structures because of their greater tissue penetration^[17].

Low-level laser diode enhances lymphatic drainage^[18], reduces pain^[19] and increases defense cell activity^[20]. Its effects on lymphatic system pathologies are the stimulation of fibroblasts, lymphocytes, and macrophages^[13], forming new vessels, re-establishing drainage, decreasing fibrosis and hypertrophic scars in armpit region and enhancing lymphatic vessel contractility^[21].

Karu^[22], found that the therapeutic effects of low-level laser diode therapy are due to primary and secondary mechanisms. The primary mechanisms (formation of singlet oxygen, modifications in the redox properties of CuA in addition to CuB or heme and also heme a₃ centers and improved superoxide anion production) happen during direct irradiation on the photoreceptors. Secondary mechanisms are responsible for the connection between a reaction to light through photoreceptors in mitochondria in addition to DNA and RNA synthesis actions located in the nucleus^[23].

In our study, the results of low-level laser diode therapy on lymphedema were examined during and after three months. One possible reason for that is re-establishing the lymphatic vessel network function. However, further studies are needed to confirm this hypothesis.

Limitations

Small sample size.

Conclusion

We found out that a combination of low-level laser diode therapy plus traditional physical therapy program had a significant effect on lymphedema reduction and decreasing shoulder pain. This combination is a novel and cheap treatment to resolve one of the most serious post modified radical mastectomy problems.

Conflict of Interest

The authors certify that they have NO affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers' bureaus; membership, employment, consultancies, stock ownership, or other equity interest; and expert testimony or patent-licensing arrangements), or non-financial interest (such as personal or professional relationships, affiliations, knowledge or beliefs) in the subject matter or materials discussed in this manuscript.

References

1. Aaronso, N.J. Immune System and cancer rehabilitation . American Fitness. 2000; 2(87):300-10.
2. Abeloff, M.D., Lichter, A.S. Breast Clinical Oncology. 3rd Ed. Churchill Livingstone Inc. 2009; 131-51.
3. Tobin MB, Lacey HJ, Meyer L, Mortimer PS. The psychological morbidity of breast cancer-related arm swelling. Psychological morbidity of lymphoedema. Cancer. 1993 Dec 1;72(11):3248-52. doi.org/10.1002/1097-0142(19931201)72:11<3248::AID-CNCR2820721119>3.0.CO;2-Z
4. Passik SD, Newman ML, Brennan M, Tunkel R. Predictors of psychological distress, sexual dysfunction and physical functioning among women with upper extremity lymphedema related to breast cancer. Psycho-oncology. 1995 Dec;4(4):255-63. doi.org/10.1002/pon.2960040402
5. Passik SD, McDonald MV. Psychosocial aspects of upper extremity lymphedema in women treated for breast carcinoma. Cancer: Interdisciplinary International Journal of the American Cancer Society. 1998 Dec 15;83(S12B):2817-20.
6. Casley-Smith J, Casley-Smith JR. The cost efficiencies of various treatments for lymphoedema. Modern Treatment for Lymphoedema. Adelaide, Australia: The Lymphoedema Association of Australia. 1997:296-304.
7. Piller NB, Thelander A. Treating chronic post-mastectomy lymphoedema with low level laser therapy: a cost effective strategy to reduce severity and improve the quality of survival. Laser therapy. 1995;7(4):163-8.
8. Piller NB, Thelander A. Treatment of chronic postmastectomy lymphedema with low level laser therapy: a 2.5 year follow-up. Lymphology. 1998 Aug 26;31(2):74-86.
9. Kaviani A, Fateh M, Nooraie RY, Alinagi-zadeh MR, Ataie-Fashtami L. Low-level laser therapy in management of postmastectomy lymphedema. Lasers in Medical Science. 2006 Jul 1;21(2):90-4.
10. Lievens P. The influence of laser-irradiation on the motricity of the lymphatical system and on the wound healing process. In International congress on laser in medicine and surgery 1985 Jun 26 (pp. 26-28).

11. Osanai T, Shiroto C, Mikami Y, Kudou E, Komatsu T, Suzuki K, Nakaji S, Kumae T, Sugawara K, Sasaki M. Measurement of GaAs Diode Laser Action on Phagocytic Activity of Human Neutrophils as a Possible Therapeutic Dosimetry Determinant. *Laser Therapy*. 1990;2(3):123-33.
12. Tadakuma T. Possible application of the laser in immunobiology. *The Keio journal of medicine*. 1993;42(4):180-2.
13. Carati CJ, Anderson SN, Gannon BJ, Piller NB. Treatment of postmastectomy lymphedema with low-level laser therapy: a double blind, placebo-controlled trial. *Cancer: Interdisciplinary International Journal of the American Cancer Society*. 2003 Sep 15;98(6):1114-22.
14. Brennan, M.J. Garden FH. Post mastectomy shoulder pain and lymphedema. *Arch Phys Med Rehabil*. 2006; (10):55- 67.
15. Cleary, JF. Management of pain in cancer of the breast. Chap 42 , Philadelphia. 2002; 905.
16. Thelander, A. Lymphedema and laser therapy. In: 4th World congress of laser therapy. Japan; 2003.
17. Ryan T, Smith RK. An investigation into the depth of penetration of low level laser therapy through the equine tendon in vivo. *Irish veterinary journal*. 2007 Dec;60(5):295.
18. Lievens PC. The effect of a combined HeNe and IR laser treatment on the regeneration of the lymphatic system during the process of wound healing. *Lasers in medical science*. 1991 Jun 1;6(2):193-9.
19. Hopkins JT, McLoda TA, Seegmiller JG, Baxter GD. Low-level laser therapy facilitates superficial wound healing in humans: a triple-blind, sham-controlled study. *Journal of athletic training*. 2004 Jul;39(3):223.
20. Posten W, Wrona DA, Dover JS, Arndt KA, Silapunt S, Alam M. Low-level laser therapy for wound healing: mechanism and efficacy. *Dermatologic surgery*. 2005 Mar;31(3):334-40.
21. Dirican A, Andacoglu O, Johnson R, McGuire K, Mager L, Soran A. The short-term effects of low-level laser therapy in the management of breast-cancer-related lymphedema. *Supportive care in cancer*. 2011 May 1;19(5):685-90.
22. Karu T. Photobiological fundamentals of low-power laser therapy. *IEEE Journal of Quantum Electronics*. 1987 Oct;23(10):1703-17.
23. Karu T. Primary and secondary mechanisms of action of visible to near-IR radiation on cells. *Journal of Photochemistry and photobiology B: Biology*. 1999 Mar 1;49(1):1-7.