

Anti-diabetic activity of cassia fistula (alpha amylase – inhibitory effect)

Hena Mariam Fathima¹, Lakshmi Thangavelu^{2*}, Anitha Roy²

¹Under Graduate Student, Saveetha Dental College, SIMATS, Saveetha University, Chennai, India, ²Associate Professor, Department of Pharmacology, Saveetha Dental College, SIMATS, Saveetha University, Chennai, India.

Correspondence: Lakshmi Thangavelu, Associate Professor, Department of Pharmacology, Saveetha Dental College, SIMATS, Saveetha University, Chennai, India. E_mail: lakshmi085@gmail.com

ABSTRACT

Aim: To evaluate the anti-diabetic activity of cassia fistula. **Materials and methods:** Cassia fistula extract used in the study here was obtained from Green Chem Herbal extracts and formulations, Bengaluru, India. DMEM, FBS, and Pioglitazone, were obtained from Sigma Aldridge co India. Acarbose, dinitrosalicylic acid, DMSO is obtained from Sigma Aldrich co. India. Alpha amylase enzymes were procured from Himedia, Mumbai. All the reagent chemicals used in the study were up to analytical grade. The study was carried out as an in vitro study by the method of Bernfeld. **Results:** cassia fistula showed high alpha amylase inhibitory activity. **Conclusion:** It can be concluded that cassia fistula shows high alpha amylase inhibitory activity and can be used as medication for diabetes.

Keywords: Alpha glycosidase, anti-diabetic, cassia fistula, diabetes mellitus, hypoglycaemia.

Introduction

Diabetes mellitus, a chronic metabolic disorder, has now become epidemic with a worldwide incidence of 5% in the general population. More than 100 million of the world's population has already reached the diabetic mark and the number of people suffering from it is expected to soar up to 366 million [1]. Decreased physical activity, increasing obesity and stress, and changes in the food consumption have been implicated in this increasing prevalence in the past two decades [2]. Overt diabetes affects 2-3% of the total world population. Type 1 and type 2 diabetes are treated with exogenous insulin and oral hypoglycemic agents; respectively, in conventional therapy [3]. The demand for using natural products with antidiabetic activities by patients has increased, although there are different kinds of oral hypoglycemic agents available along with insulin for the treatment of diabetes [4-9]. The patients with noninsulin-requiring diabetes have been treated orally in traditional medicine with a variety of plant extracts for many

years. In India, in ancient literature (Ayurveda), a number of plants are stated for the treatment of diabetic conditions.

Many traditional healers applied flowers of *Cassia fistula* Linn (Indian laburnum) belonging to the family Leguminosae widely in the most of the herbal preparations for diabetes [5,10-16]. Traditionally, it is used to cure diabetes, constipation, skin diseases, fever, abdominal pain, and leprosy [12,17,18,19]. This plant is reported to contain antibacterial, antioxidant, and antiinflammatory activities [7,20-23]. The constituents reported in this plant are alkaloids [8,24,25], flavonoids, and anthraquinone glycosides [9,26-30].

The traditional use of the flowers of *C. fistula* for diabetes has not been proven so far. Moreover, the main focus has been put on ethanol and aqueous extracts for diabetes by the researchers, but a considerable number of investigations have declared that the petroleum ether, benzene, and chloroform extracts were also found to have anti-diabetes activities [9-11]. Being aware of the effective extracts and isolating the active fraction from the effective extract would be a helpful method for developing new drugs. The standard fraction of an active extract may demonstrate to be better therapeutically, less toxic and inexpensive compared to the pure isolated compounds. Considering these facts, proving its traditional use, identifying the active antidiabetic extract of *C. fistula* in diabetes-associated complications, and identifying the active antidiabetic fraction of the active extract are the aims of the present study.

Access this article online

Website: www.japer.in

E-ISSN: 2249-3379

How to cite this article: Hena Mariam Fathima, Lakshmi Thangavelu, Anitha Roy. Anti-diabetic activity of cassia fistula (alpha amylase – inhibitory effect). *J Adv Pharm Edu Res* 2018;8(2):12-15.

Source of Support: Nil, Conflict of Interest: None declared.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Patients and Methods

Plant material:

Cassia fistula extract used in the study here was obtained from Green Chem Herbal extracts and formulations, Bengaluru, India.

Chemicals used:

DMEM, FBS, and Pioglitazone, were obtained from Sigma Aldridge co India. Acarbose, dinitrosalicylic acid, DMSO is obtained from Sigma Aldrich co. India. Alpha amylase enzymes were procured from Himedia, Mumbai. All the roger chemicals used in the study were up to analytical grade.

In – vitro α - amylase inhibitory assay (Bernfeld 1955)

In vitro amylase inhibition was studied by the method of Bernfeld^[30]. In brief, 100 μ L of different concentrations of Cassia (1, 3, 10, 30, 100, 300 and 1000 μ g/ml) were allowed to react with 200 μ L of α -amylase enzyme (Hi media RM 638) and 100 μ L of 2mM of phosphate buffer (pH-6.9). After 20-minute incubation, 100 μ L of 1% starch solution was added. 200 μ L of the enzyme was replaced by buffer to apply the same process for the controls. The reference standard was Acarbose. After incubation for 5 minutes, 500 μ L of Dinitrosalicylic acid reagent was added to both the control and the test. They were maintained in boiling water bath for 5 min. The amount of absorbance was recorded at 540 nm using spectrophotometer and the percentage inhibition of α -amylase enzyme was computed using the formula


$$\% \text{ inhibition} = [(Control - Test)/Control] * 100$$

Suitable reagent blank and inhibitor controls were simultaneously applied.

Results

α – amylase inhibitory activity

The results of the present study exhibited strong α – amylase inhibitory activity of the Cassia extract which is compared with the standard Acarbose. A maximum inhibition of 88.65% was observed at a concentration of 1000 μ g/ml by Cassia extract, which was compared with that of standard Acarbose that showed inhibition of about 93%. The IC50 of the extract was found to be 73.16 μ g/ml and for Acarbose 34.83 μ M.



Discussion

Cassia fistula Linn. also known as golden shower, Indian laburnum, is categorized in the family of Leguminosae. It is used in the treatment of hematemesis, pruritis, intestinal disorders, leucoderma, diabetes, and as antipyretic, analgesic and laxative in folk medicine. The size of Cassia fistula is moderate. And there are rich sources of flavonoids, anthraquinones and polysaccharides in various species of Cassia. The existence of flavonol and xanthone glycosides has been already reported in the bark of the plant Cassia fistula. So, isolating the phytoconstituent responsible for alpha amylase inhibition has been attempted to be done^[31].

Hasenah Ali et al.^[32] conducted a research regarding α -Amylase inhibitory activity of some Malaysian plants which are used to treat diabetes, with particular focus on Phyllanthus amarus. Six Malaysian plants' extracts which had a reputation of being useful in treating diabetes, were selected and examined for α -amylase inhibition using an in vitro model. Two different protocols (with and without pre-incubation) were used to study their inhibitory activities, and the results showed that Phyllanthus amarus hexane extract had α -amylase inhibitory properties. In the test without pre-incubation, hexane and dichloromethane extracts of Anacardium occidentale, Lagerstroemia speciosa, Averrhoa bilimbi Pithecellobium jiringa and Parkia speciosa were not active. The isolation of dotriacontanyl docosanoate, triacontanol and a mixture of oleanolic acid and ursolic acid were the results of extraction and fractionation of Phyllanthus amarus hexane extract. Dotriacontanyl docosanoate and the mixture of oleanolic acid and ursolic acid are reported from this plant species for the first time. All compounds were tested in the α -amylase inhibition assay and the results demonstrated that the oleanolic acid and ursolic acid (2:1) mixture was a potent α -amylase inhibitor with $IC_{50} = 2.01 \mu$ g/ml (4.41 μ M) which is significantly related to the α -amylase inhibition activity of the extract. Three pure pentacyclic triterpenoids, oleanolic acid, ursolic acid and lupeol demonstrated to inhibit α -amylase.

A research conducted by^[33] on in vitro α - amylase and α -glucosidase inhibitory activities of the ethanolic extracts of dioscorea villosa tubers demonstrated that the α – amylase inhibitory activity also showed a dose - dependent increase ranging from 10.71 ± 0.00 to 71.88 ± 3.77 percent with a concentration range of 1.5 μ g/ml - 1000 μ g/ml and the IC50 value was computed as 72.44 μ g / ml when compared with acarbose, IC50 value $83.23 \pm 0.39 \mu$ g/mL.

A study was conducted by^[34] on a comparative study of alpha amylase inhibitory activities of common anti-diabetic plants at Kharagpur 1 block. Acalypha indica, Allium cepa, Allium sativum, Azadirachta indica, Musa sapientum, Mangifera indica, Murraya, Ocimum sanctum, Phyllanthus amarus and Tinospora cordifolia were examined for their alpha amylase inhibitory activities to establish anti-diabetic potentials. The plant extracts were prepared sequentially with petroleum ether, hexane, chloroform, ethanol and aqueous. The obtained extracts were subjected to in vitro alpha amylase inhibitory assay using starch azure as a substrate and porcine pancreatic amylase as the enzyme. Statistical difference and linear regression analysis were performed by using Graphpad prism 5 statistical software. Ethanol extracts of Mangifera indica, Azadirachta indica and petroleum ether extract of Murraya koenigii (at a concentrations 10-100 μ g/ml) showed maximum percentage inhibition on alpha amylase activity with an IC50 value of $37.86 \pm 0.32 \mu$ g/ml,

62.99 ± 1.20 µg/ml and 59.0 ± 0.51 µg/ml; respectively, when compared with acarbose (IC₅₀ value 83.33 ± 0.75 µg/ml). The results showed that *Mangifera indica*, *Azadirachta indica* and *Murraya koenigii* might be effective in lowering post prandial hyperglycemia.

A research done by G. Nivetha et al.^[35] on Comparative Evaluation of Anti-Diabetic Activity of Lemon Grass Oil and Tulasi Oil demonstrated that the lemon grass oil and Tulasi oil have anti hypoglycaemic properties. It showed that these plants have hypoglycaemic effects. Use of lemon grass oil and Tulasi oil on a regular basis by diabetic patients can be beneficial in lowering the blood glucose and potentiates other therapies used in diabetes treatment.

A study conducted by^[36] on the topic in vitro alpha-amylase inhibition and in vivo antioxidant potential of *Amaranthus spinosus* in alloxan-induced oxidative stress in diabetic rats says that *Amaranthus spinosus* Linn. (Amaranthaceae), commonly called as “Mulluharivesoppu” in Kannada, is applied in traditional system of medicine for the treatment of diabetes in India. The current study deals with the scientific evaluation of alpha amylase and the antioxidant potential of methanol extract of *A. spinosus* (MEAS). The goal of this study was to examine in vitro alphaamylase enzyme inhibition by CNPG3 (2-chloro-4-nitrophenol a-D-maltotrioxide) and in vivo antioxidant potential of malondialdehyde (MDA), glutathione (GSH), catalase (CAT) and total thiols (TT) in alloxan-induced diabetic rats of a methanolic extract of *A. spinosus*. Blood sugar was also determined in MEAS-treated alloxan-induced diabetic rats. MEAS showed significant inhibition of alpha-amylase activity and IC₅₀ 46.02 lg/ml. Oral administration of MEAS (200 and 400 mg/kg) for 15 days showed significant reduction in the elevated blood glucose, MDA and restores GSH, CAT and TT levels as compared with a diabetic control. The present study provides evidence that the methanolic extract of *A. spinosus* has potent alpha amylase, anti-diabetic and antioxidant activities.

A research accomplished by^[37] found that preliminary phytochemical screening revealed the presence of steroids, flavonoids, polyphenols, tannins and triterpenoids in *Cassia auriculata* and sterols, triterpenoids, flavonoids, naphthalene glycosides in *Cassia angustifolia*. In the alpha amylase inhibition assay, Acarbose (at concentration 100 µg/ml) showed 50.35% inhibitory effect on alpha amylase activity with IC₅₀ values 102.76±0.65 µg/ml. Both plant extracts showed appreciable on alpha amylase inhibition effect when compared with acarbose. In the alpha glucosidase inhibition assay, the ethanolic extract of *C. auriculata* and *C. angustifolia* (at concentration 100 µg/ml) exhibited 36.2% and 30.45% on alpha glucosidase inhibitory effect with IC₅₀ values 134.9±0.54 µg/ml.

A study done by^[38] found that there was a dose-dependent increase in percentage inhibitory activity against alphaamylase enzyme. A percentage inhibition of 31.10% and 49.14 % were shown in 200µg/ml and 1200µg/ml plant extract, respectively. Glucose diffusion assay was used for the in vitro antidiabetic evaluation. Maximum decrease in glucose diffusion was observed in *Cassia Fistula L.* ethanolic extract as compared to control values. Glucose diffusion is useful in vitro index to predict the effect of plant fibers on the delay in glucose absorption in GI tract. In addition to glucose adsorption, the delay in glucose diffusion might be attributed to the physical obstacle presented by fiber particles towards glucose molecules and entrapment of glucose within the network formed by fibers^[38].

Conclusion

It can be concluded that cassia fistula shows high alpha amylase inhibitory activity and can be used as green medicine for diabetes mellitus.

References

- Hussain SA, Marouf BH. Flavonoids as alternatives in treatment of type 2 diabetes mellitus. *Acad J Med Plants*. 2013; 1:31–6.
- Shastri K. Varanasi: Chanukah Bharati; 1980. Comments on Charaka Samhita.
- Pepato MT, Mori DM, Baviera AM, Harami JB, Vendramini RC, Brunetti IL. Fruit of the Jambolan tree (*Eugenia jambolana* Lam.) and experimental diabetes. *J Ethnopharmacol*. 2005; 96:43–8.
- Venkatesh S, Reddy GD, Reddy BM, Ramesh M, Apparao AV. Antihyperglycemic activity of *Carulluma attenuate*. *Fitoterapia*. 2003; 74:274–7.
- Rajagopalan PM. Madurai: Siddha Maruthuva Gurukulam; 2000. Siddha medicine.
- Duraipandiyar V, Ignacimuthu S. Antibacterial and antifungal activity of *Cassia Fistula L.*: An ethnomedicinal plant. *J Ethnopharmacol*. 2007; 112:590–4.
- Manonmani G, Bhavapriya V, Kalpana S, Govindasamy S, Apparanantham T. Antioxidant activity of *Cassia fistula*(Linn.) flowers in alloxan-induced diabetic rats. *J Ethnopharmacol*. 2005; 97:39–42.
- Theeshan B, Vidushi SN, Okezie IA. Phytochemical constituents of *Cassia fistula*. *Afr J Biotechnol*. 2005; 4:1530–40.
- Nagarajan NS, Muruges N, Thirupathy KP, Radha N, Murali A. Antidiabetic and antihyperlipidemic effects of *Cleome feline*. *Fitoterapia*. 2005; 76:310–5.
- Rao NK, Nammi S. Antidiabetic and renoprotective effects of chloroform extract of *Terminalia chebula* seeds in streptozotocin induced diabetic rats. *BMC Complement Altern Med*. 2006; 6:17.
- Phuong ML, Ali BA, Aziz E, Abdellatif S, Yahia C, Pierre SH. The petroleum ether extract of *Nigella sativa* exerts lipid lowering and insulin-sensitizing action in the rats. *J Ethnopharmacol*. 2004; 94:251–9.
- Gandjalikhan Nassab SA, Bahrami A, Keshtkar MM, Saffaripour MH. Comparison between two methods for enhancing Focal neuropathies following percutaneous nephrolithotomy (PCNL)--preliminary study. *Ger Med Sci*. 2013 Jun 13;11:Doc07. doi: 10.3205/000175. Print 2013.
- Gandjalikhan Nassab SA, Bahrami A, Keshtkar MM, Saffaripour MH. Comparison between two methods for enhancing heat transfer in separated convection flows-second law analysis. 2014. *Iranian Journal of Science and Technology - Transactions of Mechanical Engineering*. Vol. 38, No. M1, pp 9-23.
- Nasseh H, Pourreza E, Saberi A, Kazemnejad E, Kalantari BB, Falahatkar S. Focal neuropathies following

- percutaneous nephrolithotomy (PCNL)--preliminary study. *Ger Med Sci*. 2013 Jun 13;11:Doc07. doi: 10.3205/000175. Print 2013.
15. Araban M, Kariman N, Tavafian SS, Motesaddi S, Alavimajd H, Shokravi FA. Air pollution and low birth weight: a historical cohort study from Tehran. *East Mediterr Health J*. 2012 Jun;18(6):556-60.
 16. Kariman N, Afrakhte M, Hedayati M, Fallahian M, Alavi Majd H. Diagnosis of premature rupture of membranes by assessment of urea and creatinine in vaginal washing fluid. *Iran J Reprod Med*. 2013 Feb;11(2):93-100.
 17. Keshtkar MM, Talebizadeh P. Multi-objective optimization of cooling water package based on 3E analysis: A case study. *Volume 134*, 1 September 2017, Pages 840-849.
 18. Keshtkar MM. Performance analysis of a counter flow wet cooling tower and selection of optimum operative condition by MCDM-TOPSIS method. *Applied Thermal Engineering*. *Volume 114*, 5 March 2017, Pages 776-784.
 19. Shahrzad E, Zahiri S, Ghasemi F, Kargar Jahromi H. A study of effects of l-carnitine on morphology and apoptosis in cryopreserved sperm. 2013. *Advances in Environmental Biology*, 7(9): 2126-2134.
 20. Heidari T, Kariman N, Afrakhteh M, Alavi Majd H. The study of relationship between menorrhagia and gestational diabetes. 2008, *Koomesh*. 9(2), pp. 147-153+22
 21. Alinejad V, Shadmehr A, Asfeh SK, Gholizade R, Tabbakhi E. Examine the relationship between organizational citizenship behavior and organizational health in specialty and subspecialty Seyyed-al-Shohada hospital staff, Urmia . *The Social Sciences*, 2016, Volume: 11, Issue: 6, Page No.: 910-917.
 22. Beheshtifar, M., Hoseinifar, H., Moghadam, M.N. Effect procrastination on work-related stress. . 2011. *European Journal of Economics, Finance and Administrative Sciences*. Issue 38 (2011), p59-64.
 23. Keshtkar MM, Gandjalikhan Nassab SA, Nasr MRJ. Heat transfer characteristics of a cylindrical porous radiant air heater under the influence of a two-dimensional axisymmetric radiative field. *Proceedings of the Institution of Mechanical Engineers, Part A: Journal of Power and Energy*. Vol 223, Issue 8, 2009.p913-923.
 24. Saei Ghare Naz M, Ozgoli G, Aghdashi MA, Salmani F. Prevalence and Risk Factors of Osteoporosis in Women Referring to the Bone Densitometry Academic Center in Urmia, Iran. *Glob J Health Sci*. 2015 Nov 18;8(7):135-45. doi: 10.5539/gjhs.v8n7p135.
 25. Aghdashi MA, Aghdashi MM, Rabiepoor M. Osteopoikilosis: pain as a presenting symptom in three family members. *Clin Med Insights Arthritis Musculoskelet Disord*. 2011 May 2;4:29-32. doi: 10.4137/CMAMD.S7035.
 26. Nahidi F, Kariman N, Simbar M, Mojab F. The Study on the Effects of Pimpinella anisum on Relief and Recurrence of Menopausal Hot Flashes. *Iran J Pharm Res*. 2012 Fall;11(4):1079-85.
 27. Kariman N, Simbar M, Ahmadi F, Vedadhir AA. Socioeconomic and emotional predictors of decision making for timing motherhood among Iranian women in 2013. *Iran Red Crescent Med J*. 2014 Feb;16(2):e13629. doi: 10.5812/ircmj.13629. Epub 2014 Feb 5.
 28. Haidari M, Nazer MR, Ahmadinejad M, Almasi V, Khorramabadi MS, Pournia Y. Honey in the treatment of Fournier's gangrene as an adjuvant: a cross sectional study. *J Pak Med Assoc*. 2014 May;64(5):571-3.
 29. Hemmati L, Rojhani-Shirazi Z, Ebrahimi S. Effects of Plantar Flexor Muscle Static Stretching Alone and Combined With Massage on Postural Balance. *Ann Rehabil Med*. 2016 Oct;40(5):845-850. Epub 2016 Oct 31.
 30. Bernfeld P (1955). *Amylases α and β in: Methods in Enzymology*, edited by Clowick SP and Kaplan NO. Vol 1, Academic Press Inc Publishers. New York, 149 – 152.
 31. Shrikant N. Malpani1* and KP. Manjunath2 *IJAPBC – Vol. 2(1)*, Jan-Mar, 2013.
 32. HasenahAliaP.J. HoughtonaAmalaSoumyanathb, *Journal of Ethnopharmacology*, Volume 107, Issue 3, 11 October 2006, Pages 449-455.
 33. Anitha Roy1 and R.V Geetha2, *Int J Pharm Bio Sci* 2013 Oct; 4(4): (P) 49- 54.
 34. B. Dineshkumar, Analava Mitra, M. Manjunatha, *International Journal of Green Pharmacy | April-June 2010*. |
 35. G. Nivetha*, V. Vishnupriya, R. Gayathri *Int. J. Pharm. Sci. Rev. Res.*, 39(1), July – August 2016; Article No. 43, Pages: 221-225.
 36. B.S. Ashok Kumar, K. Lakshman, R. Nandeesh, P.A. Arun Kumar, B. Manoj, Vinod Kumar, D. Sheshadri Shekar, *Saudi Journal of Biological Sciences* Volume 18, Issue 1, January 2011, Pages 1-5.
 37. Shravan Kumar Nanumala, Tulasi P and Errabelli Sujitha, *In vitro anti-diabetic activity of seed extracts of Cassia auriculata and Cassia angustifolia*, *Euro. J. Exp. Bio.*, 2015, 5(5):12-17.
 38. Soundiramani Balraj, R. Indumathy, Dr. N. Jayshree, M. Sakthi Abirami, *Evaluation of Invitro Anti-Diabetic Activity of Various Root Extract of Cassia Fistula L.*, *Imperial Journal of Interdisciplinary Research (IJIR)* Vol-2, Issue-6, 2016.