

# Efficacy of aerobic training on pulmonary functions and depression in elderly COPD patients

Gaber Sayed Soliman<sup>1,2</sup>, Walid Kamal Abdelbasset<sup>3,4\*</sup>

<sup>1</sup>Department of Physical Therapy and Health Rehabilitation, College of Applied Medical Sciences in Al-Qurayyat, Jouf University, Saudi Arabia. <sup>2</sup>Department of Physical Therapy for Cardiovascular/Respiratory Disorder and Geriatrics, Faculty of Physical Therapy, Cairo University, Giza, Egypt. <sup>3</sup>Department of Physical Therapy and Health Rehabilitation, College of Applied Medical Sciences, Prince Sattam bin Abdulaziz University, Alkharj, Saudi Arabia. <sup>4</sup>Department of Physical Therapy, Kasr Al-Aini Hospital, Cairo University, Giza, Egypt.

**Correspondence:** Walid Kamal Abdelbasset, Department of Physical Therapy and Health Rehabilitation, College of Applied Medical Sciences, Prince Sattam bin Abdulaziz University, Alkharj, Saudi Arabia. E-mail: walidkamal.wr@gmail.com, Mobile: +966561014872.

## ABSTRACT

**Background:** COPD in the elderly population is usually associated with depression that decreases quality of life and increases mortality. The main aim of the study was to assess the therapeutic effects of low to moderate-intensity aerobic training on pulmonary functions and depression levels in elderly moderate COPD patients. **Methods:** This randomized controlled study included thirty-four elderly moderate COPD patients with mild to moderate depression status, their age was  $\geq 65$  years. The patients were divided into the study and control groups randomly. The study group ( $n=17$ ) conducted a low to moderate-intensity aerobic training, three times/week for twelve weeks while the control group ( $n=17$ ) did not conduct any exercise training. Before and after the study intervention, pulmonary functions and depression scores were assessed. **Results:** Baseline characteristics presented a non-significant difference between the study and control groups ( $p>0.05$ ). A significant improvement in pulmonary functions and depression score was observed in the study group after finishing the study program ( $p<0.05$ ) while non-significant changes were shown in the control group. The findings showed significant differences between groups in pulmonary functions and depression scores in favor of the study group ( $p<0.05$ ). **Conclusions:** Low to moderate-intensity aerobic training improves pulmonary function and reduces depression in elderly COPD patients.

**Keywords:** COPD, aerobic training, depression status, pulmonary functions

## Introduction

Chronic obstructive pulmonary disease (COPD) considers a preventable and treatable common pulmonary disease characterizing by progressively persistent airflow obstruction affecting the bronchopulmonary system.<sup>[1-3]</sup> The progressive course of COPD is associated with many complications such as cardiovascular problems, impairment of

deterioration of quality of life reduced exercise capacity, and an increase in mortality.<sup>[4]</sup>

Physical therapy increases exercise capacity and mobility in daily activities, decreases dyspnea, and improves quality of life by applying various therapeutic modalities.<sup>[5-6]</sup> Decreased exercise capacity in COPD during stable and exacerbation periods increases the mortality rate accompanied by weakness of the respiratory muscles that affect the lung functions and decreases muscle mass resulting from the breakdown of muscle proteins and atrophy of skeletal muscles.<sup>[7]</sup> The spreading rate of COPD has increased throughout the past 30 years. COPD is considered the fourth common cause of death worldwide with a rate of 5.1% and it has been predicted that it will be the seventh common cause of disability in 2030. There is an increase in the economic and social burden of COPD.<sup>[8]</sup> Symptoms of COPD including productive cough and progressive exertional dyspnea lead to activity restriction and finally the inability to work efficiently.<sup>[9]</sup>

### Access this article online

Website: [www.japer.in](http://www.japer.in)

E-ISSN: 2249-3379

**How to cite this article:** Soliman GS, Abdelbasset WK. Efficacy of aerobic training on pulmonary functions and depression in elderly COPD patients. *J Adv Pharm Edu Res* 2019;9(4):39-45.  
Source of Support: Nil, Conflict of Interest: None declared.

skeletal muscle function, and depression that result in

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

COPD can be examined by different methods such as spirometer and the COPD assessment test that determine the airflow limitation severity through the forced expiratory volume in one second (FEV1) expressed as a percentage of the normal according to the person's age, gender, height, and weight as FEV1 $\geq$ 80 is mild, FEV1=50–79 is moderate, FEV1=30–49 is severe, and FEV1 $\leq$ 30 is very severe.<sup>[10]</sup>Inpatient, outpatient, younger, and elderly COPD patients usually suffer from depression ranging from depressive symptoms for short-term to dysthymia up to major depression.<sup>[11]</sup>

The COPD patients are related to depression commonly depending on the people surveyed and the assessment tools. There was an overlapping between COPD symptoms and depression symptoms that participate in the prevalence rate variations.<sup>[12]</sup> Several studies approved that outpatients with COPD have depression rates varying from 7% to 80%.<sup>[13]</sup> Also, in stable COPD patients, the range of depression rate is 10%-42%.The comorbid psychological problems in COPD patients may result in a higher functional impairment, disability, morbidity, and lower quality of life.<sup>[14]</sup>

Recently, several studies suggest that congenital factors, environmental factors including smoking and inflammation, the severity of symptoms, and lowering quality of life are considered the strongest indicators of depression among COPD patients<sup>[15]</sup> that affect the adherence to the treatment <sup>[16]</sup>, increase the risk of hospitalization, double the duration of hospitalization, and increase mortality rates. Therefore, there is recently a focus on how to assess and treat persisting depressive symptoms in COPD patients.<sup>[17]</sup>

The treatment of COPD includes combined several modalities such as pulmonary rehabilitation therapy as an independent treatment approach of the disease, cognitive behavioral therapy, anti-depressive interventions, pharmacological treatment, psychological interventions, psychotherapy, group therapy, health education, counseling, behavioral interventions, yoga, Tai Chi, relaxation therapy, and alternative modalities.<sup>[18-20]</sup>

It has been approved that aerobic training has a positive influence on COPD patients but few studies have been conducted to determine the effect of low to moderate-intensity aerobic training on pulmonary functions and depression status and in elderly patients with moderate COPD. Based on that, the hypothesis of the current study was that low to moderate-intensity aerobic training could be beneficial in improving

pulmonary functions and depression status in elderly patients with moderate COPD. The main goal of the current study was to assess the therapeutic effects of low to moderate-intensity aerobic training on pulmonary functions and depression levels in elderly moderate COPD patients.

## Subjects and Methods

### Subjects

Thirty-nine elderly patients with moderate COPD (FEV1= 50–79% predicted) and mild to moderate depression and age of more than 65 years participated in the study from January to May 2019. Each patient was medically stable. All participants were medically diagnosed with depression in accordance with a valid patient-health questionnaire-9 (PHQ-9) regarding the clinical characteristics of depression status. Each participant referred to physiotherapy and outpatient rehabilitation according to his/her chest physicians and psychiatrists of the tertiary hospital. The participant was excluded from the study if he/she suffered from severe depression, orthopedic complications, neurological dysfunctions, and severe life-limiting illness.

### Ethical approval

The study design was a randomized controlled study. The study protocol was ethically approved by the local ethics committee of the Physical Therapy Department, Cairo University Hospitals and complied with the Declaration of Helsinki. Each patient signed written informed consent before participating in the study program.

### Sample size and randomization

Thirty-nine participants were enrolled in the study in accordance with resources' availability rather than power and sample size calculation. Of the thirty-nine participants, thirty-four completed this study, two did not encounter the inclusion criteria of the study, and three declined to conduct the study program. The assignment was performed by the blinded physiotherapist before starting the study program. To avoid type II error, the 34 participants were randomly divided into 2 equal groups (n=17) using the table of random digits. The first group (10 males and 7 females) was recruited in the low to moderate-intensity aerobic training (study group) three sessions weekly for 12 weeks while the second group (9 males and 8 females) was not recruited in any exercise intervention (control group). The flow chart of the study is described in Figure 1.




Figure 1: Flow chart of the study

## Procedures of the study

### Evaluation

All participants of the study were evaluated for the depression scores consuming a valid PHQ-9 and the pulmonary functions (FVC and FEV1) using a digital spirometer (CONTEC: SP10, China) before and after completing the study program (after 12-week intervention) by the same blinded physiotherapist. The validated PHQ-9 was consumed to evaluate the depression scores.<sup>[21]</sup> The PHQ-9 scale identifies the depression level (mild=5-9, moderate=10-19, and severe $\geq$  20).

### 12-week intervention

The participants of the 2 groups were recommended to be active and behave a home-based exercise and have their medical descriptions regularly. The study group received an exercise program of low to moderate-intensity aerobic training, three sessions weekly for twelve weeks. Before starting the exercise program, the maximal heart rate ( $HR_{max}$ ) was recorded. Each exercise session was commenced with warming up (5-10 min) and finished with cooling down (5-10 min). The twelve weeks of the study were started with the initial 6 weeks of low-intensity aerobic training and ended with 6 weeks of moderate-

intensity aerobic training. The low-intensity aerobic was performed conducting the treadmill exercise with 40-50%  $HR_{max}$  for 20 to 30 min, 3 times/week while the moderate-intensity aerobics was performed conducting the treadmill exercise 50-70%  $HR_{max}$  for 30 to 40 min, 3 times/week.

### Statistical Analysis

Data were demonstrated and analyzed in the form of means $\pm$ SD. The descriptive analysis measured the differences in the mean values of FVC, FEV1, and PHQ-9 between the two study groups using unpaired t-test while the changes intragroup were assessed using the paired t-test. Statistical analysis was performed using SPSS software (SPSS v.20, Chicago, IL, USA). Significance was set at  $p < 0.05$ .

## Results

As demonstrated in Table 1, baseline characteristics presented non-significant differences between the two study groups in gender, age, body mass index (BMI), FVC, FEV1, and PHQ-9 ( $p > 0.05$ ).

Table 1. Baseline characteristics of the participants

Measures	SG (n=17)	CG (n=17)	p-value
Gender (M/F)	10/7	9/8	0.920
Age (Years)	69.5 $\pm$ 4.23	69.8 $\pm$ 4.51	0.843
BMI (Kg/m <sup>2</sup> )	26.33 $\pm$ 3.72	27.21 $\pm$ 3.84	0.502
RR (breaths/m)	21.2 $\pm$ 3.2	19.8 $\pm$ 3.6	0.239
HR (beat/minute)	102.4 $\pm$ 23.5	105.3 $\pm$ 24.2	0.725
FVC (% predicted)	72.7 $\pm$ 6.4	71.3 $\pm$ 6.8	0.541
FEV1 (% predicted)	61.8 $\pm$ 4.72	62.4 $\pm$ 4.23	0.699
PHQ-9	15.73 $\pm$ 3.25	14.88 $\pm$ 2.86	0.424

SG: Study Group; CG: Control Group; BMI: Body Mass Index; RR: Respiratory Rate; HR: Heart Rate; FVC: Forced Vital Capacity; FEV1: Forced Expiratory Volume in the first second; PHQ-9: Patient Health Questionnaire-9.

After completing the study program, there was the significant improvement of FVC, FEV1, and PHQ-9 in the exercise group ( $p < 0.05$ ) while the control group exhibited non-significant

changes in the outcome measures ( $p > 0.05$ ) as shown in Table 2.

**Table 2. Changes of mean values within each group before and after the intervention**

Measures	SG (n=17)			CG (n=17)		
	Before	After	p-value	Before	After	p-value
FVC (% predicted)	72.7±6.4	83.2±5.31	<0.0001	71.3±6.8	73.5±6.6	0.345
FEV1 (% predicted)	61.8±4.72	76.2±3.45	<0.0001	62.4±4.23	63.9±4.62	0.331
PHQ-9	15.73±3.25	7.43±2.11	<0.0001	14.88 ±2.86	13.7±2.91	0.242

SG: Study Group; CG: Control Group; FVC: Forced Vital Capacity; FEV1: Forced Expiratory Volume in the first second; PHQ-9: Patient Health Questionnaire-9.

Comparing the study group (Low to moderate-intensity exercise) to the control group, the findings demonstrated significant differences between the two study groups in all mean

values of FVC, FEV1, and PHQ-9) in favor of the study group at the end of the study program ( $p < 0.05$ ) as shown in Table 3.

**Table 3. Differences between groups after intervention**

Measures	SG (n=17)	CG (n=17)	p-value
FVC (% predicted)	83.2±5.31	73.5±6.6	<0.0001
FEV1 (% predicted)	76.2±3.45	63.9±4.62	<0.0001
PHQ-9	7.43±2.11	13.7±2.91	<0.0001

SG: Study Group; CG: Control Group; FVC: Forced Vital Capacity; FEV1: Forced Expiratory Volume in the first second; PHQ-9: Patient Health Questionnaire-9.

## Discussion

The present study was conducted to evaluate the effect of low to moderate-intensity aerobic training on depression status and pulmonary functions in elderly patients with moderate COPD. They were randomly divided into 2 equal groups; each group consisted of seventeen patients. The study group conducted the low to moderate-intensity aerobic training and the control group did not conduct any exercise program. Regular aerobic exercise training is essential for general health well-being and psychological illness and many studies approved safe and beneficial findings<sup>[22-24]</sup>, although many investigations did not encourage these results.<sup>[25,26]</sup> Previous studies compared the effects of low to moderate-intensity with moderate-intensity continuous exercises on the depressive state in patients with heart failure and found that both exercise programs had significant and beneficial effects on reducing depression status in patients with heart failure with no differences between them.<sup>[27]</sup> Several studies approved that aerobic exercises have effective results in decreasing depression illness and major symptoms of depressive disorder.<sup>[23,24,27]</sup> Other investigations confirmed the secretion of dopamine, norepinephrine, endorphins, and serotonin with aerobic training contradicting the effects of depressive disorders,<sup>[28]</sup> which confirmed that aerobic training has a positive effect on depressed subjects in controlling stress, independence, self-confidence, and normal mental state. The current study confirmed that low to moderate-intensity aerobic training is safe, reliable, and endorsed the role of physiotherapy in controlling depression in elderly COPD patients. The results

of this study were supported by Dunn et al., who approved that ten days of walking exercises (30 minutes daily) achieved the optimum decreasing in the major depression disorder depressed individuals using a depressive rate scale.<sup>[29]</sup>

In agreement with the present study results, another study approved that endurance training for the short-term results in an extreme reduction of depression disorders than anti-depressive drugs in cases with depression.<sup>[30]</sup> Similarly, the study conducted by Knubben et al. approved that in people with coronary artery disease, supervised aerobic training leads to positive effects on the functional capacity and reduction of depression manifestations more than home-based aerobic training.<sup>[31]</sup>

The present study was protocolled to assess the pulmonary functions, exercise tolerance, and psychological parameters and its relevance with COPD. Patients with COPD demonstrated decreased pulmonary function and low exercise tolerance. This study approved significant correlations between aerobic exercises and pulmonary functions. Severe COPD ( $FEV1 < 50\%$ ) can cause a considerable decrease in walk capacity.<sup>[32,33]</sup> Previous research approved that pulmonary functions are usually disturbed in COPD patients<sup>[34]</sup> explaining why the COPD patients are fatigued and inactive, which considered exercise training as a useful modality for improving pulmonary functions and physical performance.<sup>[35]</sup> In agreement with the results of this study, previous studies confirmed a significant relationship between aerobic training capacity and pulmonary functions. Consequently, decreased exercise capacity considered an important factor in severity in COPD.<sup>[36,37]</sup>

On the other hand, the results of this study were contradicted by Yohannes, who approved the low benefits of exercise in COPD patients. The pathological deteriorations that occur in the lung could be related to the lack of physical activity and negative impacts of self-confidence leading to not regularly perform aerobic training.<sup>[38]</sup>

Similarly, various investigations have confirmed improvement in the psychological aspects of people with COPD when performing aerobic training within the limit of moderate sensation.<sup>[39,40]</sup> Comparing the pre- and post-exercise values showed positive changes in COPD patients treated with a low-intensity exercise program.<sup>[41]</sup>

After a multidisciplinary pulmonary rehabilitation program, performed by Güell et al., no statistically significant increases were observed in the FVC and FEV1. This result was contributed to the increase in respiratory muscle strength but the lack of change in respiratory functions. Additionally, there were non-significant changes in the arterial blood gases (ABG) values of the patients and a statistically significant increase in the respiratory functions of the FEV1 value (from 60.4 to 70.6), FEV1/FVC ratio (from 56.2 to 62.5), and FEF 25–75 value (from 29.4 to 35.5).<sup>[42]</sup>

In addition to improvement in the musculoskeletal system at the end of the exercise, studies have shown improvement in pulmonary functions. In the study conducted by Donaire-Gonzalez et al., it was observed that low-intensity exercise decreases hospitalization and mortality rate and improves pulmonary functions and functional capacity.<sup>[7]</sup> Moreover, a recent study concluded that aerobic exercise improves pulmonary functions, aerobic capacity, and quality of life in asthmatics.<sup>[43]</sup>

Some limitations were appointed in the present study including missing depression drugs for participants of the study. Besides, exercise compliance with medications, the expenditure of energy, and physical fitness were not assessed to achieve the optimum evaluation of exercise intensity. Further proposals are necessitated to investigate the effect of various durations and intensities of exercise on pulmonary functions and depression state in elderly COPD patients.

## Conclusions

It was concluded that low to moderate-intensity aerobics has beneficial effects on improving pulmonary function and reducing the level of depression in elderly COPD patients. Approximately, this type of exercise program has been prescribed among depressed subjects particularly elderly patients with moderate COPD.

## Funding

No funds were received for this study.

## Competing interests

No competing interests to disclose.

## Acknowledgment

This publication was supported by the Deanship of Scientific Research at Prince Sattam bin Abdulaziz University. The authors acknowledge all COPD patients who were included in the study.

## References

1. American Lung Association Epidemiology and Statistics Unit. Trends in COPD (Chronic Bronchitis and Emphysema): Morbidity and Mortality; 2013. Available from: <http://www.lung.org/assets/documents/research/copd-trend-report.pdf>. [Last accessed on 2016 Aug 12].
2. Kim V, Crapo J, Zhao H, Jones PW, Silverman EK, Comellas A, Make BJ, Criner GJ. Comparison between an alternative and the classic definition of chronic bronchitis in COPD. *Gene. Annals of the American Thoracic Society*. 2015 Mar;12(3):332-9.
3. Global Initiative for Chronic Obstructive Lung Disease [webpage on the Internet]. Global strategy for the diagnosis, management, and prevention of COPD: GOLD 2017. Available from: <http://goldcopd.org/gold-2017-global-strategy-diagnosis-management-prevention-copd/> Global. Accessed December 20, 2017.
4. Abdelbasset WK, Nambi G, Mahmoud MZ. Impact of resisted exercise on chronic obstructive pulmonary disease (COPD) in elderly patients in Al-Kharj, Saudi Arabia. *Int J Med Res Health Sci* 2016;5(5):187-195.
5. Grigorova-Petrova K, Lubenova D, Dimitrova A, Baldaranov D, Lozeva J. Feasibility of early physical therapy program in-hospital patients with acute ischemic stroke. *Macedonian Journal of Medical Sciences*. 2014 Sep 1;7(3):452-5.
6. Abdelbasset WK, Elnegamy TE. Effect of chest physical therapy on pediatric hospitalized with pneumonia. *International Journal of Health and Rehabilitation Science*. 2015;4(4):219-26.
7. Donaire-Gonzalez D, Gimeno-Santos E, Balcells E, de Batlle J, Ramon MA, Rodriguez E, Farrero E, Benet M, Guerra S, Sauleda J, Ferrer A. Benefits of physical activity on COPD hospitalisation depend on intensity. *European Respiratory Journal*. 2015 Nov 1;46(5):1281-9.
8. Lopez AD, Murray CC. The global burden of disease, 1990–2020. *Nature medicine*. 1998 Nov;4(11):1241. Available from: [http://www.nature.com/articles/nm1198\\_1241](http://www.nature.com/articles/nm1198_1241). [Last accessed on 2019 Mar 28].
9. Kessler R, Partridge MR, Miravittles M, Cazzola M, Vogelmeier C, Leynaud D, Ostinelli J. Symptom variability in patients with severe COPD: a pan-European cross-sectional study. *European Respiratory Journal*. 2011 Feb 1;37(2):264-72.

10. Vestbo J, Hurd SS, Agustí AG, Jones PW, Vogelmeier C, Anzueto A, Barnes PJ, Fabbri LM, Martinez FJ, Nishimura M, Stockley RA. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *American journal of respiratory and critical care medicine*. 2013 Feb 15;187(4):347-65.
11. Hill K, Geist R, Goldstein RS, Lacasse Y. Anxiety and depression in end-stage COPD. *European respiratory journal*. 2008 Mar 1;31(3):667-77.
12. Tselebis A, Bratis D, Pachi A, et al. Anxiety and depression in patients with chronic obstructive pulmonary disease (COPD). In: Zirimis L, Papazoglakis A, editors. *Chronic Obstructive Pulmonary Disease: New Research*. New York, NY: Nova Science Publishers; 2013:15–40.
13. Katz PP, Julian LJ, Omachi TA, Gregorich SE, Eisner MD, Yelin EH, Blanc PD. The impact of disability on depression among individuals with COPD. *Chest*. 2010 Apr 1;137(4):838-45.
14. Maurer J, Rebbapragada V, Borson S, Goldstein R, Kunik ME, Yohannes AM, Hanania NA. Anxiety and depression in COPD: current understanding, unanswered questions, and research needs. *Chest*. 2008 Oct 1;134(4):43S-56S.
15. Hanania NA, Müllerova H, Locantore NW, Vestbo J, Watkins ML, Wouters EF, Rennard SI, Sharafkhanh A. Determinants of depression in the ECLIPSE chronic obstructive pulmonary disease cohort. *American journal of respiratory and critical care medicine*. 2011 Mar 1;183(5):604-11.
16. Kosmas E, Tselebis A, Bratis D, Dumitru S, Theodorakopoulou E, Harikiopoulou M, Velentzas K, Bakakos P, Alchanatis N, Koulouris N. The Relationship between the adherence to treatment and the psychological profile of patients with chronic obstructive pulmonary disease. *Chest*. 2014 Mar 1;145(3):377A.
17. Pooler A, Beech R. Examining the relationship between anxiety and depression and exacerbations of COPD which result in hospital admission: a systematic review. *International journal of chronic obstructive pulmonary disease*. 2014;9:315.
18. Spruit MA, Singh SJ, Garvey C, ZuWallack R, Nici L, Rochester C, Hill K, Holland AE, Lareau SC, Man WD, Pitta F. An official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. *American journal of respiratory and critical care medicine*. 2013 Oct 15;188(8):e13-64.
19. McCarthy B, Casey D, Devane D, Murphy K, Murphy E, Lacasse Y. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane database of systematic reviews*. 2015(2).
20. Volpato E, Banfi P, Rogers SM, Pagnini F. Relaxation techniques for people with chronic obstructive pulmonary disease: a systematic review and a meta-analysis. *Evidence-Based Complementary and Alternative Medicine*. 2015;2015.
21. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *Journal of general internal medicine*. 2001 Sep;16(9):606-13.
22. Abdelbasset WK, Alqahtani BA. A randomized controlled trial on the impact of moderate-intensity continuous aerobic exercise on the depression status of middle-aged patients with congestive heart failure. *Medicine*. 2019 Apr 1;98(17):e15344.
23. Abdelbasset WK, Elsayed SH, Abo Elyazed TI. Comparison of High-Intensity Interval to Moderate Intensity Continuous Aerobic Exercise on Ventilatory Markers in Coronary Heart Disease Patients: A Randomized Controlled Study. *Int J Physiother Res*. 2017;5(3):2013-18.
24. Abdelbasset WK, Alqahtani BA, Elshehawy AA, Tantawy SA, Elnegamy TE, Kamel DM. Examining the impacts of 12 weeks of low to moderate-intensity aerobic exercise on depression status in patients with systolic congestive heart failure-A randomized controlled study. *Clinics*. 2019;74:e1017.
25. Kerse N, Hayman KJ, Moyes SA, Peri K, Robinson E, Dowell A, Kolt GS, Elley CR, Hatcher S, Kiata L, Wiles J. Home-based activity program for older people with depressive symptoms: DeLLITE—a randomized controlled trial. *The Annals of Family Medicine*. 2010 May 1;8(3):214-23.
26. Sims J, Hill K, Davidson S, Gunn J, Huang N. Exploring the feasibility of a community-based strength training program for older people with depressive symptoms and its impact on depressive symptoms. *BMC geriatrics*. 2006 Dec;6(1):18.
27. Abdelbasset WK, Alqahtani BA, Alrawaili SM, Ahmed AS, Elnegamy TE, Ibrahim AA, Soliman GS. Similar effects of low to moderate-intensity exercise program vs moderate-intensity continuous exercise program on depressive disorder in heart failure patients: A 12-week randomized controlled trial. *Medicine*. 2019 Aug;98(32):e16820.
28. Rustad JK, Stern TA, Hebert KA, Musselman DL. Diagnosis and treatment of depression in patients with congestive heart failure: a review of the literature. *The primary care companion for CNS disorders*. 2013;15(4).
29. Dunn AL, Trivedi MH, Kampert JB, Clark CG, Chambliss HO. Exercise treatment for depression: efficacy and dose response. *American journal of preventive medicine*. 2005 Jan 1;28(1):1-8.
30. Harris AH, Cronkite R, Moos R. Physical activity, exercise coping, and depression in a 10-year cohort study of depressed patients. *Journal of Affective Disorders*. 2006 Jul 1;93(1-3):79-85.
31. Knubben K, Reischies FM, Adli M, Schlattmann P, Bauer M, Dimeo F. A randomised, controlled study on the effects of a short-term endurance training programme in

- patients with major depression. *British journal of sports medicine*. 2007 Jan 1;41(1):29-33.
32. American Thoracic Society (ATS). Standardization of spirometry. *Am J Resp Crit Care Med* 1995;152:1107–36.
  33. Casanova C, Cote C, Marin JM, Pinto-Plata V, de Torres JP, Aguirre-Jaime A, Vassaux C, Celli BR. Distance and oxygen desaturation during the 6-min walk test as predictors of long-term mortality in patients with COPD. *Chest*. 2008 Oct 1;134(4):746-52.
  34. Velloso M, Jardim JR. Study of energy expenditure during activities of daily living using and not using body position recommended by energy conservation techniques in patients with COPD. *Chest*. 2006 Jul 1;130(1):126-32.
  35. Buist AS, Vollmer WM, McBurnie MA. Worldwide burden of COPD in high-and low-income countries. Part I. The Burden of Obstructive Lung Disease (BOLD) Initiative [State of the Art Series. Chronic obstructive pulmonary disease in high-and low-income countries. Edited by G. Marks and M. Chan-Yeung. Number 6 in the series]. *The international journal of tuberculosis and lung disease*. 2008 Jul 1;12(7):703-8.
  36. Kundu A, Maji A, Sarkar S, Saha K, Jash D, Maikap M. Correlation of six minute walk test with spirometric indices in chronic obstructive pulmonary disease patients: A tertiary care hospital experience. *The Journal of Association of Chest Physicians*. 2015 Jan 1;3(1):9.
  37. Cancellero-Gaiad KM, Ike D, Soave L, Gomes EL, Dias FD, Costa D. Correlation between functional capacity and health-related quality of life in COPD: a case series. *Fisioterapia em Movimento*. 2014 Dec;27(4):505-14..
  38. Yohannes AM. Palliative care provision for patients with chronic obstructive pulmonary disease. Health and quality of life outcomes. 2007 Dec;5(1):17.
  39. Jackson BE, Coultas DB, Ashmore J, Russo R, Peoples J, Uhm M, Singh KP, Bae S. Domain-specific self-efficacy is associated with measures of functional capacity and quality of life among patients with moderate to severe chronic obstructive pulmonary disease. *Annals of the American Thoracic Society*. 2014 Mar;11(3):310-5.
  40. Manger TA, Motta RW. The impact of an exercise program on posttraumatic stress disorder, anxiety, and depression. *International journal of emergency mental health*. 2005.
  41. Cilekar S, Tülek B, Kanat F, Süerdem M, Levendoglu F, Taşpınar IT. Effect of low-intensity pulmonary rehabilitation program on quality of life and pulmonary functions in patients with stable chronic obstructive pulmonary disease. *Eurasian Journal of Pulmonology*. 2019 Jan 1;21(1):14.
  42. Güell R, Casan P, Belda J, Sangenis M, Morante F, Guyatt GH, Sanchis J. Long-term effects of outpatient rehabilitation of COPD: a randomized trial. *Chest*. 2000 Apr 1;117(4):976-83.
  43. Abdelbasset WK, Alsubaie SF, Tantawy SA, Elyazed TI, Kamel DM. evaluating pulmonary function, aerobic capacity, and pediatric quality of life following a 10-week aerobic exercise training in school-aged asthmatics: a randomized controlled trial. *Patient prefer adher*. 2018;12:1015-1023.