

The relationship between lipid profile and CRP in severe sepsis with patient prognosis

Ebrahim Hazrati¹, Mahtab Noorifard², Mohamad Afsahi³, Alireza Rohi⁴, Mohamadreza Rafiei^{5*}

¹Assistant professor, Department of Anesthesiology & Critical care, AJA University of Medical Sciences, Tehran, Iran, ²Associate Professor, Infectious Diseases Research Center, Aja university of Medical Science, ³ Assistant professor, Department of Anesthesiology, AJA University of Medical Sciences, Tehran, Iran, ⁴Researcher, Aja university of Medical Science, ⁵ Associate professor, Department of Anesthesiology, AJA University of Medical Sciences ,Tehran, Iran.

Correspondence: Mohamadreza Rafiei, Associate professor, Department of Anesthesiology, AJA University of Medical Sciences, Tehran, Iran. E_mail: m.rafiee@ajajums.ac.ir

ABSTRACT

Since severe sepsis result to mass casualties during its course, understanding the factors involved in sepsis prognosis can decrease its morbidities and mortalities. So we conduct this study to evaluate lipid profile and CRP of plasma and patient prognosis. Methods: This observational cohort study was carried out with 60 patient suffering from severe sepsis and hospitalized to Imam Reza ICU ward. Serum lipid profil and CRP were measured on first, third and seventh days and compared with patient, s SAPS. Data were analyzed on the basis of student's t – test and χ^2 test ANOVA test and within subject ANOVA with repeated measures tests and using SPSS- 18 software. Results: The results are expressed as mean \pm SE and $P < 0.05$ is considered statistically significant. According to our findings CRP level was high on the first, third and seven days in dead patients ($p=0.042$ $p=0.018$, $p=0.024$ respectively). Also triglycerids, LDL and HDL were low on the seventh day ($p=0.019$, $p=0.026$, $p=0.006$ respectively) and had not different on the first and third days($p>0.05$). Conclusion: Severe sepsis prognosis have relationship with serum lipidprofile and CRP. As with high CRP level and low triglyceride and cholestroles level the patient will have poor prognosis.

Keywords: Sever sepsis, prognosis, CRP, Lipidprofile

Introduction

The patients will have hospitalized in ICU ward, are special patients and have critical condition that susceptible to killer disease such as sepsis [1-5].

Hospital disease specially sepsis can cause ICU prolongation and increase treatment cost thus efforts to reduce the length of ICU stay cause less the hospital cost [2,6-9].

Sepsis prevention and improvement of patient condition have important role to reduce complication, for this reason understanding the factors that influence the course and prognosis of the sepsis is important [3,10-13].

The definition of sepsis was based on the presence of at least 2 of the following criteria [4,14-18]: fever or hypothermia (temperature $\geq 38^\circ\text{C}$ or $\leq 36^\circ\text{C}$), tachycardia (heart rate ≥ 90 beats per minute), tachypnea (respiratory rate ≥ 20 breaths per minute), and leukocytosis or leukopenia (leukocyte count $\geq 12000/\mu\text{L}$ or $\leq 4000/\mu\text{L}$ or $\geq 10\%$ immature forms), concurrently with the presence of confirmed infection.

Additional requirements for severe sepsis were at least 1 of the following [5,19-22]: 1-hypotension (systolic blood pressure ≤ 90 mm Hg, sustained drop in systolic blood pressure ≤ 40 mm Hg, or mean arterial pressure ≤ 65 mmHg corrected within 1 hour by fluid resuscitation), 2- arterial hypoxemia ($\text{PaO}_2 \leq 75$ mmHg without evidence of primary lung disease), 3- metabolic acidosis ($\text{pH} \leq 7.3$ or base deficit ≥ 5 mEq/L), 4- oliguria (urine output ≤ 30 mL/h for at least 2 hours despite adequate fluid replacement), 5- acute alteration of mental status, 6- oagulation abnormalities of recent onset (prothrombin time or activated partial thromboplastin time ≥ 1.2 times the upper normal limit plus D-dimers ≥ 500 or platelets $\leq 100000/\mu\text{L}$).

Lipoprotaine ability in binding to bacterial toxic substance and neutralize them more consider every day [6,23,24].

Access this article online

Website: www.japer.in

E-ISSN: 2249-3379

How to cite this article: Ebrahim Hazrati, Mahtab Noorifard, Mohamad Afsahi, Alireza Rohi, Mohamadreza Rafiei. The relationship between lipid profile and CRP in severe sepsis with patient prognosis. J Adv Pharm Edu Res 2018;8(1):103-107.

Source of Support: Nil, Conflict of Interest: None declared.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Lipoprotein have been showed to reduce cytokine response and mortality rate in vivo and vitro against sepsis [7, 8,25,26].

HDL infusion into rabbit reduced cytokine production in response to bacterial lipopolysaccharide [9,27-30].

In severly ill patients was observed that they had low serum lipid concentration [10, 11,31,32].

Serum

levels of C-reactive protein (CRP), an acute-phase protein synthesized by the liver after stimulus by various cytokines including tumor necrosis factor and interleukin (IL)-6, markedly increase within hours of infection or inflammation [12,33,34].

Numerous studies have shown increased CRP levels in patients with sepsis [13-15,33], but their relation to multiple-organ dysfunction and failure, to date, has not been well evaluated.

Severity scoring systems are designed to provide an estimate of the probability of hospital mortality

for critically ill patient one of this systems is the Simplified Acute Physiology Score (SAPS) that ability of any such systems to accurately predict mortality rate is determined by its discrimination and calibrations [16,34,35]. In this study we evaluate parameters such as CRP and lipoproteins with SAPS to determine prognostic factors in severe sepsis patients.

Material and Method:

In this cohort study 60 severe sepsis patient were studied. Our inclusion criteria were being 20-75 years old age persons with severe sepsis disease that admitted to Imam Reza hospital ICU. Exclusion criteria were DIC and renal insufficiency, died within four hours of admission to ICU or who stayed in the ICU less than 24 hours and if patients had been admitted more than once to the ICU during the study period, only the first admission was included.

Approval for the project was obtained from the faculty Ethics Committee.

Serum lipid profile was determined in all participants, after an overnight fast of 12 hours, on admission (day1) and on days 3 and 7 also CRP was determined on that days.

SAPS of patients were calculated on that days.

Data were analyzed on the basis of student's t – test and χ^2 test ANOVA test and within subject ANOVA with repeated measures tests and using SPSS- 18 software The results are expressed as mean \pm SE and $P < 0.05$ is considered statistically significant.

Results:

In this cohort observational study 60 severe sepsis patient were enrolled and their lipidprofile and CRP and SAPS compared on first and third and seventh days in order to determine prognostic condition of patients. Comparison of the demographic parameters between patients showed uniform distribution (Table 1).

In this study HDL and LDL and TG values from the first day until the seventh gradually will reduce and CRP value from the first to seventh day will increase (Table 2).

Also in this study mean values of HDL and LDL and Tricyclerides decreased in patient with high level of SAPS score and CRP value increased in patient with high level of SAPS score (Table 3).

Table 1: Demographic characteristics

		Total (n=60)
Age (years)		62 \pm 9.2
SAPS	High	12 (20%)
	Moderate	18 (30%)
	Low	30 (50%)
Sex	man	25 (41.7%)
	Woman	35 (58.3%)
Survivors		36 (60%)
Non-survivors		24 (40%)

Table 2: frequency distribution of prognosis according to measured parameters

prognosis	Triglycerids			Cholesterol						CRP		
	First day	Third day	Seventh day	HDL			LDL			First day	Third day	Seventh day
				First day	Third day	Seventh day	First day	Third day	Seventh day			
Survivors	153.1 \pm 26.5	142.5 \pm 25.4	137.4 \pm 26.1	37.2 \pm 3.9	34.4 \pm 3.8	35.7 \pm 5.1	105.25 \pm 17.2	97.16.3 \pm 16.1	93.2 \pm 16.1	45.2 \pm 40.4	49.2 \pm 49.3	53.5 \pm 51.1
Nonsurvivors	145.7 \pm 27.5	131.6 \pm 26.5	120.7 \pm 26.4	35.6 \pm 5.1	32.7 \pm 5.6	32.3 \pm 3.9	98.6 \pm 16.3	90.4 \pm 15.5	84.3 \pm 12.3	67.6 \pm 41.6	82.6 \pm 55.8	85.2 \pm 53.1
p-value	0.05<	0.05<	0.019	0.05<	0.05<	0.006	0.05<	0.05<	0.026	0.042	0.018	0.024

Table 3. frequency distribution of SAPS score according to measured parameters

SAPS Score		Triglycerids			Cholesterol						CRP		
		First day	Third day	Seventh day	HDL			LDL			First day	Third day	Seventh day
					First day	Third day	Seventh day	First day	Third day	Seventh day			
I		153.3 ±28.5	142.8 ±27.7	135.5 ±29.3	37.2 ±3.9	34.6 ±4	35.4 ±4.4	105.5 ±17.7	96.6 ±15.9	92 ±14.9	43.5 ±39.4	38.4 ±38.1	42.3 ±39.6
II		150.3 ±25.1	135.2 ±23.5	126.5 ±24.3	36.7 ±4.2	34.4 ±4.2	34.3 ±5.4	103.1 ±15.6	95.9 ±15.6	91.3 ±15.4	57.6 ±40.5	75.4 ±51.5	80.7 ±53.6
III		142.1 ±26.1	131.1 ±26.1	125.1 ±26.1	33.4 ±5.5	30.4 ±5.5	31.8 ±4.7	94.5 ±16.1	86.5 ±16.1	81.3 ±13.8	75.8 ±44.9	103.9 ±63.8	104 ±59.1
	p-value	0.05<	0.05<	0.05<	0.039	0.020	0.05<	0.05<	0.05<	0.05<	0.05<	0.0001	0.001

I= low II=moderate III=high

Discussion:

Profound changes in the concentration and composition of plasma lipids and lipoproteins have been described in critical illness [36], in intensive care patients with sepsis and in cirrhotic patients with severe sepsis, previous few studies demonstrated that a marked decline in serum HDL levels has been documented [37, 38]. Lower serum cholesterol mechanisms remain unclear [20]. One possibility is that cholesterol is consumed significantly by binding bacterial substances [39-42]. Therefore, a high production of endotoxin consumes more cholesterol and causes a rapid decline of the latter. Another possibility is that high concentrations of proinflammatory cytokines during sepsis suppress lipoprotein production and facilitate lipoprotein degradation [43, 44]. The deficiency in cholesterol renders the host tissue more susceptible to bacterial substance-mediated injury, resulting in cytokine overproduction and a further depletion of cholesterol [39].

C-reactive protein is a marker of inflammation that has been used to monitor the course of infection and inflammatory diseases [45]. C-reactive protein is predominantly produced and secreted by hepatocytes, although other cells including alveolar macrophages may also synthesize CRP. C-reactive protein is thought to represent a measure of cytokine-induced protein synthesis. The relatively short half-life of approximately 19 hours makes it a useful monitor for follow-up of inflammatory response, infection, and antibiotic treatment. In addition, laboratory tests for CRP are easily available and less costly than cytokine tests [46]. Despite its limitations, outcome predictors such as the extensively evaluated SAPS scores are helpful in identifying those septic patients who are at high risk for death and who are more likely to benefit from intervention. Morbidity and mortality in elderly patients admitted to the ICU are higher than in younger patients. The most important factors independently associated with the highest risk of death are severity of illness, 190 D. MemisS et al. impaired level of consciousness, and infection [47]. Hazrati et al (2015) showed Compared to the other centers Mean days of hospital stay was significantly shorter ($p < 0.01$, (15.1 ± 14.9 days) Vs. (23.8 ± 3 days)) and a significant higher percent of patients have history of carbapenem use. (68.8% vs 49.4%, $p = 0.00$). carbapenem resistant infections in shorter hospital stay

and a significantly higher percents of them have history of carbapenem administration [48].

In the present study, the prognostic value of cholesterol and CRP levels was comparable to that of the SAPS score. Our belief is that simultaneous use of cholesterol together with other variables may be needed to increase its prognostic value; these may include, for instance, severity of illness, age, infection, variables considered in currently available scoring systems. We conclude that hypocholesterol and increased CRP concentrations may be a valuable addition to SAPS scores in predicting risk of death, as are the SAPS scores. Serial measurements of cholesterol and CRP concentrations in critically ill patients may help to identify patients who may require more aggressive diagnostic and therapeutic interventions to avoid complications. Cholesterol and CRP concentrations also may be helpful in clinical trials, to identify high-risk patients who would benefit from new therapeutic interventions.

References

1. Kiss CM, Byham-Gray L, Denmark R, Loetscher R, Brody RA. The impact of implementation of a nutrition support algorithm on nutrition care outcomes in an intensive care unit. *Nutr Clin Pract*. 2012 Dec;27(6):793-801.
2. Farber MS, Moses J, Korn M. Reducing costs and patient morbidity in the enterally fed intensive care unit patient. *JPEN J Parenter Enteral Nutr*. 2005 Jan-Feb;29(1 Suppl): S62-9.
3. Warren DK, Quadir WW, Hollenbeak CS, Elward AM, Cox MJ, Fraser VJ. Attributable cost of catheter-associated bloodstream infections among intensive care patients in a nonteaching hospital. *Crit Care Med*. 2006 Aug;34(8):2084-9.
4. American College of Chest Physicians/Society of Critical Medicine Consensus Committee. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Crit Care Med* 1992; 20:864-74
5. Sibbald WJ, Vincent JL. Round table conference on clinical trials for the treatment of sepsis. *Crit Care Med* 1995; 23:394-9.
6. Khovichunkit W, Memon RA, Feingold KR, Grunfeld C. Infection and inflammation-induced proatherogenic

- changes of lipoproteins. *J Infect Dis* 2000;181: S462– 72 [Suppl].
7. Levine D, Parker T, Donnelly T, Walsh A, Rubin AL. In vivo protection against endotoxin by plasma high density lipoprotein. *Proc Natl Acad Sci U S A* 1993; 90:12040– 4.
 8. NeteaMG, DemackerPN, KullbergBJ, BoermanOC, VerschuereI, StalenhoefAL, et al. Low-density lipoprotein receptor-deficient mice are protected against lethal endotoxemia and severe Gram-negative infections. *J Clin Invest* 1996; 97:1366 – 72.
 9. Hubsch AP, Powell FS, Lerch PG, Doran JE. A reconstituted, apolipoprotein A-I containing lipoprotein reduces tumor necrosis factor release and attenuates shock in endotoxemic rabbits. *Circ Shock* 1993;40(1):14 – 23.
 10. Gordon BR, Parker TS, Levine DM, Saal SD, Wang JC, Sloan BJ, et al. Low lipid concentrations in critical illness: implications for preventing and treating endotoxemia. *Crit Care Med* 1996;24(4):584 – 9.
 11. Alvarez C, Ramos A. Lipids, lipoproteins and apoproteins in serum during infection. *Clin Chem* 1986; 32:142 – 5.
 12. Thijs LG, Hack CE. Time course of cytokine levels in sepsis. *Intensive Care Med* 1995;21(Suppl 2): S258-63.
 13. Po'voa P, Almeida E, Moreira P, et al. C-reactive protein as an indicator of sepsis. *Intensive Care Med* 1998; 24:1052- 6.
 14. Yentis SM, Soni N, Sheldon J. C-reactive protein as an indicator of resolution of sepsis in the intensive care unit. *Intensive Care Med* 1995; 21:602- 5.
 15. Smith RP, Lipworth BJ, Cree IA, et al. C-reactive protein: a clinical marker in community-acquired pneumonia. *Chest* 1995; 108:1288 - 91.
 16. Le Gall JR. The use of severity scores in the intensive care unit. *Intensive Care Med* 2005; 31: 1618-23.
 17. Saei Ghare Naz M, Kariman N, Ebadi A, Ozgoli G, Ghasemi V, Rashidi Fakari F. Educational interventions for cervical cancer screening behavior of women: A systematic review. *Asian Pac J Cancer Prev*. 2018 Apr 25;19(4):875-884.
 18. Shahrahmani H, Kariman N, Jannesari S, Rafieian-Kopaei M, Mirzaei M, Ghalandari S, Shahrahmani N, Mardani G. The effect of green tea ointment on episiotomy pain and wound healing in primiparous women: A randomized, double-blind, placebo-controlled clinical trial. *Phytother Res*. 2018 Mar;32(3):522-530. doi: 10.1002/ptr.5999.
 19. Alimoradi Z, Kariman N, Ahmadi F, Simbar M. Preparation for sexual and reproductive self-care in Iranian adolescent girls: a qualitative study. *Int J Adolesc Med Health*. 2017 May 17;30(1). pii: /j/ijamh.2018.30.issue-1/ijamh-2016-0094/ijamh-2016-0094.xml. doi: 10.1515/ijamh-2016-0094.
 20. Atarodi-Kashani Z, Kariman N, Ebadi A, Alavi Majd H, Beladi-Moghadam N. Sexual function and related factors in Iranian woman with epilepsy. *Seizure*. 2017 Nov;52:147-153. doi: 10.1016/j.seizure.2017.10.001. Epub 2017 Oct 5.
 21. Ghalandari S, Kariman N, Sheikhan Z, Mojab F, Mirzaei M, Shahrahmani H. Effect of Hydroalcoholic Extract of *Capsella bursa pastoris* on Early Postpartum Hemorrhage: A Clinical Trial Study. Effect of Hydroalcoholic Extract of *Capsella bursa pastoris* on Early Postpartum Hemorrhage: A Clinical Trial Study.
 22. Alimoradi Z, Kariman N, Simbar M, Ahmadi F. Empowerment of Adolescent Girls for Sexual and Reproductive Health Care: A Qualitative Study. *Afr J Reprod Health*. 2017 Dec;21(4):80-92. doi: 10.29063/ajrh2017/v21i4.9.
 23. Yazdkhasti M1, Pirak A. The effect of aromatherapy with lavender essence on severity of labor pain and duration of labor in primiparous women. *Complement Ther Clin Pract*. 2016 Nov;25:81-86. doi: 10.1016/j.ctcp.2016.08.008.
 24. Bagheri-Nesami M, Esmaili R, Tajari M. INTRAVENOUS MEDICATION ADMINISTRATION ERRORS AND THEIR CAUSES IN CARDIAC CRITICAL CARE UNITS IN IRAN. *Mater Sociomed*. 2015 Dec;27(6):442-6. doi: 10.5455/msm.2015.27.442-446.
 25. Sharifi M, Chitgar MG, Ghadamyari M, Sajedi RH, Amiri S. Characterisation of digestive protease in the rose sawfly, *Arge rosae* Linnaeus (Hymenoptera: Argidae), *Archives of Phytopathology and Plant Protection*, (2012), 45:10, 1170-1182, DOI: 10.1080/03235408.2012.658654.
 26. Gholamzadeh Chitgar M, Ghadamyari M. Effects of amitraz on the parasitoid *Encarsia formosa* (gahan) (hymenoptera: Aphelinidae) for control of *Trialetrodes vaporariorum* westwood (homoptera: Aleyrodidae): IOBC methods. *J. Entomol. Res. Soc*2012,14(2): 61-69.
 27. Gholamzadeh Chitgar M, Khosravi R, JalaliSendi J, Ghadamyari M. Sublethal effects of *Thymus vulgaris* essential oil on life-table parameters of two-spotted spider mite, *Tetranychus urticae* Koch (Acari: Tetranychidae), *Archives of Phytopathology and Plant Protection*,2013, 46:7, 781-788, DOI: 10.1080/03235408.2012.752143.
 28. Mojtaba Miladina, Amir Hosein Pishgooie, Shahla Aliyari, Elham Mousavi Nouri. The Comparison of the Effect of Two Complementary Medicine Methods (Music Therapy and Massage Therapy) On Postoperative Acute Pain After Abdominal Surgery: A Randomized Clinical Trial Study. *Iran Red Crescent Med J*. 2017 ;19(6):e14974. doi: 10.5812/ircmj.14974.
 29. Pishgooie AH, Shorakie HZ, Zareiyan A, Shoorideh FA, Forghani M. Evaluation of quality of life and its effective factors in military patients and relatives with coronary artery disease in Tehran, Iran, 2016– a cross sectional study. *Journal of Military Medicine*. 2018, Volume20, Issue 2. Pages:212-221.
 30. Talesh KT, Motamedi MH, Yazdani J, Ghavimi A, Ghoreishzadeh A. Prevention of relapse following intraoral vertical ramus osteotomy mandibular setback: can coronoidotomy help?. *Oral Surg Oral Med Oral Pathol*

- Oral Radiol Endod. 2011 May;111(5):557-60. doi: 10.1016/j.tripleo.2010.06.019.
31. Samiei M, Shahi S, Abdollahi AA, Eskandarinezhad M, Negahdari R, Pakseresht Z. The Antibacterial Efficacy of Photo-Activated Disinfection, Chlorhexidine and Sodium Hypochlorite in Infected Root Canals: An in Vitro Study. *Iran Endod J.* 2016 Summer;11(3):179-83. doi: 10.7508/icj.2016.03.006.
 32. Afkar AH, Mahbobubi M, Neyakan Shahri M, Mohammadi M, Jalilian F, Moradi F. Investigation of the relationship between illogical thoughts and dependence on others and marriage compatibility in the Iranian Veterans exposed to chemicals in Iran-Iraq War. *Glob J Health Sci.* 2014 May 8;6(5):38-45. doi: 10.5539/gjhs.v6n5p38.
 33. Kariman, N., Afrakhte, M., Hedayati, M., Fallahian, M., Alavi Majd, H. Diagnosis of premature rupture of membranes by assessment of urea and creatinine in vaginal washing fluid. 2013. *Iranian Journal of Reproductive Medicine.* 11(2), pp. 93-100.
 34. Nasimfar A, Sadeghi E, Amuzmehr A. Evaluation of knowledge, attitude, and practice of parents on the use of antibiotics for acute upper respiratory tract infections in children admitted to Motahari Hospital of Urmia in 2017–2018. *Asian Journal of Pharmaceutics,* Apr-Jun 2018 (Suppl), 12 (2) , S558.
 35. Khalatbari MR, Borghei-Razavi H, Samadian, M, Moharamzad Y, Schick U. Isolated primary craniopharyngioma in the cerebellopontine angle. 2012. *Journal of Clinical Neuroscience.* 19(11), pp. 1516-1519.
 36. Thompson PA, Kitchens RL. Native high-density lipoprotein augments monocyte responses to lipopolysaccharide by suppressing the inhibitory activity of LPSbinding protein. *J Immunol* 2006; 177:4880–7.
 37. van Leeuwen HJ, Heezius ECJM, Dallinga GM, van Strijp JA, Verhoef J, van Kessel KP. Lipoprotein metabolism in patients with severe sepsis. *Crit Care Med* 2003; 31:1359–66.
 38. Tsai MH, Peng YS, Chen YC, Lien JM, Tian YC, Fang JT, et al. Low serum concentration of apolipoprotein A-I is an indicator of poor prognosis in cirrhotic patients with severe sepsis. *J Hepatol* 2009;50(5):906–15.
 39. Chien JY, Jerng JS, Jen Yu C, Yang PC. Low serum level of highdensity lipoprotein cholesterol is a poor prognostic factor for severe sepsis. *Crit Care Med* 2005; 33:1688- 93.
 40. Levels JH, Abraham PR, van den Ende A, van Deventer SJ. Distribution and kinetics of lipoprotein-bound endotoxin. *Infect Immun* 2001; 69:2821-8.
 41. Levels JH, Abraham PR, van Barreveld EP, Meijers JC, van Deventer SJ. Distribution and kinetics of lipoprotein-bound lipoteichoic acid. *Infect Immun* 2003; 71:3280- 4.
 42. Vesj CJ, Kitchens RL, Wolfbauer G, Albers JJ, Munford RS. Lipopolysaccharide-binding protein and phospholipid transfer protein release lipopolysaccharides from gram-negative bacterial membranes. *Infect Immun* 2000; 68:2410- 7.
 43. Ettinger WH, Varma VK, Sorci-Thomas M, et al. Cytokines decrease apolipoprotein accumulation in medium from Hep G2 cells. *Arterioscler Thromb* 1994; 14:8 - 13.
 44. Fraunberger P, Pilz G, Cremer P, Werdan K, Walli AK. Association of serum tumor necrosis factor levels with decrease of cholesterol during septic shock. *Shock* 1998; 10:359- 63.
 45. Po'voa P, Almeida E, Moreira P, et al. C-reactive protein as an indicator of sepsis. *Intensive Care Med* 1998; 24:1052- 6.
 46. Dong Q, Wright JR. Expression of C-reactive protein by alveolar macrophages. *J Immunol* 1996; 156:4815 - 20.
 47. Vosylius S, Sipylaite J, Ivaskevicius J. Determinants of outcome in elderly patients admitted to the intensive care unit. *Age Ageing* 2005; 34:157- 62.
 48. Hazrati E, Najafi M, Alizadeh H, Noorifard M and Saeedi A., Comparison Between Hospital Stay and Carbapenem Use in Patients with Carbapenem Resistant Infection of a Middle-Eastern ICU with Other Centers, *Journal of Current Research in Science,* 2015, Vol. 3, No. 6, 89-92.