

Biological Activities of Some Indian medicinal plants

Khandelwal Vinoth Kumar Megraj^{1*}, Koneri Raju², R Balaraman¹, Kandhavelu Meenakshisundaram³

1. Pharmacy Department, The MS university of Baroda, Vadodara, India.
2. Karnataka College of Pharmacy, Bangalore, India.
3. Department of Signal processing, Tampere University of Technology, Tampere, Finland.

*Corresponding author: vinoth.khandelwal@yahoo.com

ABSTRACT:

There is an increasing demand for the herbal drug treatment of various ailments and many plant drugs from Ayurvedic system are being explored globally. The biological activities from various clinical and preclinical studies have been included along with some patents arising from these plants. The original report acts as a quick reference for extracting the biological activities, specially the newly reported effects, of the following selected Indian medicinal plants *Adhatoda vasica*, *Aegle marmelos*, *Aloe vera*, *Andrographis paniculata*, *Asparagus adscendents*, *Cinnamomum tamala*, *Coriandrum sativum*, *Cuminum cyminum*, *Curcuma longa*, *Embllica officinalis*, *Glycyrrhiza glabra*, *Hemidesmus indicus*, *Mucuna pruriens*, *Phyllanthus niruri*, *Solanum nigrum*, *Syzygium aromaticum*, *Terminalia chebula*, *Tinospora cordifolia*, *Withania somnifera*, *Zingiber officinale*.

Keywords: Ayurveda, alternative medicine, biological activity, herbal drug, patent, pharmacology, plant extract.

INTRODUCTION:

It is a well known fact that the demand for the herbal drug treatment of various ailments is increasing and plant drugs from the ayurvedic system are being explored more, not only in India but also globally. As a result, many research studies are being undertaken and there is a need for an update and to put them together. This article aims to give a quick reference to the biological activities of few Indian medicinal plants. We have tried to summarize most of the reported activities directly from the recent original research papers, but the list is not exhaustive. The biological activities from various clinical and preclinical studies have been included, while other uses of plants like herbicidal or insecticidal use have not been included. The main use of all these

studies is commercial utilization, so we have tried to include few patents arising from these plants, it should be noted that most of the patents are formulations and their uses. Usually the formulations are poly herbal and may contain one or more of the mentioned plants, so to avoid repetition, the formulations are included under any one plant detail only.

Adhatoda vasica

Adhatoda vasica is a well-known in Ayurvedic and Unani medicine. It has been used for the treatment of various diseases and disorders, particularly for the respiratory tract ailments. Many studies have been conducted on Adhatoda vasica and some of the reported activities of the plant include anti-diabetic [1], anti-implantation [2], anti-phlogistic, anti-allergic [3], anti-tussive [4], anti-ulcer [5], antioxidant, anti-genotoxicity [6], anti-tubercular [7], abortifacient [8], radioprotective [9], bronchodilator [10], hepatoprotective [11]. Pahwa et.al, [12] studied the chronic toxicity of some compounds of *Adhatoda vasica*. The patents consisting Adhatoda vasica are used for treating asthma (US 6746694), allergy (US 6746694), malnutrition, chronic cough, cold (US 7247322).

Aegle marmelos

Aegle marmelos has been widely used and studied for various activities like, anti-genotoxicity [13], analgesic [14], anti-cancer [15], anti-diarrhoeal [16], anti-fungal [17], anti-dyslipidemic [18], anti-inflammatory, anti-pyretic [19], anti-bacterial: Salmonella typhi, Shigella dysenteriae [20, 21], anti-viral: coxsackieviruses [22], anti-diabetic, antioxidant [23], anti-thyroid [24], anti-diarrhoeal, anti-ulcer [25], anti-asthmatic: Histamine antagonist H1 [26], immunostimulant [27], cardioprotective [28], radioprotective [29], treatment of ascites carcinoma [30], irritable bowel syndrome [31]. Acute and subacute toxicity studies have been studied by Veerappan et al. [32]. The patented formulations containing Aegle marmelos are used for treatment of HIV/AIDS and fungal infections secondary to HIV (PCT IN2004/000367), hyperglycemia (US 6989160), gastric ulcer (US 6855347).

Aloe vera

Aloe vera is one of the greatly known herbs for its cosmetic use, and it has been studied for its use in other major complications like cancer [33], myocardial ischemia [34], ulcer, diabetes [35, 36], hemorrhagic shock [37], hepatotoxicity [38], inflammation [39], allergic rhinitis [40] and oxidative stress [41]. Moreover, Aloe posses various effects like hypotensive [42], neuroprotective [43], radioprotective [44], anti-leishmanial [45], anti-

fungal [46], immunomodulatory [47], wound healing [48], cathartic [49], anti-viral: herpes simplex virus type 2 [50], angiogenic [51], anti-angiogenic [52], as a vaginal contraceptive [53], and protection against sunburn and suntan [54]. Subchronic toxicity has been studied by Zhou et.al. [55]. Some adverse effects of Aloe are cytotoxicity [56], Henoch-Schonlein purpura [57], acute toxic hepatitis [58]. There are many patented formulations available for Aloe vera specially for cosmetic use, few other patents include Aloe vera gel for the treatment of dry eye syndrome (US 6013259), ointment for skin irritations and healing of wounds (US 4725438), nasal spray for cold (US 5840278), topical treatment of pseudofolliculitis barbae (US 5034221).

Andrographis paniculata

Andrographis paniculata has been reported to encompass activities like cardioprotective [59], anti-hyperglycemic, antioxidant [60], anti-angiogenic [61], anti-cancer, immunostimulatory [62], anti-inflammatory [63], anti-malarial [64], anti-microbial [65], anti-viral: herpes simplex virus type 1 [66], upper respiratory tract infection [67], cyclophosphamide-induced toxicity [68], hepatoprotective [69], filaricidal [70], hypotensive [71], activation of TRPV4 channels [72], induction of uterus relaxation [73], inhibition of platelet aggregation [74], prevention of atherosclerotic stenosis and restenosis [75], reduction of the symptoms of common cold [76]. Andrographis paniculata has produced male anti-fertility effects [77], while another study did not find any adverse effect on fertility [78]. The patents include formulations, which are used for cancer (US 6486196), autoimmune diseases, Alzheimer's disease (PCT EP2004/005516), and improvement of liver function (US 6759061).

Asparagus adscendens

Asparagus adscendens is known for its activities like anti-cancer [79], anti-filarial [80], anti-fungal [81], anti-stress and anti-inflammatory [82]. Geriforte is a herbal formulation with Asparagus adscendens as one of the main constituents and has been widely studied for its use gastric ulcer [83], menopausal syndrome [84], neurological disorders [85], radioprotection [86], senile dermatoses [87], for stimulation of insulin secretion [88].

Cinnamomum tamala

The important studies conducted on Cinnamomum tamala are for its hypolipidemic effects [89], anti-diabetic and antioxidant [90], anti-ulcer [91], anti-inflammatory [92], anti-diarrhoeal [93], immunosuppressive [94] and anti-bacterial [95]. Few formulations

containing *Cinnamomum tamala* has been patented and are used for the treatment of diabetes (US 5886029) and antidote to tobacco (US 20060137702).

Coriandrum sativum

Various activities of *Coriandrum sativum* include anthelmintic [96], anti-implantation [97], anti-microbial [98], anti-mutagenicity [99], antioxidant and hepatoprotective [100], anxiolytic, sedative, muscle relaxant [101], hypoglycemic, hypolipidemic, cardioprotective [102], anti-dote for lead intoxication [103]. It has produced anaphylaxis in a patient [104]. Some interesting uses of patented formulations containing *Coriandrum sativum* are cigarettes which aid to stop smoking (EP1201142), hair coloring (US 7172632), sleep inducing antacid (US 20030013639), treatment of HIV-associated conditions (US 20020182272), and gastric ulcer (US 20050089583).

Cuminum cyminum

Cuminum cyminum has varied biological activities, which include anti-diabetic, antioxidant [105], anti-bacterial [106], anti-fungal [107], bronchodilatory [108], hepatoprotective, renoprotective [109], chemopreventive [110], anti-epileptic [111], galactagogue [112], hypolipidemic [113], male anti-fertility [114], memory-enhancing and anti-stress [115]. Toxicity of *Cuminum cyminum* has been studied by Haroun et al. [116]. Few patents of *Cuminum cyminum* are used for potentiation of bioefficacy of anti-infectives (PCT IN2003/000110), relieve inflammation and smooth muscle contraction (US 20060188588), cigarette for controlling body weight (PCT IB2006/000936), management of gynecological disorders and to prevent/treat anemia due to excessive bleeding associated with menstrual disorders (US 6455077), treating hepatitis B virus or hepatitis C virus (US 20020160065).

Curcuma longa

Curcuma longa is one of the most commonly used ayurvedic drugs with many of the following activities like antioxidant [117], apoptotic [118], anti-depressant [119], anti-fungal [120], anti-platelet [121], anti-spasmodic [122], anti-arthritis [123], hypoglycemic [124], hypotensive [125], anti-bacterial [126], leishmaniacidal [127], anti-genotoxicity [128], cardioprotective [129], neuroprotective [130], wound healing [131], cytoprotective by induction of heat shock protein [132]. It is also known to be effective for the treatment of pulmonary fibrosis [133], diabetic nephropathy [134], hyperlipidemia [135], encephalomyelitis [136], cancer [137], gastric ulcers by blocking H₂ receptors [138], excitotoxicity by modulation of NMDA receptors [139]. Among the adverse effects, *Curcuma longa* has been reported to arrest spermatogenesis [140] and has produced

toxicity [123] and contact urticaria [141]. The patented pharmaceutical compositions are used for reducing hyperlipidemia, anti-platelet aggregation (US 4842859), wound healing (US 5401504), control of hypertension and hypercholesterolemia (US 6162438), hepatic and splenic disorders (US 20050074507), and for the treatment of osteoarthritis (US 6492429).

Emblica officinalis

Emblica officinalis has been known for its use against acute pancreatitis [142], age related dyslipidaemia [143], memory loss [144], atherosclerosis [145], metabolic syndrome [146], inflammation [147], cancer [148], cadmium induced toxicity [149], genotoxicity [150], oxidative stress [151], hyperthyroidism [152], ulcer [153] snake venom [154], tardive dyskinesia [155], wound healing [156]. It also possesses following activities like anti-pyretic, analgesic [157], anti-microbial [158], anti-fungal [159], anti-tussive [160], chemopreventive [161], immunostimulatory [162], hepatoprotective [163], cardioprotective [164], radioprotective [165], increase haemoglobin [166]. Few of the interesting polyherbal formulations available are EuMil, Panchagavya, Hyponidd, Kalpaamruthaa, Triphala, Pepticare, Immu-21, Ophthacare. Patented formulations are used as antioxidant (US 6124268), anti-migraine (EP1284744), anti-allergic (US 20030194452), hepatoprotective (CA 2480334), for treatment of iron-deficiency anemia (US 20030198695), protection of skin against sun-induced damage (US 20050089590), inhibition of platelet aggregation (US 6290996), topical anti-inflammatory/analgesic (PCT IB2002/001306), preventing alcohol-induced hangover (PCT IN2005/000186).

Glycyrrhiza glabra

Glycyrrhiza glabra has been studied for its biological actions, including hypocholesterolaemic [167], anti-cancer [168], anti-diabetic: PPAR- γ ligand binding [169], anti-allergic [170], anti-depressive [171], anti-arthritis [172], anti-ulcer: inhibit adhesion of *Helicobacter pylori* [173], antioxidant, anti-genotoxic, anti-inflammatory [174], anti-viral: encephalitis [175], anti-microbial [176], anti-nephritis [177], anti-thrombotic [178], anti-atopic dermatitis [179], radioprotective [180], memory enhancing [181]. The herbal formulation DHC-1 has been studied for its cardio and renoprotective activities [182], other polyherbal formulations available are Revitonil, Pepticare. Few of the patented formulations containing *Glycyrrhiza glabra* are used for treating AIDS (US 5683697), prostate carcinoma (US 5665393), arthritis (US 7416748), liver associated ailments (EP1207894), as a skin whitening composition (US 5609875).

Hemidesmus indicus

Various effects of HI, such as hypoglycemic, hypolipidemic [183], antioxidant, anti-thrombotic [184], tranquilizer [185], anti-inflammatory, anti-pyretic [186], anti-ulcerogenic [187], cardioprotective [188, 189], hepatoprotective [190], renoprotective [191], genotoxic and anti-genotoxic [192], radioprotective [193], neutralization of viper venom [194], anti-bacterial and larvicidal [195], have been reported. Patented formulations with *Hemidesmus indicus* as an ingredient is used for skin disorders (US 5693327) viral infections (US 6841174), inflammation (PCT/IN2004/000148), renoprotection, and lipid lowering (PCT/IN2004/000300).

Mucuna pruriens

Some of the reported activities include hypoglycemic [196], hypocholesterolemic [197], antioxidant [198], aphrodisiac [199], neuroprotective [200], anti-Asiatic cobra venom [201], inhibition of hyperprolactinemia in man [202], management of Parkinson's disease [203]. Adverse effects like acute toxic psychosis [204], pruritus [205], has been reported. The patents of *Mucuna pruriens* are used for treatment of the nervous system, including Parkinson's disease (US 6106839) and potentiation of growth hormone (US 6340474).

Phyllanthus niruri

The known activities of *Phyllanthus niruri* are anti-babesial, anti-plasmodial [206], anti-hyperuricemic [207], anti-nociceptive [208], anti-HIV [209], antioxidant, hepatoprotective [210], vasorelaxant [211] and lipid lowering [212]. It is also used for inhibition of platelet aggregation [213], urolithiasis [214], hepatitis B virus, Woodchuck hepatitis virus [215]. The patents are pharmaceutical preparations for treating infections caused by hepatitis B virus (US 4673575), hepatitis C virus (EP1294387), adenoviruses (US 20060193907) and for treating chronic inflammatory, fibrotic processes (US 6586015).

Solanum nigrum

Solanum nigrum has been studied for its acetylcholine-like activity [216], anti-nociceptive, anti-inflammatory, anti-pyretic [217], antioxidant [218], anti-tumor [219], anti-ulcerogenic [220], anti-viral [221], hepatoprotective [222], cytoprotective [223], hypocholesterolemic [224], hypotensive [225], anti-convulsant [226]. Some alkaloids of *Solanum nigrum* have been shown to produce teratogenesis in frog [227]. The patents of *Solanum nigrum* comprise a skin treatment composition (US 20050152867), hepatoprotective and weight gain promoter (PCT IN2001/000178), formula providing

symptomatic relief for various diseases and especially for patients diagnosed with AIDS/ARC (US 20070041993).

Syzygium aromaticum

Few of the activities of *Syzygium aromaticum* are anti-carcinogenic effects [228], antioxidant [229], anti-herpesvirus [230], anti-thrombotic [231], anti-cytomegalovirus infection [232], anti-candidial [233], aphrodisiac [234], chemopreventive in lung cancer [235], inhibition of immediate hypersensitivity [236]. *Syzygium aromaticum* has been found to be cytotoxic to human skin cells [237]. A good antioxidant activity of a polyherbal composition containing *Syzygium aromaticum* has been observed in our lab [238]. *Syzygium aromaticum* is a composition of the following patents which are used for lowering cholesterol, treating arthritis, reducing blood pressure and Alzheimer's disease (US 5707631), bronchial asthma (PCT IN2002/000041), HIV infection (US 7160561).

Terminalia chebula

Terminalia chebula is also one of the widely used plants in Ayurvedic medicine. Few of the reported activities include anti-cancer [239], immunosuppressive [240], hepatoprotective [241], cardioprotective [242], radioprotective [243], anti-diabetic, renoprotective effects [244], anti- HIV [245], purgative [246], antioxidant [247], anti-aging [248], anti-ulcer: *Helicobacter pylori* [249], anti-typhoid [250], anti-caries [251], for wound healing [252], increasing gastric emptying [253]. *Triphala* is a widely used polyherbal formulation for its varied effects, which include hypolipidemic [254], immunomodulatory [255], anti-mutagenic [256], anti-bacterial [257], anti-diabetic, antioxidant [258]. *Terminalia chebula* has adverse effects like hepatic and renal toxicity [259], while it has also been shown to be hepato and renoprotective [241, 244]. Other polyherbal compositions containing *Terminalia chebula* are *Arogh*, *Rhinax*, *Triphala*. The patented compositions are used for the treatment of AIDS, flu, TB (US 5529778; US 5980903), immunocompromised conditions (US 20070122496).

Tinospora cordifolia

Tinospora cordifolia has been studied for its actions like immunomodulatory, [260], anti-allergic rhinitis [261], anti-ulcer [262], cardioprotective [263], chemopreventive [264], hepatoprotective [265], adjuvant therapy in hyperreactive malarious splenomegaly [266], hypolipidaemic [267], neuroprotective [268], obstructive jaundice [269], oxidative stress [270], radioprotective [271]. Anti-fertility effects in male rats have been reported by Gupta et al. [272]. *Pepticare* a polyherbal formulation has shown a

good anti-ulcer and antioxidant activity [273], other formulations are Hyponidd, Immu-21. There are number of patents for *Tinospora cordifolia* and are used for the treatment of renal disorders (PCT IB2005/051103), cancer (US 6780441), AIDS (PCT IN2004/000180), immune system-modulated disorders (CA 2432488), retroviral infections (PCT IN2005/000246), adenocarcinoma of the prostate (PCT IB2003/001885), allergy (US 20060141067), as anti-anxiolytic, tranquilizer, and non-narcotic sedative (US 20070122495), anti-pyretic (US 20060141069), as a health protective herbal soft drink (PCT IB2002/005555), herbal biscuits - dietary supplement for lactating mothers (US 6517861).

Withania somnifera

The biological activities of *Withania somnifera* are anxiolytic-anti-depressive [274], anti-fungal [275], anti-malarial [276], apoptotic [277], chondroprotective [278], cardioprotective [279], immunomodulatory [280], neuroprotective [281], inhibition of COX-2 enzyme [282], promoter of learning and memory in Alzheimer's disease [283]. Sharada et.al. [284] have studied toxicity of *Withania somnifera* root extract in rats and mice. The patents of the formulations containing *Withania somnifera* are used for increasing male sperm count (US 6866872), for rapid weight loss, controlling appetite, managing stress, supporting relaxation, combating fatigue and supporting mental well-being (US 20060286183), for refractory leukemias and lymphomas (US 20050058722), for cyclooxygenase-2 inhibition (US 20060247182), recovering hypofertility (EP1208845), for treatment and maintenance of hormone dependent conditions, immunostimulation (US 20070122494), as memory enhancer in Alzheimer's condition (PCT IN2004/000402), as a protection against UV irradiation (US 20060292094).

Zingiber officinale

Zingiber officinale is widely used by pregnant women for nausea and vomiting. It has been studied for various activities like cardioprotective [285], anti-convulsant, anxiolytic, anti-emetic [286], anti-diabetic, hypolipidaemic [287], anti-inflammatory [288], anti-thrombotic [289], anti-obesity [290], antioxidant [291], anti-tumor [288], anti-atherosclerotic [292], radioprotective [293], hypotensive [294], renoprotective [295], anti-reproductive toxicity [296], anti-ulcer [297], hepatoprotective [298], androgenic [299], anti-fouling [300], prokinetic [301]. inhibition of angiogenesis [302], increasing insulin sensitivity [303], molluscicidal, anti-schistosomal [304], nootropic [305], for treatment of gastrointestinal disorders [306], migraine [307], morning sickness [308]. Ginger is also non teratogenic [309]. The patented formulations of *Zingiber officinale* are used for alleviating symptoms associated with arthritis (US 5683698), inflammation,

platelet aggregation (US 6274177), relief of menopausal symptoms (US 5707630), treatment of migraine (US 6967033).

CONCLUSION:

As a consequence of increasing demand for the herbal drug treatment of various ailments, plant drugs from Ayurvedic system are being explored globally. This has resulted in many research studies with varied results, and hence there is a need to summarize them together. This review acts as a ready reference for biological activities of some Indian medicinal plants to the scientific community, in specific to researchers and students looking for sources of knowledge of medicinal plants and leads for new bioactive compounds. It is to be kept in mind that the reported activity may be shown by either the whole plant, or a part of the plant, or a particular extract, or isolated compounds. To know further about a biological action of a plant, it is advised to refer the individual research article.

We hope the article will be of immense use to the researchers and students, and result in increased interest in these medicinal plants.

ACKNOWLEDGEMENTS:

Authors thank Neelam Khandelwal for helping with editing and reviewing the manuscript.

REFERENCES:

1. Talib M, Gulfranz M, Mussaddeq Y. Effect of crude extract of *Adhatoda vasica* Nees on diabetic patients. *OnLine Journal of Biological Sciences* 2002;2: 436-437.
2. Prakash AO, Saxena V, Shukla S, Tewari RK, Mathur S, Gupta A, Sharma S, Mathur R. Anti-implantation activity of some indigenous plants in rats. *Acta Eur Fertil* 1985;16: 441-8.
3. Wagner H. Search for new plant constituents with potential antiphlogistic and antiallergic activity. *Planta Medica* 1989;55: 235-241.
4. Dhuley JN. Antitussive effect of *Adhatoda vasica* extract on mechanical or chemical stimulation-induced coughing in animals. *Journal of Ethnopharmacology* 1999;67: 361-5.
5. Shrivastava N, Srivastava A, Banerjee A, Nivsarkar M. Anti-ulcer activity of *Adhatoda vasica* Nees. *Journal of Herbal Pharmacotherapy* 2006;6: 43-49.
6. Jahangir T, Khan TH, Prasad L, Sultana S. Reversal of cadmium chloride-induced oxidative stress and genotoxicity by *Adhatoda vasica* extract in Swiss albino mice. *Biological Trace Element Research* 2006;111: 217-228.

7. Gupta R, Thakur B, Singh P, Singh HB, Sharma VD, Katoch VM, Chauhan SV. Anti-tuberculosis activity of selected medicinal plants against multi-drug resistant *Mycobacterium tuberculosis* isolates. *Indian J Med Res* 2010;131: 809-813.
8. Gupta OP, Anand KK, Ghatak BJ, Atal CK. Vasicine, alkaloid of *Adhatoda vasica*, a promising uterotonic abortifacient. *Indian J Exp Biol* 1978;16: 1075-7.
9. Kumar A, Ram J, Samarth RM, Kumar M. Modulatory influence of *Adhatoda vasica* Nees leaf extract against gamma irradiation in Swiss albino mice. *Phytomedicine* 2005;12: 285-93.
10. Amin AH, Mehta DR. Bronchodilator alkaloid (vasicinone) from *Adhatoda vasica* Nees. *Nature* 1959;184: 1317.
11. Jayakumar T, Ilayaraja M, Sridhar MP, Sivalokanathan S, Prasad TRB, Balasubramanian MP. Protective effect of *Adhatoda vasica* on D-galactosamine induced liver damage in rats with reference to lipid peroxidation and antioxidant status. *Journal of Natural Remedies* 2009;9: 91-98.
12. Pahwa GS, Zutshi U, Atal CK. Chronic toxicity studies with vasicine from *Adhatoda vasica* Nees. in rats and monkeys. *Indian J Exp Biol* 1987;25: 467-70.
13. Venkatesh P, Shantala B, Jagetia GC, Rao KK, Baliga MS. Modulation of Doxorubicin-induced genotoxicity by *Aegle marmelos* in mouse bone marrow: A micronucleus study. *Integrative Cancer Therapies* 2007;6: 42.
14. Shankarananth V, Balakrishnan N, Suresh D, Sureshpandian G, Edwin E, Sheeja E. Analgesic activity of methanol extract of *Aegle marmelos* leaves. *Fitoterapia* 2007;78: 258-259.
15. Costa-Lotufo LV, Khan MTH, Ather A, Wilke DV, Jimenez PC, Pessoa C, Moraes MEA, Moraes MO. Studies of the anticancer potential of plants used in Bangladeshi folk medicine. *Journal of Ethnopharmacology* 2005;99: 21-30.
16. Mazumder R, Bhattacharya S, Mazumder A, Pattnaik AK, Tiwary PM, Chaudhary S. Antidiarrhoeal evaluation of *Aegle Marmelos* (Correa) Linn. root extract. *Phytother Res* 2006;20: 82-4.
17. Rana BK, Singh UP, Taneja V. Antifungal activity and kinetics of inhibition by essential oil isolated from leaves of *Aegle marmelos*. *Journal of Ethnopharmacology* 1997;57: 29-34.
18. Narender T, Shweta S, Tiwari P, Papi Reddy K, Khaliq T, Prathipati P, Puri A, Srivastava AK, Chander R, Agarwal SC. Antihyperglycemic and antidyslipidemic agent from *Aegle marmelos*. *Bioorganic & Medicinal Chemistry Letters* 2007;17: 1808-1811.

19. Arul V, Miyazaki S, Dhananjayan R. Studies on the anti-inflammatory, antipyretic and analgesic properties of the leaves of *Aegle marmelos* Corr. *Journal of Ethnopharmacology* 2005;96: 159-163.
20. Rani P, Khullar N. Antimicrobial evaluation of some medicinal plants for their anti-enteric potential against multi-drug resistant *Salmonella typhi*. *Phytotherapy Research* 2004;18: 670-673.
21. Raja SB, Murali MR, Kumar NK, Devaraj SN. Isolation and Partial Characterisation of a Novel Lectin from *Aegle marmelos* Fruit and Its Effect on Adherence and Invasion of *Shigellae* to HT29 Cells. *PloS one* 2011;6: e16231.
22. Badam L, Bedekar SS, Sonawane KB, Joshi SP. In vitro antiviral activity of bael (*Aegle marmelos* Corr) upon human coxsackieviruses B1-B6. *J Commun Dis* 2002;34: 88-99.
23. Kamalakkannan N, Prince SMP. Antidiabetic and anti-oxidant activity of *Aegle marmelos* extract in streptozotocin-induced diabetic rats. *Pharmaceutical Biology* 2004;42: 125-130.
24. Panda S, Kar A. Evaluation of the antithyroid, antioxidative and antihyperglycemic activity of scopoletin from *Aegle marmelos* leaves in hyperthyroid rats. *Phytotherapy Research* 2006;20: 1103-1105.
25. Rao C, Amresh R, Irfan A, Rawat AKS, Pushpangadan P. Protective Effect of *Aegle marmelos* Fruit in Gastrointestinal Dysfunction in rats. *Pharmaceutical Biology* 2003;41: 558-563.
26. Arul V, Miyazaki S, Dhananjayan R. Mechanisms of the contractile effect of the alcoholic extract of *Aegle marmelos* Corr. on isolated guinea pig ileum and tracheal chain. *Phytomedicine* 2004;11: 679-83.
27. Patel P, Asdaq SMB. Immunomodulatory activity of methanolic fruit extract of *Aegle marmelos* in experimental animals. *Saudi Pharmaceutical Journal* 2010;18: 161-165.
28. Rajadurai M, Prince PSM. Comparative effects of *Aegle marmelos* extract and alpha-tocopherol on serum lipids, lipid peroxides and cardiac enzyme levels in rats with isoproterenol-induced myocardial infarction. *Singapore Medical Journal* 2005;46: 78-81.
29. Jagetia GC, Venkatesh P, Baliga MS. Evaluation of the radioprotective effect of *Aegle marmelos* (L.) Correa in cultured human peripheral blood lymphocytes exposed to different doses of {gamma}-radiation: a micronucleus study. *Mutagenesis* 2003;18: 387-393.

30. Jagetia GC, Venkatesh P, Baliga MS. Aegle marmelos (L.) CORREA Inhibits the Proliferation of Transplanted Ehrlich Ascites Carcinoma in Mice. *Biological & Pharmaceutical Bulletin* 2005;28: 58-64.
31. Yadav SK, Jain AK, Tripathi SN, Gupta JP. Irritable bowel syndrome: therapeutic evaluation of indigenous drugs. *Indian J Med Res* 1989;90: 496-503.
32. Veerappan A, Miyazaki S, Kadarkaraisamy M, Ranganathan D. Acute and subacute toxicity studies of Aegle marmelos Corr., an Indian medicinal plant. *Phytomedicine* 2007;14: 209-215.
33. Akev N, Turkay G, Can A, Gurel A, Yildiz F, Yardibi H, Ekiz EE, Uzun H. Effect of Aloe vera leaf pulp extract on Ehrlich ascites tumours in mice. *European Journal of Cancer Prevention* 2007;16: 151-157.
34. Sakai T, Repko BM, Griffith BP, Waters JH, Kameneva MV. IV infusion of a drag-reducing polymer extracted from aloe vera prolonged survival time in a rat model of acute myocardial ischaemia. *British Journal of Anaesthesia* 2007;98: 23.
35. Okyar A, Can A, Akev N, Baktir G, Sutlupinar N. Effect of Aloe vera leaves on blood glucose level in type I and type II diabetic rat models. *Phytother Res* 2001;15: 157-61.
36. Koo MWL. Aloe vera: antiulcer and antidiabetic effects. *Phytother Res* 1994;8: 461-464.
37. Macias CA, Kameneva MV, Tenhunen JJ, Puyana JC, Fink MP. Survival in a rat model of lethal hemorrhagic shock is prolonged following resuscitation with a small volume of a solution containing a drag-reducing polymer derived from Aloe vera. *Shock* 2004;22: 151-156.
38. Chandan BK, Saxena AK, Shukla S, Sharma N, Gupta DK, Suri KA, Suri J, Bhadauria M, Singh B. Hepatoprotective potential of Aloe barbadensis Mill. against carbon tetrachloride induced hepatotoxicity. *Journal of Ethnopharmacology* 2007;111: 560-566.
39. Langmead L, Makins RJ, Rampton DS. Anti-inflammatory effects of aloe vera gel in human colorectal mucosa in vitro. *Alimentary Pharmacology & Therapeutics* 2004;19: 521-527.
40. Yu H, Dong Z, Yang Z. Molecular biological study of aloe vera in the treatment of experimental allergic rhinitis in rat. *Lin Chuang Er Bi Yan Hou Ke Za Zhi* 2002;16: 229-31.
41. Saada HN, Ussama ZS, Mahdy AM. Effectiveness of Aloe vera on the antioxidant status of different tissues in irradiated rats. *Pharmazie* 2003;58: 929-931.

42. Saleem R, Faizi S, Siddiqui BS, Ahmed M, Hussain SA, Qazi A, Dar A, Ahmad SI, Qazi MH, Akhtar S. Hypotensive effect of chemical constituents from Aloe barbadensis. *Planta Med* 2001;67: 757-60.
43. Lin H, Lai CC, Chao P, Fan SS, Tsai Y, Huang SYI, Wan LEI, Tsai F. Aloe-Emodin Metabolites Protected N-Methyl-D-Aspartate-Treated Retinal Ganglion Cells by Cu-Zn Superoxide Dismutase. *JOURNAL OF OCULAR PHARMACOLOGY AND THERAPEUTICS* 2007;23.
44. Wang ZW, Zhou JM, Huang ZS, Yang AP, Liu ZC, Xia YF, Zeng YX, Zhu XF. Aloe polysaccharides mediated radioprotective effect through the inhibition of apoptosis. *Journal of Radiation Research* 2004;45: 447-454.
45. Dutta A, Mandal G, Mandal C, Chatterjee M. In vitro antileishmanial activity of Aloe vera leaf exudate: A potential herbal therapy in leishmaniasis. *Glycoconjugate Journal* 2007;24: 81-86.
46. Rosca-Casian O, Parvu M, Vlase L, Tamas M. Antifungal activity of Aloe vera leaves. *Fitoterapia* 2007;78: 219-222.
47. Zhang X-f, Wang H-m, Song Y-l, Nie L-h, Wang L-f, Liu B, Shen P-p, Liu Y. Isolation, structure elucidation, antioxidative and immunomodulatory properties of two novel dihydrocoumarins from Aloe vera. *Bioorganic & Medicinal Chemistry Letters* 2006;16: 949-953.
48. Atiba A, Nishimura M, Kakinuma S, Hiraoka T, Goryo M, Shimada Y, Ueno H, Uzuka Y. Aloe vera oral administration accelerates acute radiation-delayed wound healing by stimulating transforming growth factor-[beta] and fibroblast growth factor production. *The American Journal of Surgery* 2011;201: 809-818.
49. Ishii Y, Tanizawa H, Takino Y. Studies of aloe. V. Mechanism of cathartic effect. *Biol Pharm Bull* 1994;17: 651-3.
50. Zandi K, Zadeh MA, Sartavi K, Rastian Z. Antiviral activity of Aloe vera against herpes simplex virus type 2: An in vitro study. *African Journal of Biotechnology* 2007;6: 1770-1773.
51. Moon EJ, Lee YM, Lee OH, Lee MJ, Lee SK, Chung MH, Park YI, Sung CK, Choi JS, Kim KW. A novel angiogenic factor derived from Aloe vera gel: β -sitosterol, a plant sterol. *Angiogenesis* 1999;3: 117-123.
52. Cárdenas C, Quesada AR, Medina MA. Evaluation of the anti-angiogenic effect of aloe-emodin. *Cellular and Molecular Life Sciences (CMLS)* 2006;63: 3083-3089.
53. Fahim MS, Wang M. Zinc acetate and lyophilized aloe barbadensis as vaginal contraceptive. *Contraception* 1996;53: 231-6.
54. Puvabanditsin P, Vongtongsri R. Efficacy of aloe vera cream in prevention and treatment of sunburn and suntan. *J Med Assoc Thai* 2005;88: S173-6.

55. Zhou Y, Feng Y, Wang H, Yang H. 90-day subchronic toxicity study of aloe whole-leaf powder. *Wei Sheng Yan Jiu* 2003;32: 590-3.
56. Avila H, Rivero J, Herrera F, Fraile G. Cytotoxicity of a low molecular weight fraction from Aloe vera (*Aloe barbadensis* Miller) gel. *Toxicon* 1997;35: 1423-30.
57. Evangelos C, Spyros K, Spyros D. Henoch-Schonlein purpura associated with Aloe vera administration. *Eur J Intern Med* 2005;16: 59-60.
58. Kanat O, Ozet A, Ataergin S. Aloe vera-induced acute toxic hepatitis in a healthy young man. *European Journal of Internal Medicine* 2006;17: 589.
59. Guo ZL, Zhao HY, Zheng XH. The effect of andrographis paniculata nees (APN) in alleviating the myocardial ischemic reperfusion injury. *J Tongji Med Univ* 1994;14: 49-51.
60. Zhang XF, Tan BKH. Antihyperglycaemic And Anti-Oxidant Properties Of Andrographis Paniculata In Normal And Diabetic Rats. *Clinical and Experimental Pharmacology and Physiology* 2000;27: 358-363.
61. Sheeja K, Guruvayoorappan C, Kuttan G. Antiangiogenic activity of Andrographis paniculata extract and andrographolide. *Int Immunopharmacol* 2007;7: 211-21.
62. Kumar RA, Sridevi K, Kumar NV, Nanduri S, Rajagopal S. Anticancer and immunostimulatory compounds from Andrographis paniculata. *Journal of Ethnopharmacology* 2004;92: 291-295.
63. Madav S, Tandan SK, Lal J, Tripathi HC. Anti-inflammatory activity of andrographolide. *Fitoterapia(Milano)* 1996;67: 452-458.
64. Najila MJS, Rain AN, Kamel AGM, Zahir SIS, Khozirah S, Hakim SL, Zakiah I, Azizol AK. The screening of extracts from *Goniothalamus scortechinii*, *Aralidium pinnatifidum* and *Andrographis paniculata* for anti-malarial activity using the lactate dehydrogenase assay. *Journal of Ethnopharmacology* 2002;82: 239-242.
65. Singha PK, Roy S, Dey S. Antimicrobial activity of *Andrographis paniculata*. *Fitoterapia* 2003;74: 692-4.
66. Wiert C, Kumar K, Yusof MY, Hamimah H, Fauzi ZM, Sulaiman M. Antiviral properties of ent-labdene diterpenes of *Andrographis paniculata* nees, inhibitors of herpes simplex virus type 1. *Phytother Res* 2005;19: 1069-70.
67. Saxena RC, Singh R, Kumar P, Yadav SC, Negi MPS, Saxena VS, Joshua AJ, Vijayabalaji V, Goudar KS, Venkateshwarlu K. A randomized double blind placebo controlled clinical evaluation of extract of *Andrographis paniculata* (KalmCold (TM)) in patients with uncomplicated upper respiratory tract infection. *Phytomedicine* 2010;17: 178-185.

68. Sheeja K, Kuttan G. Ameliorating effects of andrographis paniculata extract against cyclophosphamide-induced toxicity in mice. *Asian Pac J Cancer Prev* 2006;7: 609-14.
69. Singha PK, Roy S, Dey S. Protective activity of andrographolide and arabinogalactan proteins from *Andrographis paniculata* Nees. against ethanol-induced toxicity in mice. *Journal of Ethnopharmacology* 2007;111: 13-21.
70. Dutta A, Sukul NC. Filaricidal properties of a wild herb, *Andrographis paniculata*. *J Helminthol* 1982;56: 81-4.
71. Zhang CY, Tan BK. Mechanisms of cardiovascular activity of *Andrographis paniculata* in the anaesthetized rat. *Journal of Ethnopharmacology* 1997;56: 97-101.
72. Smith PL, Maloney KN, Pothen RG, Clardy J, Clapham DE. Bisandrographolide from *Andrographis paniculata* activates TRPV4 channels. *Journal of Biological Chemistry* 2006;281: 29897-29904.
73. Burgos RA, Aguila MJ, Santiesteban ET, Sánchez NS, Hancke JL. *Andrographis paniculata* (Nees) Induces Relaxation of Uterus by Blocking Voltage Operated Calcium Channels and Inhibits Ca²⁺ Influx. *Phytother. Res* 2001;15: 235-239.
74. Thisoda P, Rangkadilok N, Pholphana N, Worasuttayangkurn L, Ruchirawat S, Satayavivad J. Inhibitory effect of *Andrographis paniculata* extract and its active diterpenoids on platelet aggregation. *European Journal of Pharmacology* 2006;553: 39-45.
75. Wang DW, Zhao HY. Prevention of atherosclerotic arterial stenosis and restenosis after angioplasty with *Andrographis paniculata* nees and fish oil. *Experimental studies of effects and mechanisms. Chin Med J (Engl)* 1994;107: 464-70.
76. Caceres DD, Hancke JL, Burgos RA, Sandberg F, Wikman GK. Use of visual analogue scale measurements (VAS) to assess the effectiveness of standardized *Andrographis paniculata* extract SHA-10 in reducing the symptoms of common cold. A randomized double blind-placebo study. *Phytomedicine* 1999;6: 217-23.
77. Akbarsha MA, Manivannan B, Hamid KS, Vijayan B. Antifertility effect of *Andrographis paniculata* (Nees) in male albino rat. *Indian J Exp Biol* 1990;28: 421-6.
78. Allan JJ, Pore MP, Deepak M, Murali B, Mayachari AS, Agarwal A. Reproductive and fertility effects of an extract of *Andrographis paniculata* in male Wistar rats. *International journal of toxicology* 2009;28: 308.
79. [79] Singh M, Singh S, Kale RK. Chemomodulatory potential of *Asparagus adscendens* against murine skin and forestomach papillomagenesis. *European Journal of Cancer Prevention* 2011;20: 240-247 10.1097/CEJ.0b013e3283447410.

80. Singh R, Khan NU, Singhal KC. Potential antifilarial activity of roots of *Asparagus adscendens* Roxb, against *Setaria cervi* in vitro. *Indian J Exp Biol* 1997;35: 168-72.
81. Singh R, Rai B. Antifungal potential of some higher plants against *Fusarium udum* causing wilt disease of *Cajanus cajan*. *Microbios* 2000;102: 165-73.
82. Kanwar AS, Bhutani KK. Effects of *Chlorophytum arundinaceum*, *Asparagus adscendens* and *Asparagus racemosus* on pro-inflammatory cytokine and corticosterone levels produced by stress. *Phytotherapy Research* 2010;24: 1562-1566.
83. Padma P, Chansouria JPN, Khosa RL. Effect of some indigenous drugs on cold immobilization stress induced gastric ulcer. *Phytotherapy Research* 1998;12: 127-128.
84. Saxena SC. Use of Geriforte in the Menopausal Syndrome. *Probe* 1978;1: 20-27.
85. Velmurugendran CU, Virudhagirinatha BS. Geriforte in Neurological Disorders. *Probe* 1980;1: 13-17.
86. Jagetia GC, Baliga MS. Evaluation of the radioprotective action of geriforte in mice exposed to different doses of γ -radiation. *American Journal of Chinese Medicine* 2004;32: 551-567.
87. Bhargava RK, Soni V. Geriforte—An Indigenous Drug in Senile Pruritus. *Antiseptic* 1989;86: 486.
88. Mathews JN, Flatt PR, Abdel-Wahab YH. *Asparagus adscendens* (Shweta musali) stimulates insulin secretion, insulin action and inhibits starch digestion. *British Journal of Nutrition* 2007;95: 576-581.
89. Dhulasavant V, Shinde S, Pawar M, Naikwade S. Antihyperlipidemic Activity of *Cinnamomum tamala* Nees. on High Cholesterol Diet Induced Hyperlipidemia. *Int. J. PharmTech Res* 2010;2: 2517-2521.
90. Chakraborty U, Das H. Antidiabetic and Antioxidant Activities of *Cinnamomum tamala* Leaf Extracts in Stz-Treated Diabetic Rats. *Global Journal of Biotechnology & Biochemistry* 2010;5: 12-18.
91. Eswaran MB, Surendran S, Vijayakumar M, Ojha SK, Rawat AKS, Rao CV. Gastroprotective activity of *Cinnamomum tamala* leaves on experimental gastric ulcers in rats. *Journal of Ethnopharmacology* 2010;128: 537-540.
92. Gambhire MN, Juvekar AR, Wankhede SS. Anti-inflammatory activity of aqueous extract of *Cinnamomum tamala* leaves by in vivo and in vitro methods. *Journal of Pharmacy Research* 2009;2: 1521-1524.
93. Rao C, Vijayakumar M, Sairam K, Kumar V. Antidiarrhoeal activity of the standardised extract of *Cinnamomum tamala* in experimental rats. *Journal of Natural Medicines* 2008;62: 396-402.

94. Chaurasia JK, Mishra A, Tripathi YB. Immunomodulation property of hexane fraction of leaves of *Cinnamomum tamala* Linn. in rats. *Cell Biochemistry and Function* 2010;28: 454-460.
95. Goyal P, Chauhan A, Kaushik P. Laboratory evaluation of crude extracts of *Cinnamomum tamala* for potential antibacterial activity. *Electronic Journal of Biology* 2009;5: 75-79.
96. Eguale T, Tilahun G, Debella A, Feleke A, Makonnen E. In vitro and in vivo anthelmintic activity of crude extracts of *Coriandrum sativum* against *Haemonchus contortus*. *Journal of Ethnopharmacology* 2007;110: 428-433.
97. Al-Said MS, Al-Khamis KI, Islam MW, Parmar NS, Tariq M, Ageel AM. Post-coital antifertility activity of the seeds of *Coriandrum sativum* in rats. *Journal of Ethnopharmacology* 1987;21: 165-73.
98. Begnami AF, Duarte MCT, Furletti V, Rehder VLG. Antimicrobial potential of *Coriandrum sativum* L. against different *Candida* species in vitro. *Food Chemistry* 2010;118: 74-77.
99. Cortes-Eslava J, Gomez-Arroyo S, Villalobos-Pietrini R, Espinosa-Aguirre JJ. Antimutagenicity of coriander (*Coriandrum sativum*) juice on the mutagenesis produced by plant metabolites of aromatic amines. *Toxicology Letters* 2004;153: 283-292.
100. Samojlik I, Lakic N, Mimica-Dukic N, akovic -S vajcer K, Boz in B. Antioxidant and hepatoprotective potential of essential oils of coriander (*Coriandrum sativum* L.) and caraway (*Carum carvi* L.)(Apiaceae). *Journal of Agricultural and Food Chemistry* 2010.
101. Emamghoreishi M, Khasaki M, Aazam MF. *Coriandrum sativum*: evaluation of its anxiolytic effect in the elevated plus-maze. *Journal of Ethnopharmacology* 2005;96: 365-370.
102. Aissaoui A, Zizi S, Israili ZH, Lyoussi B. Hypoglycemic and hypolipidemic effects of *Coriandrum sativum* L. in *Meriones shawi* rats. *Journal of Ethnopharmacology* 2011;In Press, Uncorrected Proof.
103. Aga M, Iwaki K, Ueda Y, Ushio S, Masaki N, Fukuda S, Kimoto T, Ikeda M, Kurimoto M. Preventive effect of *Coriandrum sativum* (Chinese parsley) on localized lead deposition in ICR mice. *Journal of Ethnopharmacology* 2001;77: 203-208.
104. Ebo DG, Bridts CH, Mertens MH, Stevens WJ. Coriander anaphylaxis in a spice grinder with undetected occupational allergy. *Acta Clin Belg* 2006;61: 152-6.
105. Jagtap AG, Patil PB. Antihyperglycemic activity and inhibition of advanced glycation end product formation by *Cuminum cyminum* in streptozotocin induced diabetic rats. *Food and Chemical Toxicology* 2010;48: 2030-2036.

106. Iacobellis NS, Cantore PL, Capasso F, Senatore F. Antibacterial activity of *Cuminum cyminum* L. and *Carum carvi* L. essential oils. *Journal of Agricultural and Food Chemistry* 2005;53: 57-61.
107. Romagnoli C, Andreotti E, Maietti S, Mahendra R, Mares D. Antifungal activity of essential oil from fruits of Indian *Cuminum cyminum*. *Pharmaceutical Biology* 2010;48: 834-838.
108. Boskabady M, Kiani S, Azizi H. Relaxant effect of *Cuminum cyminum* on guinea pig tracheal chains and its possible mechanism (s). *Indian Journal of Pharmacology* 2005;37: 111-115.
109. Aruna K, Rukkumani R, Varma PS, Menon VP. Therapeutic role of *Cuminum cyminum* on ethanol and thermally oxidized sunflower oil induced toxicity. *Phytotherapy Research* 2005;19: 416-421.
110. Gagandeep DS, Mendiz E, Rao AR, Kale RK. Chemopreventive effects of *Cuminum cyminum* in chemically induced forestomach and uterine cervix tumors in murine model systems. *Nutr Cancer* 2003;47: 171-80.
111. Janahmadi M, Niazi F, Danyali S, Kamalinejad M. Effects of the fruit essential oil of *Cuminum cyminum* Linn.(Apiaceae) on pentylenetetrazol-induced epileptiform activity in F1 neurones of *Helix aspersa*. *Journal of Ethnopharmacology* 2006;104: 278-282.
112. Agrawala IP, Achar MV, Boradkar RV, Roy N. Galactagogue action of *Cuminum cyminum* and *Nigella staiva*. *Indian J Med Res* 1968;56: 841-4.
113. Dhandapani S, Subramanian VR, Rajagopal S, Namasivayam N. Hypolipidemic effect of *Cuminum cyminum* L. On alloxan-induced diabetic rats. *Pharmacological Research* 2002;46: 251-255.
114. Gupta RS, Saxena P, Gupta R, Kachhawa JBS. Evaluation of reversible contraceptive activities of *Cuminum cyminum* in male albino rats. *Contraception* 2011;84: 98-107.
115. Koppula S, Choi DK. *Cuminum cyminum* extract attenuates scopolamine-induced memory loss and stress-induced urinary biochemical changes in rats: A noninvasive biochemical approach. *Pharmaceutical Biology* 2011;49: 702-708.
116. Haroun EM, Mahmoud OM, Adam SE. Effect of feeding *Cuminum cyminum* fruits, *Thymus vulgaris* leaves or their mixture to rats. *Vet Hum Toxicol* 2002;44: 67-9.
117. Nishinaka T, Ichijo Y, Ito M, Kimura M, Katsuyama M, Iwata K, Miura T, Terada T, Yabe-Nishimura C. Curcumin activates human glutathione S-transferase P1 expression through antioxidant response element. *Toxicology Letters* 2007;170: 238-247.

118. Anto RJ, Mukhopadhyay A, Denning K, Aggarwal BB. Curcumin (diferuloylmethane) induces apoptosis through activation of caspase-8, BID cleavage and cytochrome c release: its suppression by ectopic expression of Bcl-2 and Bcl-xl. *Carcinogenesis* 2002;23: 143-150.
119. Yu ZF, Kong LD, Chen Y. Antidepressant activity of aqueous extracts of *Curcuma longa* in mice. *Journal of Ethnopharmacology* 2002;83: 161-165.
120. Wuthi-udomlert M, Grisanapan W, Luanratana O, Caichompoo W. Antifungal activity of *Curcuma longa* grown in Thailand. *Southeast Asian J Trop Med Public Health* 2000;31: 178-82.
121. Lee HS. Antiplatelet property of *Curcuma longa* L. rhizome-derived ar-turmerone. *Bioresource Technology* 2006;97: 1372-1376.
122. Itthipanichpong C, Ruangrunsi N, Kemsri W, Sawasdipanich A. Antispasmodic effects of curcuminoids on isolated guinea-pig ileum and rat uterus. *J Med Assoc Thai* 2003;86: S299-309.
123. Funk JL, Frye JB, Oyarzo JN, Zhang H, Timmermann BN. Anti-Arthritic Effects and Toxicity of the Essential Oils of Turmeric (*Curcuma longa* L.). *Journal of Agricultural and Food Chemistry* 2009;58: 842-849.
124. Kuroda M, Mimaki Y, Nishiyama T, Mae T, Kishida H, Tsukagawa M, Takahashi K, Kawada T, Nakagawa K, Kitahara M. Hypoglycemic effects of turmeric (*Curcuma longa* L. Rhizomes) on genetically diabetic KK-A y mice. *Biological & Pharmaceutical Bulletin* 2005;28: 937-939.
125. Goto H, Sasaki Y, Fushimi H, Shibahara N, Shimada Y, Komatsu K. Effect of *Curcuma* herbs on vasomotion and hemorheology in spontaneously hypertensive rat. *American Journal of Chinese Medicine* 2005;33: 449-457.
126. Singh R, Chandra R, Bose M, Luthra PM. Antibacterial activity of *Curcuma longa* rhizome extract on pathogenic bacteria. *Current science(Bangalore)* 2002;83: 737-740.
127. Saleheen D, Ali SA, Ashfaq K, Siddiqui AA, Agha A, Yasinzai MM. Latent Activity of Curcumin against Leishmaniasis in Vitro. *Biological & Pharmaceutical Bulletin* 2002;25: 386-389.
128. El-Ashmawy IM, Ashry KM, El-Nahas AF, Salama OM. Protection by Turmeric and Myrrh against Liver Oxidative Damage and Genotoxicity Induced by Lead Acetate in Mice. *Basic & Clinical Pharmacology & Toxicology* 2006;98: 32-37.
129. Mohanty I, Singh Arya D, Dinda A, Joshi S, Talwar KK, Gupta SK. Protective effects of *Curcuma longa* on ischemia-reperfusion induced myocardial injuries and their mechanisms. *Life Sci* 2004;75: 1701-11.

130. Zbarsky V, Datla KP, Parkar S, Rai DK, Aruoma OI, Dexter DT. Neuroprotective properties of the natural phenolic antioxidants curcumin and naringenin but not quercetin and fisetin in a 6-OHDA model of Parkinson's disease. *Free Radical Research* 2005;39: 1119-1125.
131. Phan TT, See P, Lee ST, Chan SY. Protective Effects of Curcumin against Oxidative Damage on Skin Cells In Vitro: Its Implication for Wound Healing. *The Journal of Trauma: Injury, Infection, and Critical Care* 2001;51: 927-931.
132. Dunsmore KE, Chen PG, Wong HR. Curcumin, a medicinal herbal compound capable of inducing the heat shock response. *Crit Care Med* 2001;29: 2199-204.
133. Punithavathi D, Venkatesan N, Babu M. Protective effects of curcumin against amiodarone-induced pulmonary fibrosis in rats. *British Journal of Pharmacology* 2003;139: 1342-1350.
134. Babu PS, Srinivasan K. Amelioration of renal lesions associated with diabetes by dietary curcumin in streptozotocin diabetic rats. *Molecular and Cellular Biochemistry* 1998;181: 87-96.
135. Rukkumani R, Balasubashini MS, Menon VP. Protective effects of curcumin and photo-irradiated curcumin on circulatory lipids and lipid peroxidation products in alcohol and polyunsaturated fatty acid-induced toxicity. *Phytotherapy Research* 2003;17: 925-929.
136. Natarajan C, Bright JJ. Curcumin Inhibits experimental allergic encephalomyelitis by blocking IL-12 signaling through Janus kinase-STAT Pathway in T lymphocytes 1. *The Journal of Immunology* 2002;168: 6506-6513.
137. Cho WCSCS, Bar-Sela G, Schaffer M. An Evidence-Based Perspective of *Curcuma Longa* (Turmeric) for Cancer Patients. In: *Evidence-based Anticancer Materia Medica*: Springer Netherlands; 2011, p. 225-243.
138. Kim DC, Kim SH, Choi BH, Baek NI, Kim D, Kim MJ, Kim KT. *Curcuma longa* extract protects against gastric ulcers by blocking H₂ histamine receptors. *Biological & Pharmaceutical Bulletin* 2005;28: 2220-2224.
139. Matteucci A, Frank C, Domenici MR, Balduzzi M, Paradisi S, Carnovale-Scalzo G, Scorcio G, Malchiodi-Albedi F. Curcumin treatment protects rat retinal neurons against excitotoxicity: effect on N-methyl-D-aspartate-induced intracellular Ca²⁺ increase. *Experimental Brain Research* 2005;167: 641-648.
140. Ashok P, Meenakshi B. Contraceptive effect of *Curcuma longa* (L.) in male albino rat. *Asian Journal of Andrology* 2004;6: 71-74.
141. Liddle M, Hull C, Liu C, Powell D. Contact urticaria from Curcumin. *Dermatitis* 2004;2006: 196-7.

142. Thorat SP, Rege NN, Naik AS, Thatte UM, Joshi A, Panicker KN, Bapat RD, Dahanukar SA. *Emblica officinalis*: a novel therapy for acute pancreatitis--an experimental study. *HPB Surg* 1995;9: 25-30.
143. Yokozawa T, Young Kim H, Ju Kim H, Okubo T, Chu DC, Raj Juneja L. *Amla* (*Emblica officinalis* Gaertn.) prevents dyslipidaemia and oxidative stress in the ageing process. *British Journal of Nutrition* 2007;97: 1187-1195.
144. Vasudevan M, Parle M. Memory enhancing activity of *Anwala churna* (*Emblica officinalis* Gaertn.): An Ayurvedic preparation. *Physiology & Behavior* 2007;91: 46-54.
145. Thakur CP, Mandal K. Effect of *Emblica officinalis* on cholesterol-induced atherosclerosis in rabbits. *Indian J Med Res* 1984;79: 142-6.
146. Kim HY, Okubo T, Juneja LR, Yokozawa T. The protective role of *amla* (*Emblica officinalis* Gaertn.) against fructose-induced metabolic syndrome in a rat model. *British Journal of Nutrition* 2010;103: 502-512.
147. Asmawi MZ, Kankaanranta H, Moilanen E, Vapaatalo H. Anti-inflammatory activities of *Emblica officinalis* Gaertn leaf extracts. *J Pharm Pharmacol* 1993;45: 581-4.
148. Jose JK, Kuttan G, Kuttan R. Antitumour activity of *Emblica officinalis*. *Journal of Ethnopharmacology* 2001;75: 65-69.
149. Khandelwal S, Shukla LJ, Shanker R. Modulation of acute cadmium toxicity by *Emblica officinalis* fruit in rat. *Indian J Exp Biol* 2002;40: 564-70.
150. Banu SM, Selvendiran K, Singh J, Sakthisekaran D. Protective effect of *Emblica officinalis* ethanolic extract against 7, 12-dimethylbenz (a) anthracene (DMBA) induced genotoxicity in Swiss albino mice. *Human & Experimental Toxicology* 2004;23: 527.
151. Pozharitskaya ON, Ivanova SA, Shikov AN, Makarov VG. Original Paper Separation and evaluation of free radical-scavenging activity of phenol components of *Emblica officinalis* extract by using an HPTLC-DPPH method. *J. Sep. Sci* 2007;30: 1250-1254.
152. Panda S, Kar A. Fruit extract of *Emblica officinalis* ameliorates hyperthyroidism and hepatic lipid peroxidation in mice. *Pharmazie* 2003;58: 753-5.
153. Sairam K, Rao CV, Babu MD, Kumar KV, Agrawal VK, Goel RK. Antiulcerogenic effect of methanolic extract of *Emblica officinalis*: An experimental study. *Journal of Ethnopharmacology* 2002;82: 1-9.
154. Alam MI, Gomes A. Snake venom neutralization by Indian medicinal plants (*Vitex negundo* and *Emblica officinalis*) root extracts. *Journal of Ethnopharmacology* 2003;86: 75-80.

155. Bhattachary SK, Bhattacharya D, Muruganandam AV. Effect of *Emblica officinalis* tannoids on a rat model of tardive dyskinesia. *Indian J Exp Biol* 2000;38: 945-7.
156. Sumitra M, Manikandan P, Gayathri VS, Mahendran P, Suguna L. *Emblica officinalis* exerts wound healing action through up-regulation of collagen and extracellular signal-regulated kinases (ERK1/2). *Wound Repair and Regeneration* 2009;17: 99-107.
157. Perianayagam JB, Sharma SK, Joseph A, Christina AJM. Evaluation of anti-pyretic and analgesic activity of *Emblica officinalis* Gaertn. *Journal of Ethnopharmacology* 2004;95: 83-85.
158. Vimala Y, Rachel KV, Sankar AU. Antimicrobial activity of *Emblica officinalis* seeds on human pathogens. *Journal of Pure and Applied Microbiology* 2009;3: 219-222.
159. Dutta BK, Rahman I, Das TK. Antifungal activity of Indian plant extracts. *Mycoses* 1998;41: 535-6.
160. NosalOva G. Antitussive activity of the fruit extract of *Emblica officinalis* Gaertn.(Euphorbiaceae). *Phytomedicine* 2003;10: 583-589.
161. Sancheti G, Jindal A, Kumari R, Goyal PK. Chemopreventive action of *emblica officinalis* on skin carcinogenesis in mice. *Asian Pac J Cancer Prev* 2005;6: 197-201.
162. Suja RS, Nair AMC, Sujith S, Preethy J, Deepa AK. Evaluation of immunomodulatory potential of *Emblica officinalis* fruit pulp extract in mice. *Indian Journal of Animal Research* 2009;43: 103-106.
163. Sultana S, Ahmad S, Khan N, Jahangir T. Effect of *Emblica officinalis* (Gaertn) on CCl₄ induced hepatic toxicity and DNA synthesis in Wistar rats. *Indian J Exp Biol* 2005;43: 430-6.
164. Rajak S, Banerjee SK, Sood S, Dinda AK, Gupta YK, Gupta SK, Maulik SK. *Emblica officinalis* causes myocardial adaptation and protects against oxidative stress in ischemic-reperfusion injury in rats. *Phytotherapy Research* 2004;18: 54-60.
165. Singh I, Sharma A, Nunia V, Goyal PK. Radioprotection of Swiss albino mice by *Emblica officinalis*. *Phytotherapy Research* 2005;19: 444-446.
166. Anila L, Vijayalakshmi NR. Beneficial effects of flavonoids from *Sesamum indicum*, *Emblica officinalis* and *Momordica charantia*. *Phytotherapy Research* 2000;14: 592-595.
167. Visavadiya NP, Narasimhacharya AV. Hypocholesterolaemic and antioxidant effects of *Glycyrrhiza glabra* (Linn) in rats. *Mol Nutr Food Res* 2006;50: 1080-6.

168. Rafi MM. Elucidating the role of nutraceuticals in overexpressing antiapoptotic proteins in prostate cancer. *Nutrition* 2004;20: 78-82.
169. Kuroda M, Mimaki Y, Honda S, Tanaka H, Yokota S, Mae T. Phenolics from *Glycyrrhiza glabra* roots and their PPAR-[gamma] ligand-binding activity. *Bioorganic & Medicinal Chemistry* 2010;18: 962-970.
170. Shin Y, Bae E, Lee B, Lee SH, Kim JA, Kim Y, Kim D. In vitro and in vivo antiallergic effects of *Glycyrrhiza glabra* and its components. *Planta Medica* 2007;73: 257.
171. Dhingra D, Sharma A. Antidepressant-like activity of *Glycyrrhiza glabra* L. in mouse models of immobility tests. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 2006;30: 449-454.
172. Gujral ML, Sareen KN, Tangri KK, Roy AK, Gupta GP, Amma MK. Antiarthritic activity of *glycyrrhiza glabra* Linn. *Indian J Psychol* 1959;3: 39-47.
173. Wittschier N, Faller G, Hensel A. Aqueous extracts and polysaccharides from Licorice roots (*Glycyrrhiza glabra* L.) inhibit adhesion of *Helicobacter pylori* to human gastric mucosa. *Journal of Ethnopharmacology* 2009;125: 218-223.
174. Siracusa L, Saija A, Cristani M, Cimino F, D'Arrigo M, Trombetta D, Rao F, Ruberto G. Phytocomplexes from licorice (*Glycyrrhiza glabra* L.) leaves -- Chemical characterization and evaluation of their antioxidant, anti-genotoxic and anti-inflammatory activity. *Fitoterapia* 2011;82: 546-556.
175. Badam L. In vitro antiviral activity of indigenous glycyrrhizin, licorice and glycyrrhizic acid (Sigma) on Japanese encephalitis virus. *J Commun Dis* 1997;29: 91-9.
176. Mitscher LA, Park YH, Clark D, Beal JL. Antimicrobial Agents From Higher Plants. Antimicrobial Isoflavanoids and Related Substances From *Glycyrrhiza glabra* L. var. *typica*. *Journal of Natural Products* 1980;43: 259-269.
177. Fukai T, Satoh K, Nomura T, Sakagami H. Preliminary evaluation of antinephritis and radical scavenging activities of glabridin from *Glycyrrhiza glabra*. *Fitoterapia* 2003;74: 624-9.
178. Francischetti IM, Monteiro RQ, Guimaraes JA, Francischetti B. Identification of glycyrrhizin as a thrombin inhibitor. *Biochem Biophys Res Commun* 1997;235: 259-63.
179. Saeedi M, Morteza-Semnani K, Ghoreishi MR. The treatment of atopic dermatitis with licorice gel. *Journal of Dermatological Treatment* 2003;14: 153-157.
180. Shetty TK, Satav JG, Nair CK. Protection of DNA and microsomal membranes in vitro by *Glycyrrhiza glabra* L. against gamma irradiation. *Phytother. Res* 2002;16: 576-578.

181. Dhingra D, Parle M, Kulkarni SK. Comparative brain cholinesterase-inhibiting activity of Glycyrrhiza glabra, Myristica fragrans, ascorbic acid, and metrifonate in mice. *J Med Food* 2006;9: 281-3.
182. Bafna PA, Balaraman R. Antioxidant activity of DHC-1, an herbal formulation, in experimentally-induced cardiac and renal damage. *Phytotherapy Research* 2005;19: 216-221.
183. Murshed S, Rokeya B, Nahar N, Khan AKA, Mosihuzzaman M, Banerjee A, Maiti S, Ali L. Hypoglycemic and hypolipidemic effect of Hemidesmus indicus root on diabetic model rats. *Diabetes Research* 2005;39: 15-24.
184. Mary NK, Achuthan CR, Babu BH, Padikkala J. In vitro antioxidant and antithrombotic activity of Hemidesmus indicus (L) R. Br. *Journal of Ethnopharmacology* 2003;87: 187-191.
185. Chidrawar VR, Patel NJ, Ushir YV, Patel KN. Pre-clinical evolutionary study of Hemidesmus indicus Linn as minor tranquilizer. *J. Res. Educ. Indian Med* 2009.
186. Lakshman K, Shivaprasad HN, Jaiprakash B, Mohan S. Anti-Inflammatory and Antipyretic activities of Hemidesmus indicus root extract. *African Journal. Traditional* 2006;3: 90-94.
187. Anoop A, Jegadeesan M. Biochemical studies on the anti-ulcerogenic potential of Hemidesmus indicus R. Br. var. indicus. *Journal of Ethnopharmacology* 2003;84: 149-156.
188. Khandelwal VKM, Balaraman R, Ondrejčáková M, Pancza D, Ravingerová T. Effect of Hemidesmus indicus on ischemia-reperfusion injury in the isolated rat heart. *Pharmaceutical Biology* 2010;48: 611-614.
189. Khandelwal VKM, Balaraman R, Pancza D, Ravingerová T. Hemidesmus indicus and Hibiscus rosa-sinensis Affect Ischemia Reperfusion Injury in Isolated Rat Hearts. *Evidence-Based Complementary and Alternative Medicine* 2011;2011.
190. Baheti JR, Goyal RK, Shah GB. Hepatoprotective activity of Hemideomus indicus R. Br. in rats. *Indian journal of experimental biology* 2006;44: 399.
191. Kotnis MS, Patel P, Menon SN, Sane RT. Renoprotective effect of Hemidesmus indicus, a herbal drug used in gentamicin induced renal toxicity. *Nephrology* 2004;9: 142-152.
192. Ananthi R, Chandra N, Santhiya ST, Ramesh A. Genotoxic and antigenotoxic effects of Hemidesmus indicus R. Br. root extract in cultured lymphocytes. *Journal of Ethnopharmacology* 2010;127: 558-560.
193. Shetty TK, Satav JG, Nair CKK. Radiation protection of DNA and membrane in vitro by extract of Hemidesmus indicus. *Phytotherapy Research* 2005;19: 387-390.

194. Alam MI, Auddy B, Gomes A. Viper venom neutralization by Indian medicinal plant (*Hemidesmus indicus* and *Pluchea indica*) root extracts. *Phytotherapy Research* 1996;10: 58-61.
195. Joseph B, Sujatha S, Anusha JR. Bioactivity of *Hemidesmus indicus* (L.) on Human Pathogenic Bacteria and *Culex quinquefasciatus* (Diptera: Culicidae). *Research Journal of Medicinal Plant* 2011;5: 613-620.
196. Bhaskar A, Vidhya VG, Ramya M. Hypoglycemic effect of *Mucuna pruriens* seed extract on normal and streptozotocin-diabetic rats. *Fitoterapia* 2008;79: 539-543.
197. Ratnawati H, Widowati W. Anticholesterol Activity of Velvet Bean (*Mucuna pruriens* L.) towards Hypercholesterolemic Rats. *Sains Malaysiana* 2011;40: 317-321.
198. Tripathi YB, Upadhyay AK. Effect of the alcohol extract of the seeds of *Mucuna pruriens* on free radicals and oxidative stress in albino rats. *Phytotherapy Research* 2002;16: 534-538.
199. Suresh S, Prithviraj E, Prakash S. Dose- and time-dependent effects of ethanolic extract of *Mucuna pruriens* Linn. seed on sexual behaviour of normal male rats. *Journal of Ethnopharmacology* 2009;122: 497-501.
200. Manyam BV, Dhanasekaran M, Hare TA. Neuroprotective effects of the antiparkinson drug *Mucuna pruriens*. *Phytotherapy Research* 2004;18: 706-712.
201. Tan NH, Fung SY, Sim SM, Marinello E, Guerranti R, Aguiyi JC. The protective effect of *Mucuna pruriens* seeds against snake venom poisoning. *Journal of Ethnopharmacology* 2009;123: 356-358.
202. Vaidya RA, Aloorkar SD, Sheth AR, Pandya SK. Activity of bromoergocryptine, *Mucuna pruriens* and L-dopa in the control of hyperprolactinaemia. *Neurol India* 1978;26: 179-82.
203. Katzenschlager R, Evans A, Manson A, Patsalos PN, Ratnaraj N, Watt H, Timmermann L, Van der Giessen R, Lees AJ. *Mucuna pruriens* in Parkinson's disease: a double blind clinical and pharmacological study. *Journal of Neurology, Neurosurgery & Psychiatry* 2004;75: 1672-1677.
204. Infante ME, Perez AM, Simao MR, Manda F, Baquete EF, Fernandes AM, Cliff JL. Outbreak of acute toxic psychosis attributed to *Mucuna pruriens*. *Lancet* 1990;336: 1129.
205. Shelley WB, Arthur RP. Studies on cowhage (*Mucuna pruriens*) and its pruritogenic proteinase, mucunain. *AMA Arch Derm* 1955;72: 399-406.
206. Subeki S, Matsuura H, Takahashi K, Yamasaki M, Yamato O, Maede Y, Katakura K, Kobayashi S, Trimurningsih C, Chairul C. Anti-babesial and anti-

- plasmodial compounds from *Phyllanthus niruri*. *Journal of Natural Products* 2005;68: 537-539.
207. Murugaiyah V, Chan K-L. Mechanisms of antihyperuricemic effect of *Phyllanthus niruri* and its lignan constituents. *Journal of Ethnopharmacology* 2009;124: 233-239.
208. Santos Filho AR, Yunes RA, Calixto JB. Analysis of the mechanisms underlying the antinociceptive effect of the extracts of plants from the genus *Phyllanthus*. *Gen Pharmacol* 1995;26: 1499-1506.
209. Naik A, Juvekar A. Effects of alkaloidal extract of *Phyllanthus niruri* on HIV replication. *Indian Journal of Medical Sciences* 2003;57: 387-93.
210. Chatterjee M, Sarkar K, Sil PC. Herbal (*Phyllanthus niruri*) protein isolate protects liver from nimesulide induced oxidative stress. *Pathophysiology* 2006;13: 95-102.
211. Iizuka T, Moriyama H, Nagai M. Vasorelaxant Effects of Methyl Brevifolincarboxylate from the Leaves of *Phyllanthus niruri*. *Biological & Pharmaceutical Bulletin* 2006;29: 177-179.
212. Khanna AK, Rizvi F, Chander R. Lipid lowering activity of *Phyllanthus niruri* in hyperlipemic rats. *Journal of Ethnopharmacology* 2002;82: 19-22.
213. Iizuka T, Nagai M, Taniguchi A, Moriyama H, Hoshi K. Inhibitory Effects of Methyl Brevifolincarboxylate Isolated from *Phyllanthus niruri* L. on Platelet Aggregation. *Biological & Pharmaceutical Bulletin* 2007;30: 382-384.
214. Sampaio F. Effect of extract of *Phyllanthus niruri* on crystal deposition in experimental urolithiasis. *Int Braz J Urol* 2007;33: 114-5.
215. Venkateswaran PS, Millman I, Blumberg BS. Effects of an Extract from *Phyllanthus niruri* on Hepatitis B and Woodchuck Hepatitis Viruses: In vitro and in vivo Studies. *Proceedings of the National Academy of Sciences* 1987;84: 274-278.
216. De Melo AC, Perec CJ, Rubio MC. Acetylcholine-like activity in the fruit of the black nightshade (Solanaceae). *Acta Physiol Lat Am* 1978;28: 19-26.
217. Zakaria ZA, Gopalan HK, Zainal H, Pojan NHM, Morsid NA, Aris A, Sulaiman MR. Antinociceptive, anti-inflammatory and antipyretic effects of *Solanum nigrum* chloroform extract in animal models. *YAKUGAKU ZASSHI* 2006;126: 1171-1178
218. Heo KS, Lim KT. Antioxidative effects of glycoprotein Isolated from *Solanum nigrum* L. *Journal of Medicinal Food* 2004;7: 349-357.
219. Huang H-C, Syu K-Y, Lin J-K. Chemical Composition of *Solanum nigrum* Linn Extract and Induction of Autophagy by Leaf Water Extract and Its Major Flavonoids in AU565 Breast Cancer Cells. *Journal of Agricultural and Food Chemistry* 2010;58: 8699-8708.

220. Jainu M, Devi CSS. Antiulcerogenic and ulcer healing effects of *Solanum nigrum* (L.) on experimental ulcer models: Possible mechanism for the inhibition of acid formation. *Journal of Ethnopharmacology* 2006;104: 156-163.
221. Javed T, Ashfaq U, Riaz S, Rehman S, Riazuddin S. In-vitro antiviral activity of *Solanum nigrum* against Hepatitis C Virus. *Virology Journal* 2011;8: 26.
222. Raju K, Anbuganapathi G, Gokulakrishnan V, Rajkapoor B, Jayakar B, Manian S. Effect of dried fruits of *Solanum nigrum* LINN against CCl 4-Induced hepatic damage in rats. *Biological & Pharmaceutical Bulletin* 2003;26: 1618-1619.
223. Prashanth Kumar V, Shashidhara S, Kumar MM, Sridhara BY. Cytoprotective role of *Solanum nigrum* against gentamicin-induced kidney cell (Vero cells) damage in vitro. *Fitoterapia* 2001;72: 481-6.
224. Lee SJ, Ko JH, Lim K, Lim KT. 150 kDa glycoprotein isolated from *Solanum nigrum* Linne enhances activities of detoxicant enzymes and lowers plasmic cholesterol in mouse. *Pharmacol. Res* 2005;51: 399-408.
225. Ye JR. The hypotensive effect of *Solanum nigrum*. *Zhong Yao Tong Bao* 1984;9: 35-6.
226. Wannang NN, Anuka JA, Kwanashie HO, Gyang SS, Auta A. Anti-seizure activity of the aqueous leaf extract of *Solanum nigrum* linn (solanaceae) in experimental animals. *African Health Sciences* 2008;8: 74.
227. Friedman M, Rayburn JR, Bantle JA. Structural relationships and development toxicity of *Solanum* alkaloids in the frog embryo teratogenesis assay-Xenopus. *Journal of Agricultural and Food Chemistry* 1992;40: 1617-1624.
228. Banerjee S, Das S. Anticarcinogenic Effects of an Aqueous Infusion of Cloves on Skin Carcinogenesis. *Asian Pacific J. Cancer Prev* 2005;6: 304-308.
229. Shyamala MP, Venukumar MR, Latha MS. Antioxidant potential of the *Syzygium aromaticum* (gaertn.) Linn.(cloves) in rats fed with high fat diet. *Indian Journal of Pharmacology* 2003;35: 99-103.
230. Kurokawa M, Hozumi T, Basnet P, Nakano M, Kadota S, Namba T, Kawana T, Shiraki K. Purification and Characterization of Eugeniiin as an Anti-herpesvirus Compound from *Geum japonicum* and *Syzygium aromaticum*. *Journal of Pharmacology and Experimental Therapeutics* 1998;284: 728-735.
231. Lee JI, Lee HS, Jun WJ, Yu KW, Shin DH, Hong BS, Cho HY, Yang HC. Purification and Characterization of Antithrombotics from *Syzygium aromaticum* (L.) MERR. & PERRY. *Biological & Pharmaceutical Bulletin* 2001;24: 181-187.
232. Shiraki K, Yukawa T, Kurokawa M, Kageyama S. Cytomegalovirus infection and its possible treatment with herbal medicines. *Nippon Rinsho* 1998;56: 156-60.

233. Taguchi Y, Ishibashi H, Takizawa T, Inoue S, Yamaguchi H, Abe S. Protection of oral or intestinal candidiasis in mice by oral or intragastric administration of herbal food, clove (*Syzygium aromaticum*). *Jpn. J. Med. Mycol.* 2005;46: 27-33.
234. Tajuddin SA, Latif A, Qasmi IA. Effect of 50% ethanolic extract of *Syzygium aromaticum* (L.) Merr. & Perry.(clove) on sexual behaviour of normal male rats. *BMC Complementary and Alternative Medicine* 2004;4: 17-23.
235. Banerjee S, Panda CK, Das S. Clove (*Syzygium aromaticum* L.), a potential chemopreventive agent for lung cancer. *Carcinogenesis* 2006;27: 1645.
236. Kim HM, Lee EH, Hong SH, Song HJ, Shin MK, Kim SH, Shin TY. Effect of *Syzygium aromaticum* extract on immediate hypersensitivity in rats. *Journal of Ethnopharmacology* 1998;60: 125-131.
237. Prashar A, Locke IC, Evans CS. Cytotoxicity of clove (*Syzygium aromaticum*) oil and its major components to human skin cells. *Cell Proliferation* 2006;39: 241-248.
238. Balaraman R, Bafna PA, Kolhapure SA. Antioxidant activity of DHC-1 a herbal formulation. *Journal of Ethnopharmacology* 2004;94: 135-141.
239. Saleem A, Husheem M, Harkonen P, Pihlaja K. Inhibition of cancer cell growth by crude extract and the phenolics of *Terminalia chebula* retz. fruit. *Journal of Ethnopharmacology* 2002;81: 327-36.
240. Hamada S, Kataoka T, Woo JT, Yamada A, Yoshida T, Nishimura T, Otake N, Nagai K. Immunosuppressive effects of gallic acid and chebulagic acid on CTL-mediated cytotoxicity. *Biol Pharm Bull* 1997;20: 1017-9.
241. Gopi KS, Reddy AG, Jyothi K, Kumar BA. Acetaminophen-induced Hepato-and Nephrotoxicity and Amelioration by Silymarin and *Terminalia chebula* in Rats. *Toxicology International* 2010;17: 64.
242. Suchalatha S, Shyamala Devi CS. Protective effect of *Terminalia chebula* against experimental myocardial injury induced by isoproterenol. *Indian J Exp Biol* 2004;42: 174-8.
243. Gandhi NM, Nair CKK. Radiation protection by *Terminalia chebula*: Some mechanistic aspects. *Molecular and Cellular Biochemistry* 2005;277: 43-48.
244. Rao N, Nammi S. Antidiabetic and renoprotective effects of the chloroform extract of *Terminalia chebula* Retz. seeds in streptozotocin-induced diabetic rats. *BMC Complementary and Alternative Medicine* 2006;6: 17.
245. Ahn MJ, Kim CY, Lee JS, Kim TG, Kim SH, Lee CK, Lee BB, Shin CG, Huh H, Kim J. Inhibition of HIV-1 Integrase by Galloyl Glucoses from *Terminalia chebula* and Flavonol Glycoside Gallates from *Euphorbia pekinensis*. *Planta Medica* 2002;68: 457-459.

246. Miglani BD, Sen P, Sanyal RK. Purgative action of an oil obtained from *Terminalia chebula*. *Indian J Med Res* 1971;59: 281-3.
247. Senthilkumar GP. Evaluation of Antioxidant Potential of *Terminalia chebula* Fruits Studied in Streptozotocin-Induced Diabetic Rats. *Pharmaceutical Biology* 2007;45: 511-518.
248. Manosroi A, Jantrawut P, Akihisa T, Manosroi W, Manosroi J. In vitro anti-aging activities of *Terminalia chebula* gall extract. *Pharmaceutical Biology* 2010;48: 469-481.
249. Malekzadeh F, Ehsanifar H, Shahamat M, Levin M, Colwell RR. Antibacterial activity of black myrobalan(*Terminalia chebula* Retz) against *Helicobacter pylori*. *International Journal of Antimicrobial Agents* 2001;18: 85-88.
250. Khan KH, Jain SK. Regular intake of *Terminalia chebula* can reduce the risk of getting typhoid fever. *Advanced Biotech* 2009;8: 10-15.
251. Jagtap AG, Karkera SG. Potential of the aqueous extract of *Terminalia chebula* as an anticaries agent. *Journal of Ethnopharmacology* 1999;68: 299-306.
252. Suguna L, Singh S, Sivakumar P, Sampath P, Chandrakasan G. Influence of *Terminalia chebula* on dermal wound healing in rats. *Phytotherapy Research* 2002;16: 227-231.
253. Tamhane M, Thorat S, Rege N, Dahanukar S. Effect of oral administration of *Terminalia chebula* on gastric emptying: an experimental study. *Journal of Postgraduate Medicine* 1997;43: 12-13.
254. Saravanan S, Srikumar R, Manikandan S, Jeya Parthasarathy N, Sheela Devi R. Hypolipidemic effect of Triphala in experimentally induced hypercholesteremic rats. *YAKUGAKU ZASSHI* 2007;127: 385-388.
255. Srikumar R, Jeya Parthasarathy N, Sheela Devi R. Immunomodulatory activity of Triphala on neutrophil functions. *Biological & Pharmaceutical Bulletin* 2005;28: 1398-1403.
256. Kaur S, Arora S, Kaur K, Kumar S. The in vitro antimutagenic activity of Triphala--an Indian herbal drug. *Food Chem Toxicol* 2002;40: 527-34.
257. Srikumar R, Parthasarathy NJ, Shankar EM, Manikandan S, Vijayakumar R, Thangaraj R, Vijayananth K, Sheeladevi R, Rao UA. Evaluation of the growth inhibitory activities of triphala against common bacterial isolates from HIV infected patients. *Phytother Res* 2007.
258. Sabu MC, Kuttan R. Anti-diabetic activity of medicinal plants and its relationship with their antioxidant property. *Journal of Ethnopharmacology* 2002;81: 155-160.

259. Arseculeratne SN, Gunatilaka AA, Panabokke RG. Studies of medicinal plants of Sri Lanka. Part 14: Toxicity of some traditional medicinal herbs. *Journal of Ethnopharmacology* 1985;13: 323-35.
260. Purandare H, Supe A, Statistics A, Comments R, Downloaded PDF. Immunomodulatory role of *Tinospora cordifolia* as an adjuvant in surgical treatment of diabetic foot ulcers: A prospective randomized controlled study. *Indian J Med Sci* 2007;61: 347-55.
261. Badar VA, Thawani VR, Wakode PT, Shrivastava MP, Gharpure KJ, Hingorani LL, Khiyani RM. Efficacy of *Tinospora cordifolia* in allergic rhinitis. *Journal of Ethnopharmacology* 2005;96: 445-449.
262. Sarma DNK, Khosa RL, Chansouria JPN, Sahai M. Antiulcer activity of *Tinospora cordifolia* Miers and *Centella asiatica* Linn extracts. *Phytother Res* 1995;9: 589-590.
263. Rao PR, Kumar VK, Viswanath RK, Subbaraju GV. Cardioprotective Activity of Alcoholic Extract of *Tinospora cordifolia* in Ischemia-Reperfusion Induced Myocardial Infarction in Rats. *Biological & Pharmaceutical Bulletin* 2005;28: 2319-2322.
264. Singh RP, Banerjee S, Kumar PV, Raveesha KA, Rao AR. *Tinospora cordifolia* induces enzymes of carcinogen/drug metabolism and antioxidant system, and inhibits lipid peroxidation in mice. *Phytomedicine* 2006;13: 74-84.
265. Bishayi B, Roychowdhury S, Ghosh S, Sengupta M. Hepatoprotective and immunomodulatory properties of *Tinospora cordifolia* in CCl₄ intoxicated mature albino rats. *The Journal of Toxicological Sciences* 2002;27: 139-146.
266. Singh RK. *Tinospora cordifolia* as an adjuvant drug in the treatment of hyper-reactive malarious splenomegaly-case reports. *Journal of Vector Borne Diseases* 2005;42: 36-38.
267. Prince PSM, Menon VP, Gunasekaran G. Hypolipidaemic action of *Tinospora cordifolia* roots in alloxan diabetic rats. *Journal of Ethnopharmacology* 1998;64: 53-57.
268. Rawal AK, Muddeshwar MG, Biswas SK. *Rubia cordifolia*, *Fagonia cretica* linn and *Tinospora cordifolia* exert neuroprotection by modulating the antioxidant system in rat hippocampal slices subjected to oxygen glucose deprivation. *feedback* 2005.
269. Rege N, Bapat RD, Koti R, Desai NK, Dahanukar S. Immunotherapy with *Tinospora cordifolia*: a new lead in the management of obstructive jaundice. *Indian J Gastroenterol* 1993;12: 5-8.

270. Prince PSM, Menon VP. Antioxidant action of *Tinospora cordifolia* root extract in alloxan diabetic rats. *Phytother. Res* 2001;15: 213-218.
271. Goel HC, Prem Kumar I, Rana SV. Free radical scavenging and metal chelation by *Tinospora cordifolia*, a possible role in radioprotection. *Indian J Exp Biol* 2002;40: 727-34.
272. Gupta RS, Sharma A. Antifertility effect of *Tinospora cordifolia* (Willd.) stem extract in male rats. *Indian J Exp Biol* 2003;41: 885-9.
273. Bafna PA, Balaraman R. Anti-ulcer and anti-oxidant activity of pepticare, a herbomineral formulation. *Phytomedicine* 2005;12: 264-70.
274. Bhattacharya SK, Bhattacharya A, Sairam K, Ghosal S. Anxiolytic-antidepressant activity of *Withania somnifera* glycowithanolides: an experimental study. *Phytomedicine* 2000;7: 463-9.
275. Girish KS, Machiah KD, Ushanandini S, Harish Kumar K, Nagaraju S, Govindappa M, Vedavathi M, Kemparaju K. Antimicrobial properties of a non-toxic glycoprotein (WSG) from *Withania somnifera* (Ashwagandha). *J Basic Microbiol* 2006;46: 365-74.
276. Dikasso D, Makonnen E, Debella A, Abebe D, Urga K, Makonnen W, Melaku D, Kassa M, Guta M. Anti-malarial activity of *withania somnifera* L. Dunal extracts in mice. *Ethiop Med J* 2006;44: 279-85.
277. Senthil V, Ramadevi S, Venkatakrishnan V, Giridharan P, Lakshmi BS, Vishwakarma RA, Balakrishnan A. Withanolide induces apoptosis in HL-60 leukemia cells via mitochondria mediated cytochrome c release and caspase activation. *Chemico-Biological Interactions* 2007;167: 19-30.
278. Sumantran VN, Kulkarni A, Boddul S, Chinchwade T, Koppikar SJ, Harsulkar A, Patwardhan B, Chopra A, Wagh UV. Chondroprotective potential of root extracts of *Withania somnifera* in osteoarthritis. *Journal of Biosciences* 2007;32: 299-307.
279. Hamza A, Amin A, Daoud S. The protective effect of a purified extract of *Withania somnifera* against doxorubicin-induced cardiac toxicity in rats. *Cell Biol Toxicol* 2007.
280. Davis L, Kuttan G. Immunomodulatory activity of *Withania somnifera*. *Journal of Ethnopharmacology* 2000;71: 193-200.
281. Sankar SR, Manivasagam T, Krishnamurti A, Ramanathan M. The neuroprotective effect of *Withania somnifera* root extract in MPTP-intoxicated mice: An analysis of behavioral and biochemical variables. *Cell Mol Biol Lett* 2007.
282. Jayaprakasam B, Nair MG. Cyclooxygenase-2 enzyme inhibitory withanolides from *Withania somnifera* leaves. *Tetrahedron* 2003;59: 841-849.

283. Bhattacharya SK, Kumar A, Ghosal S. Effects of glycowithanolides from *Withania somnifera* on an animal model of Alzheimer's disease and perturbed central cholinergic markers of cognition in rats. *PTR. Phytotherapy research* 1995;9: 110-113.
284. Sharada AC, Emerson Solomon F, Uma Devi P. Toxicity of *Withania somnifera* root extract in rats and mice. *International journal of pharmacognosy* 1993;31: 205-212.
285. Ansari MN, Bhandari U, Pillai KK. Ethanolic *Zingiber officinale* R. extract pretreatment alleviates isoproterenol-induced oxidative myocardial necrosis in rats. *Indian J Exp Biol* 2006;44: 892-7.
286. Vishwakarma SL, Pal SC, Kasture VS, Kasture SB. Anxiolytic and antiemetic activity of *Zingiber officinale*. *Phytother Res* 2002;16: 621-6.
287. Al-Amin ZM, Thomson M, Al-Qattan KK, Peltonen-Shalaby R, Ali M. Anti-diabetic and hypolipidaemic properties of ginger (*Zingiber officinale*) in streptozotocin-induced diabetic rats. *British Journal of Nutrition* 2007;96: 660-666.
288. Sang S, Hong J, Wu H, Liu J, Yang CS, Pan M-H, Badmaev V, Ho C-T. Increased Growth Inhibitory Effects on Human Cancer Cells and Anti-inflammatory Potency of Shogaols from *Zingiber officinale* Relative to Gingerols. *Journal of Agricultural and Food Chemistry* 2009;57: 10645-10650.
289. Thomson M, Al-Qattan KK, Al-Sawan SM, Alnaqeeb MA, Khan I, Ali M. The use of ginger (*Zingiber officinale* Rosc.) as a potential anti-inflammatory and antithrombotic agent. *Prostaglandins Leukot Essent Fatty Acids* 2002;67: 475-8.
290. Goyal RK, Kadnur SV. Beneficial effects of *Zingiber officinale* on goldthioglucose induced obesity. *Fitoterapia* 2006;77: 160-163.
291. Masuda Y. Antioxidant properties of gingerol related compounds from ginger. *BioFactors* 2004;21: 293-296.
292. Fuhrman B, Rosenblat M, Hayek T, Coleman R, Aviram M. Ginger extract consumption reduces plasma cholesterol, inhibits LDL oxidation and attenuates development of atherosclerosis in atherosclerotic, apolipoprotein e-deficient mice. *Journal of Nutrition* 2000;130: 1124-1131.
293. Haksar A, Sharma A, Chawla R, Kumar R, Arora R, Singh S, Prasad J, Gupta M, Tripathi RP, Arora MP. *Zingiber officinale* exhibits behavioral radioprotection against radiation-induced CTA in a gender-specific manner. *Pharmacol Biochem Behav* 2006;84: 179-88.
294. Ghayur MN, Gilani AH. Ginger lowers blood pressure through blockade of voltage-dependent calcium channels. *Journal of Cardiovascular Pharmacology* 2005;45: 74-80.

295. Ajith TA, Nivitha V, Usha S. Zingiber officinale Roscoe alone and in combination with a-tocopherol protect the kidney against cisplatin-induced acute renal failure. *Food and Chemical Toxicology* 2007;45: 921-927.
296. Amin A, Hamza AEA. Effects of Roselle and Ginger on cisplatin-induced reproductive toxicity in rats. *Asian J Androl* 2006;8: 607-612.
297. Siddaraju MN, Dharmesh SM. Research Article Inhibition of gastric H, K-ATPase and Helicobacter pylori growth by phenolic antioxidants of Zingiber officinale. *Mol. Nutr. Food Res* 2007;51: 324-332.
298. Yemitan OK, Izegbu MC. Protective effects of Zingiber officinale (Zingiberaceae) against carbon tetrachloride and acetaminophen-induced hepatotoxicity in rats. *Phytother. Res* 2006;20: 997-1002.
299. Kamtchouing P, Mbongue Fandio GY, Dimo T, Jatsa HB. Evaluation of androgenic activity of Zingiber officinale and Pentadiplandra brazzeana in male rats. *Asian J Androl* 2002;4: 299-301.
300. Etoh H, Kondoh T, Noda R, Singh IP, Sekiwa Y, Morimitsu K, Kubota K. Shogaols from Zingiber officinale as Promising Antifouling Agents. *Bioscience, Biotechnology, and Biochemistry* 2002;66: 1748-1750.
301. Ghayur MN, Gilani AH. Species differences in the prokinetic effects of ginger. *International Journal of Food Sciences and Nutrition* 2006;57: 65-73.
302. Kim EC, Min JK, Kim TY, Lee SJ, Yang HO, Han S, Kim YM, Kwon YG. [6]-Gingerol, a pungent ingredient of ginger, inhibits angiogenesis in vitro and in vivo. *Biochem Biophys Res Commun* 2005;335: 300-8.
303. Sekiya K. Enhancement of insulin sensitivity in adipocytes by ginger. *BioFactors* 2004;22: 153-156.
304. Adewunmi CO, Oguntimein BO, Furu P. Molluscicidal and antischistosomal activities of Zingiber officinale. *Planta Med* 1990;56: 374-6.
305. Joshi H, Parle M. Zingiber officinale: Evaluation of its Nootropic effect in mice. *African Journal. Traditional* 2005;3: 64-74.
306. [Ghayur MN, Gilani AH. Pharmacological Basis for the Medicinal Use of Ginger in Gastrointestinal Disorders. *Digestive Diseases and Sciences* 2005;50: 1889-1897.
307. Mustafa T, Srivastava KC. Ginger (Zingiber officinale) in migraine headache. *Journal of Ethnopharmacology* 1990;29: 267-73.
308. Willetts KE, Ekangaki A, Eden JA. Effect of a ginger extract on pregnancy-induced nausea: A randomised controlled trial. *The Australian and New Zealand Journal of Obstetrics and Gynaecology* 2003;43: 139-144.
309. Weidner MS, Sigwart K. Investigation of the teratogenic potential of a zingiber officinale extract in the rat. *Reprod Toxicol* 2001;15: 75-80.